

## ON THE TOPOLOGY OF RIEMANN SPACES OF QUASI-CONSTANT CURVATURE

Sorin Dragomir and Renata Grimaldi

**Abstract.** We compute Betti numbers of certain Riemann spaces of  $q$ -constant curvature such as conformally flat hypersurfaces of elliptic-space-forms, subprojective spaces and para-Sasakian manifolds.

### 1. Harmonic forms on manifolds of quasi-constant curvature

Let  $(M^n, g_{ij})$  be an  $n$ -dimensional space and  $V$  a unit vector field on  $M^n$ , with (local) components  $v^i$ . Let  $v_i = g_{ij}v^j$ ; if  $(M^n, g_{ij})$  is conformally flat and for some real valued smooth functions  $a, b \in C^\infty(M^n)$  the curvature of  $(M^n, g_{ij})$  is expressed by:

$$(1.1) \quad R_{kji}^h = a\{\delta_k^h g_{ji} - \delta_j^h g_{ki}\} \\ + b\{(\delta_k^h v_j - \delta_j^h v_k)v_i + (v_k g_{ji} - v_j g_{ki})v^h\}$$

then  $(M^n, g_{ij})$  is said to be a Riemann space of  $q$ -constant curvature, cf. [7], [8]. We suppose throughout that  $b \neq 0$ , (for  $b = 0$ ,  $M^n$  falls into nothing but a real space-form). Further contraction of indices in (1.1) gives the Ricci form:

$$(1.2) \quad R_{ji} = \{(n-1)a + b\}g_{ji} + b(n-2)v_j v_i.$$

If  $\alpha_{i_1 \dots i_p}, \beta_{j_1 \dots j_p}$  are two  $p$ -forms on  $M^n$ , we put as usual

$$\langle \alpha, \beta \rangle = \alpha^{(i_1 \dots i_p)} \beta_{(i_1 \dots i_p)},$$

$|\alpha|^2 = \langle \alpha, \alpha \rangle$ . We proceed by establishing the following:

**THEOREM 1.** *Let  $(M^n, g_{ij})$  be a compact orientable Riemann space of  $q$ -constant curvature. The following relation holds:*

$$(1.3) \quad \int_{M^n} \{p[p!\{(n-p)a + b\}|\alpha|^2 + \\ + (p-1)!(n-2p)b|i(V)\alpha|^2] + \nabla_j \alpha_{i_1 \dots i_p} \nabla^j \alpha^{i_1 \dots i_p}\} * 1 = 0$$

for each harmonic  $p$ -form  $\alpha$  on  $M^n$ .

*Proof.* Let  $(p!)^{-1}\alpha_{i_1\dots i_p}dx^{i_1}\wedge\dots\wedge dx^{i_p}$  be a differential  $p$ -form on  $M^n$ . Note that:

$$(1.4) \quad v_i\alpha^{i i_2\dots i_p} = (i(V)\alpha)^{i_2\dots i_p},$$

where  $i(V)$  denotes the interior product with  $V$ . Taking into account (1.4), (1.2) one obtains:

$$(1.5) \quad R_{ji}\alpha^{j i_2\dots i_p}\alpha_{i_2\dots i_p}^i = p!\{(n-1)a+b\}|\alpha|^2 + (p-1)!(n-2)b|i(V)\alpha|^2.$$

Using  $R_{ijks} = g_{ih}R_{kshj}^h$  and (1.1) one derives:

$$(1.6) \quad R_{ijkh}\alpha^{j i_3\dots i_p}\alpha_{i_3\dots i_p}^{kh} = 2ap!|\alpha|^2 + 4b(p-1)!|i(V)\alpha|^2$$

By (3.2.9) in [6, p. 88], if  $\Delta\alpha = 0$  then

$$(1.7) \quad \int_{M^n} \{pF_p(\alpha) + \nabla_j\alpha_{i_1\dots i_p}\nabla^j\alpha^{i_1\dots i_p}\} * 1 = 0,$$

where the quadratic form  $F_p(\alpha)$  is given by (3.2.10) in [6]. Finally (1.5)–(1.6) furnish:

$$(1.8) \quad F_p(\alpha) = p!\{(n-p)a+b\}|\alpha|^2 + (p-1)!(n-2p)b|i(V)\alpha|^2$$

and (1.7) leads to our (1.3), QED.

Betti numbers of conformally flat (compact, orientable) Riemann spaces are known to vanish (cf. Th. 3.9.1. in [6, p. 118]) provided that the Ricci curvature is positive definite. Yet, if  $M^n$  is a space of  $q$ -constant curvature, by (1.2) one has  $R_{ji}v^jv^i = (n-1)(a+b)$ , i.e.  $R_{ji}$  is degenerate along the distribution generated by  $V$ , provided  $b = -a$ . As an application of Theorem 1 we get

**THEOREM 2.** *Let  $M^n$  be a compact orientable connected Riemann space of  $q$ -constant curvature,  $n > 2$ . If  $a = \text{const.} > 0$  and  $db = V(b)v$  then  $M^n$  has the homology type of  $S^1 \times S^{n-1}$ .*

*Proof.* By a result of Wang and Adati, i.e. Th. 2.3. in [8, p. 101], if  $a = \text{const.}$  and  $b_j = b_iv^iv_j$ ,  $b_i = \nabla_ib$ , then  $b = -a$  and  $V$  is parallel. Therefore, since  $a > 0$ , the coefficients in (1.8) are subject to  $(n-p)a+b > 0$ ,  $(n-2p)b \geq 0$  iff  $n/2 \leq p < n-1$ . If  $n$  is even, i.e.  $n = 2m$ , we have  $b_p(M^n) = 0$  for all  $m \leq p < 2m-1$ . By Poincaré duality one obtains also  $b_p(M^n) = 0$ ,  $1 < p \leq m$ . A similar argument for  $n$  odd shows that all Betti numbers of  $M^n$  vanish except for  $b_1(M^n) = b_{n-1}(M^n)$ . Since  $v_i$  is parallel, it is harmonic. Thus  $b_1(M^n) \geq 1$ . Let  $\beta$  be any other harmonic 1-form on  $M^n$ . By applying the Hodge operator one obtains a harmonic  $(n-1)$ -form  $*\beta$ . At this point we may use (1.8) to get

$$(1.9) \quad F_{n-1}(*\beta) = (n-1)!(a+b)|*\beta|^2 - (n-2)!(n-2)b|i(V)*\beta|^2$$

Since  $b = -a$  and  $*\beta$  is harmonic, (1.3) or (1.7) furnishes  $i(V) * \beta = 0$ , which by applying once more the Hodge operator gives  $v \wedge \beta = 0$  or  $\beta = fv$  for some everywhere non-vanishing  $f \in C^\infty(M^n)$ . Since  $\beta$  is harmonic it is closed, so that  $df \wedge v = 0$  or  $df = \lambda v$  for some  $\lambda \in C^\infty(M^n)$ . But  $\beta$  is coclosed, too, such that  $(df, \beta) = (f, \delta\beta) = 0$ , by (2.9.3) in [6, p. 74], i.e.  $df$  and  $\beta$  are orthogonal. Thus  $0 = (df, \beta) = \lambda f \text{vol}(M^n)$  yields  $\lambda = 0$ ; since  $M^n$  is connected one obtains  $f = \text{const.}$ , i.e.  $b_1(M^n) = 1$ . Consequently,  $M^n$  has the same Betti numbers as the product  $S^1 \times S^{n-1}$ , QED.

### 2. Conformally flat real hypersurfaces

Let  $M^{n+1}(k)$  be a real space-form, i.e. a Riemann space of constant sectional curvature  $k$ . Let  $M^n$  be a conformally flat hypersurface of  $M^{n+1}(k)$ . Let  $h$  be the second fundamental form of the given immersion of  $M^n$  in  $M^{n+1}(k)$ . By a classical result of J. A. Schouten,  $h = \alpha_0 g + \beta_0 v \otimes v$ , for some  $\alpha_0, \beta_0 \in C^\infty(M^n)$  and some unit tangent vector field  $V = \sharp v$ . Here  $\sharp$  means raising of indices with respect to the induced metric  $g$  on  $M^n$ . Then, B. Y. Chen and K. Yano, [4], have shown (via the Gauss equation, e.g. (2.6) in [3, p. 45]) that  $M^n$  is a space of  $q$ -constant curvature with  $a = k + \alpha_0^2$ ,  $b = \alpha_0 \beta_0$  and unit vector  $(V)$ . This allows us to apply Theorem 1 to get

**THEOREM 3.** *Let  $M^n$  be a conformally flat compact orientable real hypersurface of the real projective space  $IRP^{n+1}(k)$ ,  $k > 0$ . Then  $M^n$  is a real homology sphere, provided  $\alpha_0 \beta_0 \geq 0$ .*

*Proof.* Let  $\alpha$  be a harmonic  $p$ -form on  $M^n$ . By (1.8) we obtain  $\alpha = 0$  if  $(n - p)(k + \alpha_0^2) + \alpha_0 \beta_0 > 0$ , and  $(n - 2p)\alpha_0 \beta_0 \geq 0$ . Under the hypothesis of Theorem 3, it is sufficient to impose  $p \leq n/2$ . We finally use Poincaré duality to prove that all Betti numbers of  $M^n$  vanish, QED.

### 3. Subprojective spaces

Cf. T. Adati, [1], if  $M^n$ ,  $n \geq 3$ , is conformally flat and the tensor field:

$$(3.1) \quad L_{ji} = -\frac{1}{n-2} R_{ji} + \frac{R}{2(n-1)(n-2)} g_{ji}$$

is expressed by:

$$(3.2) \quad L_{ji} = R\rho g_{ji} + \rho_j \sigma_i,$$

where  $\rho_j = \nabla_j \rho$ ,  $\sigma_i = \nabla_i \sigma$  and  $\sigma$  is a function of  $\rho$ ,  $\rho \neq 0$ , then  $M^n$  is said to be a subprojective space. Here  $R$  denotes the scalar curvature of  $(M^n, g_{ij})$  cf. [8, p. 96],  $M^n$  follows to be a manifold of  $q$ -constant curvature with  $a = -2$ ,  $b = -K$ ,  $v_i = \gamma^{-1} \rho_i$ , where  $\gamma = |d\rho|$ . Also if  $\sigma = f(\rho)$ , for some  $C^2$ -smooth function  $f$  of one variable, then (3.2) gives  $K = \gamma^2 f'(\rho)$ . Using (1.3) we obtain:

**THEOREM 4.** *Let  $M^n, n \geq 3$  be a compact orientable subprojective space with  $\rho < 0$  and  $\sigma = f(\rho)$  for some  $C^2$ -smooth decreasing function  $f: (-\infty, 0) \rightarrow \mathbf{R}$ . Then  $M^n$  is a real homology sphere.*

*Proof.* Using (1.3) for the harmonic  $p$ -form  $\alpha$  on  $M^n$  we note that

$$2(n - p)\rho + \gamma^2 f'(\rho) < 0, \quad (2p - n)\gamma 2f'(\rho) \geq 0,$$

provided that  $p \geq n/2$ . As in Th. 3, by separately analysing the cases  $n = \text{odd}$  and  $n = \text{even}$ , and by Poincaré duality one obtains  $b_p(M^n) = 0$ , for all  $p$ , QED.

**4. Special para-Sasakian manifolds with vanishing  $D$ -concircular tensor**

Let  $(M^n, g_{ij})$  be a Riemann space,  $n \geq 3$ , and  $\xi^i$  a given unit vector field with  $\nabla_j \xi^i = \varepsilon(-\delta_j^i + \eta_j \xi^i)$ ,  $\eta_j = g_{ji} \xi^i$ ,  $\varepsilon = \pm 1$ . Then  $M^n$  carries a para-contact metric structure  $(\phi_j^i, \xi^i, \eta_j, g_{ij})$ ,  $\phi_j^i = \nabla_j \xi^i$ , [2]. On the other hand, cf. G. Chuman, [5], one may consider the  $D$ -concircular tensor:

$$(4.1) \quad U_{kji}^h = R_{kji}^h + \frac{R + 2(n - 1)}{(n - 1)(n - 2)} \{g_{ki} \delta_j^h - g_{ji} \delta_k^h\} - \\ - \frac{R + n(n - 1)}{(n - 1)(n - 2)} \{g_{ki} \eta_j \xi^h - g_{ji} \eta_k \xi^h + \eta_k \eta_i \delta_j^h - \eta_j \eta_i \delta_k^h\}.$$

Consequently, if  $U_{kji}^h = 0$  then the curvature of  $M^n$  is expressed by (1.1) with  $a = \frac{R + 2(n - 1)}{(n - 1)(n - 2)}$ ,  $b = -\frac{R + n(n - 1)}{(n - 1)(n - 2)}$ ,  $v_i = \eta_i$ . Using (1.3) one obtains:

**THEOREM 5.** *Let  $M^n$  be a compact orientable special para-Sasakian manifold with a vanishing  $D$ -concircular tensor. Then there exists  $R_0 > 0$  such that if  $R \geq R_0$  then  $b_p(M^n) = 0$ ,  $1 < p < n - 1$ .*

*Proof.* Let  $\alpha$  be a harmonic  $p$ -form on  $M^n$ . By (1.8) if  $(n - p)(R + 2(n - 1)) - (R + n(n - 1)) > 0$ ,  $(2p - n)(R + n(n - 1)) \geq 0$  then  $F_p(\alpha)$  is positive definite. Suppose  $R \geq 0$ . Define  $f_n(R) = n - \frac{R + n(n - 1)}{R + 2(n - 1)}$ . Then  $f_n$  is strictly increasing,  $f_n(0) = n/2$ ,  $\lim_{R \rightarrow +\infty} f_n(R) = n - 1$ . This makes clear the contents of the assumption on the scalar curvature in Theorem 5. Indeed, as  $R \rightarrow +\infty$ , i.e. if  $R \geq R_0$  for some constant  $R_0 > 0$ , the condition  $n/2 \leq p < n - 1$  is equivalent to  $n/2 \leq p < f_n(R)$ , QED.

Note that  $M^n$  in Theorem 5 is not necessarily conformally flat, but the form (1.1) is sufficient for our purpose; by [8, p. 98], the additional condition  $R = \text{const.}$  implies  $M^n$  is a space of  $q$ -constant curvature.

If  $M^n$  is a space of  $q$ -constant curvature with  $R = \text{const.} \neq 0$  and at the same time an  $S$ -manifold (i.e.  $\nabla_m \nabla_s R_{kji}^h = \nabla_s \nabla_m R_{kji}^h$ ) then  $b_1(M^n) \geq 1$ , since from Theorem 3.3. in [8, p. 102] it follows that  $v_i$  is parallel, and thus harmonic. Nevertheless, this situation may not be used in our Theorem 5. Indeed, by a result

of Wang, [7], a space of  $q$ -constant curvature is an  $S$ -manifold iff  $a + b = 0$  (while any special para-Sasakian manifold with  $U_{kji}^h = 0$  has  $a + b = -1$ ).

## REFERENCES

- [1] T. Adati, *On subprojective spaces*, Tohoku Math. J. (2) **3-2** (1951), 159–173.
- [2] T. Adati, G. Chuman, *Special para-Sasakian manifolds and concircular transformations*, TRU Math. (1) **20** (1984), 111–123.
- [3] B. Y. Chen, *Geometry of submanifolds*, M. Dekker, New York, 1973.
- [4] B. Y. Chen, K. Yano, *Hypersurfaces of conformally flat spacec*, Tensor, N.S. **26** (1972), 318–322.
- [5] G. Chuman, *D-conformal changes in para-Sasakian manifolds*, Tensor, N.S. **39** (1982), 117–123.
- [6] S. Goldberg, *Curvature and Homology*, Academic Press, New York, 1962.
- [7] Y. Wang, *On some roperties of Riemann spaces of quasi-constant curvature*, Tensor, N.S. **35** (1981), 173–176.
- [8] Y. Wang, T. Adati, *Manifolds of quasi-constant curvature*, TRU Math. (1) **21** (1985), 95–103.

Sorin Dragomir  
Università degli Studi di Bari  
Dipartimento di Matematica  
Via G. Fortunato  
Bari, Italia

(Received 22 09 1988)

Renata Grimaldi  
Università di Palermo  
Dipartimento di Matematica ed Applicazioni  
Via Archirafi  
Palermo, Italia