

## AN ESTIMATE FOR COEFFICIENTS OF POLYNOMIALS IN $L^2$ NORM. II

G. V. Milovanović and L. Z. Rančić

*Dedicated to the memory of Professor S. Aljančić*

**Abstract.** Let  $\mathcal{P}_n$  be the class of algebraic polynomials  $P(x) = \sum_{k=0}^n a_k x^k$  of degree at most  $n$  and  $\|P\|_{d\sigma} = (\int_{\mathbb{R}} |P(x)|^2 d\sigma(x))^{1/2}$ , where  $d\sigma(x)$  is a nonnegative measure on  $\mathbb{R}$ . We determine the best constant in the inequality  $|a_k| \leq C_{n,k}(d\sigma)\|P\|_{d\sigma}$ , for  $k = 0, 1, \dots, n$ , when  $P \in \mathcal{P}_n$  and such that  $P(\xi_k) = 0$ ,  $k = 1, \dots, m$ . The cases  $C_{n,n}(d\sigma)$  and  $C_{n,n-1}(d\sigma)$  were studied by Milovanović and Guessab [6]. In particular, we consider the case when the measure  $d\sigma(x)$  corresponds to generalized Laguerre orthogonal polynomials on the real line.

### 1. Introduction

Let  $\mathcal{P}_n$  be the class of algebraic polynomials  $P(x) = \sum_{k=0}^n a_k x^k$  of degree at most  $n$ . The first inequality of the form  $|a_k| \leq C_{n,k}\|P\|$  was given by Markov [3]. Namely, if  $\|P\| = \|P\|_{\infty} = \max_{x \in [-1,1]} |P(x)|$  and  $T_n(x) = \sum_{\nu=0}^n t_{n,\nu} x^{\nu}$  denotes the  $n$ -th Chebyshev polynomial of the first kind, then Markov proved that

$$|a_k| \leq \begin{cases} |t_{n,k}| \cdot \|P\|_{\infty} & \text{if } n-k \text{ is even,} \\ |t_{n-1,k}| \cdot \|P\|_{\infty} & \text{if } n-k \text{ is odd.} \end{cases} \quad (1.1)$$

For  $k = n$  (1.1) reduces to the well-known Chebyshev inequality

$$|a_n| \leq 2^{n-1} \|P\|_{\infty}. \quad (1.2)$$

---

AMS Subject Classification (1991): Primary 26 C 05, 26 D 05, 33 C 45, 41 A 44.

\*Research partly supported by Science Fund of Serbia under grant 0401F.

Using a restriction on the polynomial class like  $P(1) = 0$  or  $P(-1) = 0$ , Schur [8] found the following improvement of (1.2)

$$|a_n| \leq 2^{n-1} \left( \cos \frac{\pi}{4n} \right)^{2n} \|P\|_{\infty}.$$

This result was extended by Rahman and Schmeisser [7] for polynomials with real coefficients, which have at most  $n - 1$  distinct zeros in  $(-1, 1)$ .

Similarly in  $L^2$  norm,

$$\|P\| = \|P\|_2 = \left( \int_{-1}^1 |P(x)|^2 dx \right)^{1/2},$$

Tariq [10] improved the following result of Labelle [2]

$$|a_k| \leq \frac{1 \cdot 3 \cdot 5 \cdots (2k-1)}{k!} \left( k + \frac{1}{2} \right)^{1/2} \binom{[(n-k)/2] + k + 1/2}{[(n-k)/2]} \|P\|_2. \quad (1.3)$$

for  $P \in \mathcal{P}_n$  and  $0 \leq k \leq n$ , where the symbol  $[x]$  denotes as usual the integral part of  $x$ . Equality in this case is attained only for the constant multiples of the polynomial

$$\sum_{\nu=0}^{[(n-k)/2]} (-1)^{\nu} (4\nu + 2k + 1) \binom{k + \nu - 1/2}{\nu} P_{k+2\nu}(x),$$

where  $P_m(x)$  denotes the Legendre polynomial of degree  $m$ .

Under restriction  $P(1) = 0$ , Tariq [10] proved that

$$|a_n| \leq \frac{n}{n+1} \cdot \frac{(2n)!}{2^n (n!)^2} \left( \frac{2n+1}{2} \right)^{1/2} \|P\|_2, \quad (1.4)$$

with equality case  $P(x) = P_n(x) - \frac{1}{n^2} \sum_{\nu=0}^{n-1} (2\nu+1) P_{\nu}(x)$ . Also, he obtained that

$$|a_{n-1}| \leq \frac{(n^2+2)^{1/2}}{n+1} \cdot \frac{(2n-2)!}{2^{n-1} ((n-1)!)^2} \left( \frac{2n-1}{2} \right)^{1/2} \|P\|_2, \quad (1.5)$$

with equality case

$$P(x) = \frac{2n+1}{n^2+2} P_n(x) - P_{n-1}(x) + \frac{1}{n^2+2} \sum_{\nu=0}^{n-2} (2\nu+1) P_{\nu}(x).$$

In the absence of the hypothesis  $P(1) = 0$  the factor  $(n^2 + 2)^{1/2}/(n + 1)$  appearing on the right-hand side of (1.5) is to be dropped.

This result was extended by Milovanović and Guessab [4] for polynomials with real coefficients, which have  $m$  zeros on real line.

In this paper we consider more general problem including  $L^2$  norm of polynomials with respect to a nonnegative measure on the real line  $\mathbb{R}$ . The generalized Laguerre measure is also included.

### 2. Main results

Let  $d\sigma(x)$  be a given nonnegative measure on the real line  $\mathbb{R}$ , with compact or infinite support, for which all moments  $\mu_k = \int_{\mathbb{R}} x^k d\sigma(x)$ ,  $k = 0, 1, \dots$ , exist and are finite, and  $\mu_0 > 0$ . In that case, there exist a unique set of orthonormal polynomials  $\pi_n(\cdot) = \pi_n(\cdot; d\sigma)$ ,  $n = 0, 1, \dots$ , defined by

$$\begin{aligned} \pi_n(x) &= b_n^{(n)}(d\sigma)x^n + b_{n-1}^{(n)}(d\sigma)x^{n-1} + \dots + b_0^{(n)}(d\sigma), & b_n^{(n)}(d\sigma) > 0, \\ (\pi_n, \pi_m) &= \delta_{nm}, & n, m \geq 0, \end{aligned}$$

where

$$(f, g) = \int_{\mathbb{R}} f(x)\overline{g(x)} d\sigma(x) \quad (f, g \in L^2(\mathbb{R})). \tag{2.1}$$

For  $P \in \mathcal{P}_n$ , we define

$$\|P\|_{d\sigma} = \sqrt{(P, P)} = \left( \int_{\mathbb{R}} |P(x)|^2 d\sigma(x) \right)^{1/2}. \tag{2.2}$$

Also, for  $\xi_k \in \mathbb{C}$ ,  $k = 1, \dots, m$ , we define a restricted polynomial class

$$\mathcal{P}_n(\xi_1, \dots, \xi_m) = \{P \in \mathcal{P}_n \mid P(\xi_k) = 0, k = 1, \dots, m\} \quad (0 \leq m \leq n).$$

In the case  $m = 0$  this class of polynomials reduces to  $\mathcal{P}_n$ . The case  $m = n$  is trivial. If  $\xi_1 = \dots = \xi_k = \xi$  ( $1 \leq k \leq m$ ) then the restriction on polynomials at the point  $x = \xi$  becomes  $P(\xi) = P'(\xi) = \dots = P^{(k-1)}(\xi) = 0$ .

Let

$$\prod_{i=1}^m (x - \xi_i) = x^m - s_1 x^{m-1} + \dots + (-1)^{m-1} s_{m-1} x + (-1)^m s_m$$

where  $s_k$  denotes elementary symmetric functions of  $\xi_1, \dots, \xi_m$ , i.e.,

$$s_k = \sum \xi_1 \cdots \xi_k \quad \text{for } k = 1, \dots, m. \tag{2.3}$$

For  $k = 0$  we have  $s_0 = 1$ , and  $s_k = 0$  for  $k > m$  or  $k < 0$ .

**THEOREM 2.1.** *Let  $P \in \mathcal{P}_n(\xi_1, \dots, \xi_m)$  and  $s_1, \dots, s_m$  be given by (2.3). If the measure  $d\hat{\sigma}(x)$  is given by*

$$d\hat{\sigma}(x) = \prod_{k=1}^m |x - \xi_k|^2 d\sigma(x) \quad (2.4)$$

and  $\|P\|_{d\sigma}$  is defined by (2.2), then

$$|a_{n-k}| \leq \left( \sum_{j=0}^k \left( \sum_{i=j}^k (-1)^{k-i} s_{k-i} \hat{b}_{n-m-i}^{(n-m-j)} \right)^2 \right)^{1/2} \|P\|_{d\sigma}, \quad (2.5)$$

for  $k = 0, 1, \dots, n$ , where  $\hat{b}_\nu^\mu = b_\nu^\mu(d\hat{\sigma})$ ,  $\nu = 0, 1, \dots, \mu$ , are the coefficients in the orthonormal polynomial  $\hat{\pi}_\mu(\cdot) = \pi_\mu(\cdot; d\hat{\sigma})$ .

*Inequality (2.5) is sharp and becomes an equality if and only if  $P(x)$  is a constant multiple of the polynomial*

$$\left( \sum_{j=0}^k \hat{\pi}_{n-m-j}(x) \sum_{i=j}^k (-1)^{k-i} s_{k-i} \hat{b}_{n-m-i}^{(n-m-j)} \right) \prod_{k=1}^m (x - \xi_k).$$

*Proof.* At first we consider the inner product (2.1). Then the polynomial  $P(x) = \sum_{\nu=0}^n a_\nu x^\nu \in \mathcal{P}_n$  can be represented in the form  $P(x) = \sum_{\nu=0}^n \alpha_\nu \pi_\nu(x; d\sigma)$ , where  $\alpha_\nu = (P, \pi_\nu)$ ,  $\nu = 0, 1, \dots, n$ . Then we have

$$a_{n-k} = \sum_{i=0}^k \alpha_{n-i} b_{n-k}^{(n-i)}(d\sigma) = \left( P, \sum_{i=0}^k b_{n-k}^{(n-i)}(d\sigma) \pi_{n-i} \right), \quad k = 0, 1, \dots, n, \quad (2.6)$$

where  $\pi_\nu(\cdot) = \pi_\nu(\cdot; d\sigma)$ .

Suppose now that  $P \in \mathcal{P}_n(\xi_1, \dots, \xi_m)$ . Then we can write

$$P(x) = Q(x) \prod_{k=1}^m (x - \xi_k), \quad (2.7)$$

where  $Q(x) = a'_{n-m} x^{n-m} + a'_{n-m-1} x^{n-m-1} + \dots + a'_0 \in \mathcal{P}_{n-m}$ . Also, we have

$$\prod_{i=1}^m (x - \xi_i) = x^m - s_1 x^{m-1} + \dots + (-1)^{m-1} s_{m-1} x + (-1)^m s_m$$

where  $s_k$ ,  $k = 0, 1, \dots, m$ , denotes elementary symmetric functions (2.3). Now, putting this in (2.7), we obtain

$$P(x) = \sum_{i=0}^{n-m} \sum_{\nu=0}^m a'_i (-1)^\nu s_\nu x^{m+i-\nu} = \sum_{k=0}^n a_{n-k} x^{n-k},$$

where

$$a_{n-k} = \sum_{i=0}^k a'_{n-m-i} (-1)^{k-i} s_{k-i}, \quad k = 0, 1, \dots, n, \quad (2.8)$$

and  $a'_k = 0$  for  $k < 0$  and  $k > n - m$ .

Now, the corresponding equalities (2.6) for polynomial  $Q$  in the measure  $d\hat{\sigma}(x)$ , given by (2.4), become

$$a'_{n-m-i} = \left( Q, \sum_{j=0}^i \hat{b}_{n-m-i}^{(n-m-j)} \hat{\pi}_{n-m-j} \right), \quad i = 0, 1, \dots, n - m, \quad (2.9)$$

where  $\hat{\pi}_\nu(\cdot) = \pi_\nu(\cdot; d\hat{\sigma})$ .

According to (2.7), we have

$$a_{n-k} = \sum_{i=0}^k (-1)^{k-i} s_{k-i} \left( Q, \sum_{j=0}^i \hat{b}_{n-m-i}^{(n-m-j)} \hat{\pi}_{n-m-j} \right) = (Q, W_{n-m}) \quad (2.10)$$

where

$$\begin{aligned} W_{n-m}(x) &= \sum_{i=0}^k (-1)^{k-i} s_{k-i} \sum_{j=0}^i \hat{b}_{n-m-i}^{(n-m-j)} \hat{\pi}_{n-m-j}(x) \\ &= \sum_{j=0}^k \hat{\pi}_{n-m-j}(x) \sum_{i=j}^k (-1)^{k-i} s_{k-i} \hat{b}_{n-m-i}^{(n-m-j)} \end{aligned}$$

and  $\hat{b}_\nu^{(\mu)} = 0$  for  $\nu < 0$ . Now, using Cauchy inequality we get

$$|a_{n-k}| \leq C_{n,n-k} \|Q\|_{d\hat{\sigma}}$$

where  $C_{n,n-k} = \|W_{n-m}\|_{d\hat{\sigma}} = \left( \sum_{j=0}^k \left( \sum_{i=j}^k (-1)^{k-i} s_{k-i} \hat{b}_{n-m-i}^{(n-m-j)} \right)^2 \right)^{1/2}$ . Since

$$\|Q\|_{d\hat{\sigma}}^2 = \int_{\mathbb{R}} |Q(x)|^2 d\hat{\sigma}(x) = \int_{\mathbb{R}} |P(x)|^2 d\sigma(x) = \|P\|_{d\sigma}^2$$

we obtain inequality (2.5).

The extremal polynomial is  $x \mapsto W_{n-m}(x) \prod_{k=1}^m (x - \xi_k)$ .  $\square$

*Remark 2.1.* For  $k = 0$  and  $k = 1$  Theorem 2.1 gives the results obtained by Milovanović and Guessab [4] (see also [6, pp. 432–439]).

Consider now the generalized Laguerre measure  $d\sigma(x) = x^\alpha e^{-x} dx$ ,  $\alpha > -1$ , on  $(0, +\infty)$ . With  $\tilde{L}_n^{(\alpha)}(x)$  we denote the generalized orthonormal Laguerre polynomial. The coefficient  $b_k^{(n)}$  of  $x^k$  in  $\tilde{L}_n^{(\alpha)}(x)$  is given by

$$b_k^{(n)} = (-1)^{n-k} \binom{n}{k} \frac{(\alpha + k + 1)_{n-k}}{\sqrt{n! \Gamma(n + \alpha + 1)}}.$$

As a direct corollary of Theorem 2.1, we have:

COROLLARY 2.2. *Under restriction  $P^{(i)}(0) = 0$ ,  $i = 0, 1, \dots, m-1$ , we have that*

$$|a_{n-k}| \leq \sqrt{A_{n,k}} \|P\|_{d\sigma},$$

where

$$A_{n,k} = \frac{1}{(n-m-k)! \Gamma(n+m-k+\alpha+1)} \sum_{j=0}^k \binom{n+m-j+\alpha}{k-j} \binom{n-m-j}{k-j}$$

for  $n-k \geq m$ , and  $A_{n,k} = 0$  for  $n-k < m$ . The equality is attained if and only if  $P(x)$  is a constant multiple of the polynomial

$$x^m \sum_{j=0}^k \hat{b}_{n-m-k}^{(n-m-j)} \tilde{L}_{n-m-j}^{(\alpha+2m)}(x).$$

#### REFERENCES

- [1] A. Giroux and Q.I. Rahman, *Inequalities for polynomials with a prescribed zero*, Trans. Amer. Math. Soc. **193** (1974), 67–98.
- [2] G. Labelle, *Concerning polynomials on the unit interval*, Proc. Amer. Math. Soc. **20** (1969), 321–326.
- [3] V.A. Markov, *On functions deviating least from zero in a given interval*, Izdat. Imp. Akad. Nauk, St. Petersburg 1892 (Russian) [German transl. Math. Ann. **77** (1916), 218–258].
- [4] G.V. Milovanović and A. Guessab, *An estimate for coefficients of polynomials in  $L^2$  norm*, Proc. Amer. Math. Soc. **120** (1994), 165–171.
- [5] G.V. Milovanović and L.Z. Marinković, *Extremal problems for coefficients of algebraic polynomials*, Facta Univ. Ser. Math. Inform. **5** (1990), 25–36.
- [6] G.V. Milovanović, D.S. Mitrinović and Th.M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific, Singapore-New Jersey-London-Hong Kong, 1994.
- [7] Q.I. Rahman and G. Schmeisser, *Inequalities for polynomials on the unit interval*, Trans. Amer. Math. Soc. **231** (1977), 93–100.
- [8] I. Schur, *Über das Maximum des absoluten Betrages eines Polynoms in einem gegebenen Intervall*, Math. Z. **4** (1919), 217–287.
- [9] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ., vol. 23, 4th ed., Amer. Math. Soc., Providence, R.I., 1975.
- [10] Q.M. Tariq, *Concerning polynomials on the unit interval*, Proc. Amer. Math. Soc. **99** (1987), 293–296.

Elektronski fakultet  
Katedra za matematiku  
18001 p.p. 73  
Yugoslavia  
e-mail: grade@efnis.elfak.ni.ac.yu

(Received 29 06 1995)