

BANDS OF NIL-EXTENSIONS OF RIGHT SIMPLE SEMIGROUPS

Petar V. Protić

Communicated by Žarko Mijajlović

Abstract. We prove a general theorem when the semigroup is a band of nil-extensions of right simple semigroups. We also describe normal bands of right simple semigroups, normal bands of right Archimedean semigroups and normal bands of nil-extensions of right simple semigroups.

Nil-extensions of some classes of semigroups were considered by many authors, for example by L.N. Ševrin, J.L. Galbiati, M.L. Veronesi, S. Bogdanović and M. Ćirić. All of these results are presented in a survey paper [2].

Let N be the set of all positive integers. A semigroup S is *right Archimedean* (or *r-Archimedean*) if, for every $a, b \in S$, there exists $n \in N$ such that $a^n \in bS$. A semigroup S is *right simple* if $a \in bS$ for every $a, b \in S$. A semigroup S is *right π -regular semigroup* if for all $a \in S$ there exists $n \in N$ such that $a^n \in a^{n+1}S$. A semigroup S with the zero 0 is *nil* if for every $a \in S$ there exists $n \in N$ such that $a^n = 0$. An ideal extension S of T is a *nil-extension* if S/T is a nil semigroup. A semigroup B is a *band* if $x^2 = x$ for each $x \in B$.

A semigroup S is a *band Y of semigroups S_α* if $S = \bigcup_{\alpha \in Y} S_\alpha$, Y is a band, $S_\alpha \cap S_\beta = \emptyset$ for $\alpha, \beta \in Y$ with $\alpha \neq \beta$ and $S_\alpha S_\beta \subseteq S_{\alpha\beta}$. A congruence ρ on S is called *band congruence* if S/ρ is a band.

For undefined notions and notation we refer to [1], [3] and [4].

THEOREM 1. [6] *A semigroup S is a band of r-Archimedean semigroups if and only if*

$$(1) \quad (\forall a \in S)(\forall x, y \in S^1)(\exists i, j \in N)(xay)^i \in xa^2yS, (xa^2y)^j \in xayS. \quad \square$$

In [6] is proved that if (1) holds, then the relation ρ defined on S by

$$(2) \quad a\rho b \iff (\forall x, y \in S^1)(\exists i, j \in N)(xay)^i \in xbyS, (xby)^j \in xayS$$

is a band congruence on S .

THEOREM 2. [1] *A semigroup S is a nil-extension of a right simple semigroup if and only if S is r -Archimedean and right π -regular. \square*

THEOREM 3. *The following conditions on a semigroup S are equivalent:*

- (i) S is a band of nil-extensions of right simple semigroups,
- (ii) $(\forall a \in S)(\forall x, y \in S^1)(\exists m, n \in N)(\forall p, q \in N)$

$$(xay)^m \in (xa^2y)^pS, (xa^2y)^n \in (xay)^qS,$$

- (iii) $(\forall a \in S)(\forall x, y \in S^1)(\exists m, n \in N)$

$$(xay)^m \in (xa^2y)^mS, (xa^2y)^n \in (xay)^nS.$$

Proof. (i) \implies (ii). Let S be a band Y of nil-extensions S_α of right simple semigroups K_α . Then S^1 is a band Y^1 of nil-extensions S_α of right simple semigroups K_α , $\alpha \in Y^1$ and $S_1 = 1 = K_1$. If $x \in S_\alpha$, $a \in S_\beta$, $y \in S_\gamma$ then $xay, xa^2y \in S_{\alpha\beta\gamma}$. Now there exist $m, n \in N$ such that $(xay)^m, (xa^2y)^n \in K_{\alpha\beta\gamma}$. Also, $(xa^2y)^{np} \in K_{\alpha\beta\gamma}$ for every $p \in N$. Since $K_{\alpha\beta\gamma}$ is a right simple semigroup it follows that

$$(xay)^m \in (xa^2y)^{np}K_{\alpha\beta\gamma} \subseteq (xa^2y)^pS.$$

Similarly, $(xa^2y)^n \in (xay)^qS$ for every $q \in N$, and so (ii) holds.

(ii) \implies (iii). This follows trivially.

(iii) \implies (i). From (iii) we have

$$(\forall a \in S)(\forall x, y \in S^1)(\exists m, n \in N) (xay)^m \in xa^2yS, (xa^2y)^n \in xayS,$$

and by Theorem 1 we conclude that semigroup S is a band of r -Archimedean semigroups S_α , $\alpha \in Y$. Also, from (iii) for $x = a = y \in S_\alpha$, it follows that $a^{3m} \in a^{4m}S$ and so there exists $u \in S_\beta$ such that $a^{3m} = a^{4m}u$. From above we have $\alpha = \alpha\beta$. Now $a^{2m-1}u^2 \in S_\alpha$ and

$$a^{3m} = a^{4m}u = a^m a^{3m}u = a^m a^{4m}uu = a^{3m+1}a^{2m-1}u^2 \in a^{3m+1}S_\alpha.$$

Hence, S_α is a right π -regular r -Archimedean semigroup and by Theorem 2 it follows that S_α is a nil-extension of a right simple semigroup, whence semigroup S is a band of nil-extensions of right simple semigroups. \square

Recall that a band B is a *normal band* if $efge = egfe$ for every $e, f, g \in B$.

THEOREM 4. *A semigroup S is a normal band of r -Archimedean semigroups if and only if*

$$(3) \quad (\forall u, v, w, t \in S)(\exists n \in N) (uvt)^n \in uvvtS.$$

Proof. Let S be a normal band Y of r -Archimedean semigroups S_α . If $u \in S_\alpha$, $v \in S_\beta$, $w \in S_\gamma$, $t \in S_\delta$ then $uvt \in S_{\alpha\beta\gamma\delta}$. Since Y is a normal band we have $uvtvuwt \in S_{\alpha\beta\gamma\delta}$ and, since $S_{\alpha\beta\gamma\delta}$ is a right Archimedean semigroup, there exists $n \in N$ such that

$$(uvt)^n \in uvtvuwtS_{\alpha\beta\gamma\delta} \subseteq uvvtS$$

and so (3) holds.

Conversely, let statement (3) hold in S and let $a \in S$, $x, y \in S^1$. Then, by (3), for $u = xa$, $v = ayxa^2$, $w = yxa$, $t = ay$, there exists $n \in N$ such that

$$(xa^2y)^{3n} = (xa \cdot ayxa^2 \cdot yxa \cdot ay)^n \in xa \cdot yxa \cdot ayxa^2 \cdot ayS \subseteq xayS.$$

Also, for $u = xa$, $v = xayax$, $w = ay$, $t = xay$, there exists $m \in N$ such that

$$(xay)^{4m} = (xa \cdot xayax \cdot ay \cdot xay)^m \in xa \cdot ay \cdot xayax \cdot xayS \in xa^2yS.$$

By Theorem 1 we have that S is a band Y of r -Archimedean semigroups S_α .

We shall prove that the congruence ρ defined by (2) is a normal band congruence on S . Let $a, b, c \in S$ and $x, y \in S^1$. For $u = xa$, $v = b$, $w = c$, $t = ay$ by (3) there exists $n \in N$ such that $(xabcay)^n \in xabcayS$. Similarly, $(xacbay)^m \in xacbayS$ for some $m \in N$. Hence $abcapacba$ and ρ is a normal band congruence on S . It follows that S is a normal band of r -Archimedean semigroups. \square

THEOREM 5. *The following conditions on a semigroup S are equivalent:*

- (i) S is a normal band of nil-extensions of right simple semigroups,
- (ii) $(\forall u, v, w, t \in S)(\exists n \in N)(\forall k \in N) (uvt)^n \in (uvtvuwt)^k S$,
- (iii) $(\forall u, v, w, t \in S)(\exists n \in N) (uvt)^n \in (uvtvuwt)^n S$.

Proof. (i) \implies (ii). Let S be a normal band Y of nil-extensions S_α of right simple semigroups K_α . If $u \in S_\alpha$, $v \in S_\beta$, $w \in S_\gamma$, $t \in S_\delta$, then $uvt \in S_{\alpha\beta\gamma\delta}$. Since $\alpha\beta\gamma\delta = \alpha\gamma\beta\delta\beta\alpha\gamma\delta$ we have that $uvtvuwt \in S_{\alpha\beta\gamma\delta}$. Now, there exist $n, m \in N$ such that $(uvt)^n, (uvtvuwt)^m \in K_{\alpha\beta\gamma\delta}$ and, also $(uvtvuwt)^{mk} \in K_{\alpha\beta\gamma\delta}$ for every $k \in N$. Since $K_{\alpha\beta\gamma\delta}$ is a right simple semigroup, we conclude that

$$(uvt)^n \in (uvtvuwt)^{mk} K_{\alpha\beta\gamma\delta} \subseteq (uvtvuwt)^k S.$$

(ii) \implies (iii). This follows trivially.

(iii) \implies (i). From (iii) we have

$$(\forall u, v, w, t \in S)(\exists n \in N) (uvwt)^n \in (uwtvuw)^n S \subseteq uvwtS$$

and by Theorem 4 S is a normal band Y of r -Archimedean semigroups S_α . If $a \in S_\alpha$, then from (iii) for $a = u = v = w = t$ there exists $n \in N$ such that $u^{4n} \in u^{8n}S$ and $u^{4n} = u^{8n}s$ for some $s \in S$. If $s \in S_\beta$, then it follows that $\alpha = \alpha\beta$. Now $u^{4n-1}s \in S_\alpha$ and so

$$u^{4n} = u^{8n}s = u^{4n+1}u^{4n-1}s \in u^{4n+1}S_\alpha.$$

Hence S_α is right π -regular and by Theorem 2 we conclude that S_α is a nil-extension of right simple semigroup and S is a normal band of nil-extensions of right simple semigroups. \square

THEOREM 6. *A semigroup S is a normal band of right simple semigroups if and only if*

$$(4) \quad (\forall u, v, w, t \in S) uvwt \in uwvtS, u \in u^2S.$$

Proof. Let S be a normal band Y of right simple semigroups S_α . If $u \in S_\alpha$, $v \in S_\beta$, $w \in S_\gamma$, $t \in S_\delta$, then $uvwt$, $uwvtvuw$ $\in S_{\alpha\beta\gamma\delta}$. Since $S_{\alpha\beta\gamma\delta}$ is a right simple semigroup we conclude that

$$uvwt \in uwvtvuwS \subseteq uwvtS.$$

Since $u, u^2 \in S_\alpha$, we have $u \in u^2S$.

Conversely, let (4) hold. We define on S a relation σ by

$$a\sigma b \iff a \in bS, b \in aS.$$

Clearly $\sigma \subseteq \mathcal{R}$ where \mathcal{R} is the Green \mathcal{R} -relation. Conversely, let $a\mathcal{R}b$. If $a \neq b$, then $a \in bS$, $b \in aS$ and so $a\sigma b$. If $a = b$, then by (4) we have $a \in a^2S \subseteq aS$, and so $a\sigma a$. Hence $\mathcal{R} \subseteq \sigma$ and so $\sigma = \mathcal{R}$.

Let $a\mathcal{R}b$, $c \in S$; then $ac \in bSc$. Hence $ac = btc$ for some $t \in S$. Now by (4) we have $ac = btc \in btc^2S \subseteq bctcS \subseteq bcS$. Similarly, $bc \in acS$ and so $ac\mathcal{R}bc$. Since \mathcal{R} is a left congruence on an arbitrary semigroup, we conclude that \mathcal{R} is a congruence relation on S .

Let $a \in S$; then $a^2 \in aS$. By (4) we have $a \in a^2S$ and so $a\mathcal{R}a^2$. Hence \mathcal{R} is a band congruence. If $a, b, c \in S$, then by (4) it follows that $abca\mathcal{R}acba$ and so \mathcal{R} is a normal band congruence on S . Now, $S = \bigcup_{\alpha \in Y} S_\alpha$, Y is a normal band and S_α are \mathcal{R} -classes.

Let $a\mathcal{R}b$; then $a\mathcal{R}b^2$, whence $a = b^2x$ for some $x \in S$. If $a, b \in S_\alpha$ and $x \in S_\beta$, then $\alpha = \alpha\beta$. By (4) from $a \in a^2S$ we have $a = a^2y$ for some $y \in S$. If $y \in S_\gamma$ then $\alpha = \alpha\gamma$. Now from above we have

$$a = a^2y = aay = b^2xay = bbxay \in bS_{\alpha\beta\alpha\gamma} = bS_\alpha.$$

Hence S_α is a right simple semigroup and so the semigroup S is a normal band of right simple semigroups. \square

REFERENCES

- [1] S. Bogdanović, *Semigroups with a system of subsemigroups*, Inst. of Math. Novi Sad, 1985.
- [2] S. Bogdanović and M. Ćirić, *Semilattices of Archimedean semigroups and (completely) π -regular semigroups I (A survey)*, Zb. Rad. Fil. Fak. (Niš), Ser. Mat. **7** (1993), 1–40.
- [3] M. Petrich, *Introduction to Semigroups*, Merrill, Columbus, 1973.
- [4] M. Petrich, *Lectures in Semigroups*, Akad. Verlag, Berlin, 1977.
- [5] P. V. Protić, *On some band decompositions of semigroups*, Publ. Math. Debrecen **45** (1994), 205–211.
- [6] M. S. Putcha, *Band of t -Archimedean semigroups*, Semigroup Forum **6**(1973), 232–239.

Građevinski Fakultet
18000 Niš
Yugoslavia
pvprotic@ziux.grafak.ni.ac.yu

(Received 10 03 1997)
(Revised 16 07 1997)