

DISTANCE OF THORNY GRAPHS

Ivan Gutman

Communicated by Slobodan Simić

Abstract. Let G be a connected graph on n vertices. The thorn graph G^* of G is obtained from G by attaching to its i -th vertex p_i new vertices of degree one, $p_i \geq 0$, $i = 1, 2, \dots, n$. Let $d(G)$ be the sum of distances of all pairs of vertices of G . We establish relations between $d(G)$ and $d(G^*)$ and examine several special cases of this result. In particular, if $p_i = \gamma - \gamma_i$, where γ is a constant and γ_i the degree of the i -th vertex in G , and if G is a tree, then there is a linear relation between $d(G^*)$ and $d(G)$, namely $d(G^*) = (\gamma - 1)^2 d(G) + [(\gamma - 1)n + 1]^2$.

Introduction

In this paper we consider connected finite graphs without loops and multiple edges. Let G be such a graph, $V = V(G)$ its vertex set, $E = E(G)$ its edge set, and let its vertices, whose number is n , be labeled by u_1, u_2, \dots, u_n . The distance (= length of a shortest path) between the vertices u_i and u_j of G is denoted by $d(u_i, u_j|G)$. The sum of the distances between all pairs of vertices of G is *the distance of the graph G* and is denoted by $d(G)$.

In the mathematical literature the distance of a graph was first introduced by Entringer, Jackson and Snyder [2], although the chemical applications of this quantity are somewhat older [8]. For results concerning the distance of compound graphs as well as for additional references see [3, 6, 9].

Let p_1, p_2, \dots, p_n be non-negative integers.

Definition 1. The *thorn graph* of the graph G , with parameters p_1, p_2, \dots, p_n , is obtained by attaching p_i new vertices of degree one to the vertex u_i of the graph G , $i = 1, 2, \dots, n$.

The thorn graph of the graph G will be denoted by G^* , or if the respective parameters need to be specified, by $G^*(p_1, p_2, \dots, p_n)$.

AMS Subject Classification (1991): Primary 05C12

Supported by Ministry of Science and Technology of Serbia, grant number 04M03/C

In this work we examine the relation between $d(G)$ and $d(G^*)$. The motivation for this study comes from a particular special case, namely $G^*(\gamma - \gamma_1, \gamma - \gamma_2, \dots, \gamma - \gamma_n)$, where γ_i is the degree of the i -th vertex of G and γ is a constant ($\gamma \geq \gamma_i$ for all $i = 1, 2, \dots, n$). Then the vertices of G^* are either of degree γ or of degree one. If, in addition, $\gamma = 4$, then the thorn graph is just what Cayley [1] calls a “plerogram” and Pólya [7] a “C–H graph”. (The parent graph G would then be referred to as a “kenogram” [1] or a “C–graph” [7]. Clearly, these notions have their origins in the attempts to represent molecular structure by means of graphs [4].) It is also worth mentioning that the so-called “caterpillars” [5] are thorn graphs whose parent graph is a path.

Denote the vertex set of G^* by V^* . Further, the set of degree-one vertices of G^* , attached to the vertex u_i is V_i . Its cardinality is p_i and, clearly,

$$V^* = V \cup V_1 \cup V_2 \cup \dots \cup V_n \quad \text{and} \quad V_i \cap V_j = \emptyset \quad \text{for } i \neq j$$

The main results

Let $\{x, y\} \subseteq V^*$. In order to compute $d(G^*)$ we distinguish between four types of pairs of vertices of G^* :

Type 1. $x \in V$, $y \in V$;

Type 2. $x \in V_i$, $y \in V$, for some i , $1 \leq i \leq n$;

Type 3. $x \in V_i$, $y \in V_j$, for some i, j , $1 \leq i < j \leq n$;

Type 4. $x \in V_i$, $y \in V_i$, for some i , $1 \leq i \leq n$.

Let the contributions of all such vertex pairs to $d(G^*)$ be denoted by F_1, F_2, F_3 and F_4 , respectively. Then,

$$d(G^*) = F_1 + F_2 + F_3 + F_4 \tag{1}$$

If $\{x, y\}$ is a vertex pair of Type 1, then $d(x, y|G^*) = d(x, y|G)$ and therefore

$$F_1 = d(G)$$

There are p_i vertex pairs $\{x, y\}$ of Type 2, and for each of them $d(x, y|G^*) = d(u_i, y|G) + 1$. Therefore

$$F_2 = \sum_{i=1}^n \sum_{y \in V} p_i [d(u_i, y|G) + 1] = \sum_{1 \leq i < j \leq n} (p_i + p_j) d(u_i, u_j|G) + n \sum_{i=1}^n p_i$$

There are $p_i \cdot p_j$ vertex pairs $\{x, y\}$ of Type 3, and for each of them $d(x, y|G^*) = d(u_i, u_j|G) + 2$. Therefore

$$F_3 = \sum_{1 \leq i < j \leq n} p_i p_j [d(u_i, u_j|G) + 2] = \sum_{1 \leq i < j \leq n} p_i p_j d(u_i, u_j|G) + \left(\sum_{i=1}^n p_i \right)^2 - \sum_{i=1}^n p_i^2$$

There are $\binom{p_i}{2}$ vertex pairs $\{x, y\}$ of Type 4, each of them at distance 2. Therefore

$$F_4 = \sum_{i=1}^n 2 \binom{p_i}{2} = \sum_{i=1}^n p_i^2 - \sum_{i=1}^n p_i$$

Substituting the above relations back into Eq. (1) we arrive at the general expression for the distance of a thorn graph:

THEOREM 1. *If G^* is the thorn graph of the graph G , with parameters p_i , $p_i \geq 0$, $i = 1, 2, \dots, n$, then*

$$\begin{aligned} d(G^*) &= d(G) + \sum_{1 \leq i < j \leq n} (p_i + p_j) d(u_i, u_j | G) + \sum_{1 \leq i < j \leq n} p_i p_j d(u_i, u_j | G) \\ &\quad + \left(\sum_{i=1}^n p_i \right)^2 + (n-1) \sum_{i=1}^n p_i \end{aligned} \quad (2)$$

i.e.,

$$d(G^*) = \sum_{1 \leq i < j \leq n} (p_i + 1)(p_j + 1) d(u_i, u_j | G) + \left(\sum_{i=1}^n p_i \right)^2 + (n-1) \sum_{i=1}^n p_i \quad (3)$$

COROLLARY 1.1. *If G^* is the thorn graph of the graph G , with parameters $p_1 = p_2 = \dots = p_n = p$, then $d(G^*)$ and $d(G)$ are related as:*

$$d(G^*) = (p+1)^2 d(G) + np(np+n-1)$$

Theorem 1 has been obtained by a routine combinatorial reasoning and the form of Eqs. (2) and (3) is neither appealing nor unexpected. Also, the existence of a simple linear connection between $d(G^*(p, p, \dots, p))$ and $d(G)$, as specified in Corollary 1.1, is by no means a surprise. The following two special cases seem, however, to be somewhat less self-evident and hardly could have been anticipated.

Let T be an n -vertex tree and let γ_i be the degree of its i -th vertex. Recall that

$$\gamma_1 + \gamma_2 + \dots + \gamma_n = 2n - 2$$

COROLLARY 1.2. *If T^* is the thorn graph of the tree T , with parameters $p_i = \gamma_i$, $i = 1, 2, \dots, n$, then $d(T^*)$ and $d(T)$ are related as:*

$$d(T^*) = 9 d(T) + (n-1)(3n-5)$$

COROLLARY 1.3. *Let γ be an integer with the property $\gamma \geq \gamma_i$, $i = 1, 2, \dots, n$. If T^* is the thorn graph of the tree T , with parameters $p_i = \gamma - \gamma_i$, $i = 1, 2, \dots, n$, then $d(T^*)$ and $d(T)$ are related as:*

$$d(T^*) = (\gamma-1)^2 d(T) + [(\gamma-1)n+1]^2$$

Proofs

Theorem 1 has already been verified, and Corollary 1.1 is immediate. What remains is to demonstrate the validity of Corollaries 1.2 and 1.3. For this we need:

LEMMA 1. *If T is a tree on n vertices u_1, u_2, \dots, u_n and if the degree of u_i is γ_i , $i = 1, 2, \dots, n$, then*

$$\sum_{1 \leq i < j \leq n} (\gamma_i + \gamma_j) d(u_i, u_j | T) = 4d(T) - n(n-1) \quad (4)$$

and

$$\sum_{1 \leq i < j \leq n} \gamma_i \gamma_j d(u_i, u_j | T) = 4d(T) - (n-1)(2n-1) \quad (5)$$

Proof. Let $V(T)$ and $E(T)$ be the vertex and edge sets, respectively, of the tree T . Let $e \in E(T)$. Then the subgraph whose vertex set is $V(T)$ and whose edge set is $E(T) \setminus \{e\}$ consists of two components, $T_1 = T_1(e)$ and $T_2 = T_2(e)$, possessing $n_1 = n_1(e|T)$ and $n_2 = n_2(e|T)$ vertices, respectively. Recall that $n_1(e|T) + n_2(e|T) = n$ holds for all $e \in E(T)$.

It is long known [8] that the distance $d(T)$ of a tree T (in which the path between any two vertices is unique) may be calculated by counting the paths of T which contain the edge e , and summing this count over all edges of T . Now, the number of paths of T containing e is $n_1(e|T) \cdot n_2(e|T)$ and therefore,

$$d(T) = \sum_{e \in E(T)} n_1(e|T) n_2(e|T) \quad (6)$$

which may be rewritten as

$$d(T) = \sum_{e \in E(T)} \sum_{u_i \in V(T_1)} \sum_{u_j \in V(T_2)} 1 \quad (7)$$

Associate to each pair of vertices $u_i, u_j \in V(G)$ a weight ω_{ij} and define a generalized distance-sum $d_\omega(G)$ as

$$d_\omega(G) = \sum_{1 \leq i < j \leq n} \omega_{ij} d(u_i, u_j | G)$$

Clearly, if $\omega_{ij} = 1$ for all i, j , $1 \leq i < j \leq n$ then $d_\omega(G) = d(G)$.

Now, repeating the reasoning leading to Eq. (7), and bearing in mind that for T being a tree, in $d_\omega(T)$ the distance between the vertices u_i, u_j has to be counted with weight ω_{ij} , we obtain

$$d_\omega(T) = \sum_{e \in E(T)} \sum_{u_i \in V(T_1)} \sum_{u_j \in V(T_2)} \omega_{ij} \quad (8)$$

If we choose $\omega_{ij} = \gamma_i + \gamma_j$ then Eq. (8) yields

$$\sum_{1 \leq i < j \leq n} (\gamma_i + \gamma_j) d(u_i, u_j | T) = \sum_{e \in E(T)} \left[n_2 \sum_{u_i \in V(T_1)} \gamma_i + n_1 \sum_{u_j \in V(T_2)} \gamma_j \right] \quad (9)$$

Because T_1 and T_2 have $n_1 - 1$ and $n_2 - 1$ edges, respectively,

$$\sum_{u_i \in V(T_1)} \gamma_i = 2(n_1 - 1) + 1 \quad (10)$$

and

$$\sum_{u_j \in V(T_2)} \gamma_j = 2(n_2 - 1) + 1 \quad (11)$$

Recall that γ_i and γ_j in Eqs. (10) and (11) are the degrees of the vertices of the tree T ; they coincide with the degrees of the vertices of the trees T_1 and T_2 , respectively, except for one particular vertex of T_1 and for one particular vertex of T_2 . These “exceptional” vertices of T_1 and T_2 have degrees by one less than in T . The terms $+1$ on the right-hand sides of Eqs. (10) and (11) occur because of this difference between the vertex degrees of T and the vertex degrees of T_1 and T_2 .

Substituting (10) and (11) back into Eq. (9) results in

$$\sum_{1 \leq i < j \leq n} (\gamma_i + \gamma_j) d(u_i, u_j | T) = \sum_{e \in E(T)} [4n_1 n_2 - (n_1 + n_2)] \quad (12)$$

Using the fact that T has $n - 1$ edges, that $n_1 + n_2 = n$ and that $d(T)$ obeys Eq. (6), Eq. (4) is directly obtained from Eq. (12).

This proves the first part of Lemma 1.

If, on the other hand, we choose $\omega_{ij} = \gamma_i \cdot \gamma_j$, then from Eq. (8),

$$\begin{aligned} \sum_{1 \leq i < j \leq n} \gamma_i \gamma_j d(u_i, u_j | T) &= \sum_{e \in E(T)} \left[\sum_{u_i \in V(T_1)} \gamma_i \right] \left[\sum_{u_j \in V(T_2)} \gamma_j \right] \\ &= \sum_{e \in E(T)} [2(n_1 - 1) + 1][2(n_2 - 1) + 1] \end{aligned}$$

Combining the above with Eq. (6) leads to Eq. (5).

This completes the proof of Lemma 1. \square

Proof of Corollary 1.2. Set $p_i = \gamma_i$ into Eq. (1) and use Eqs. (4) and (5) of Lemma 1. \square

Proof of Corollary 1.3. Set $p_i = \gamma - \gamma_i$ into Eq. (1) and use Lemma 1. The calculation is somewhat lengthier than, but fully analogous to the previous case. \square

REFERENCES

- [1] A. Cayley, *On the mathematical theory of isomers*, Phil. Magazine **47** (1874), 444–446.
- [2] R. C. Entringer, D. E. Jackson, D. A. Snyder, *Distance in graphs*, Czech. Math. J. **26** (1976), 283–296.
- [3] I. Gutman, *On the distance of some compound graphs*, Publ. Elektrotehn. Fak. (Beograd) Ser. Mat. **5** (1994), 29–34.
- [4] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986, pp. 19–20.
- [5] F. Harary, A. J. Schwenk, *The number of caterpillars*, Discrete Math. **6** (1973), 359–365.
- [6] J. Plesnik, *On the sum of all distances in a graph or digraph*, J. Graph Theory **8** (1984), 1–21.
- [7] G. Pólya, *Kombinatorische Anzahlbestimmungen für Gruppen, Graphen und chemische Verbindungen*, Acta Math. **68** (1937), 145–254, sections 35–37, 55.
- [8] H. Wiener, *Structural determination of paraffin boiling points*, J. Amer. Chem. Soc. **69** (1947), 17–20.
- [9] Y. N. Yeh, I. Gutman, *On the sum of all distances in composite graphs*, Discrete Math. **135** (1994), 359–365.

Prirodno-matematički fakultet
Univerzitet u Kragujevcu
34000 Kragujevac, p.p. 60
Yugoslavia

(Received 19 08 1997)