

FINITE DIFFERENCE APPROXIMATION OF A PARABOLIC PROBLEM WITH VARIABLE COEFFICIENTS

Boško S. Jovanović and Zorica Milovanović

Communicated by Gradimir Milovanović

ABSTRACT. We study the convergence of a finite difference scheme that approximates the third initial-boundary-value problem for a parabolic equation with variable coefficients on a unit square. We assume that the generalized solution of the problem belongs to the Sobolev space $W_2^{s,s/2}$, $s \leq 3$. An almost second-order convergence rate estimate (with additional logarithmic factor) in the discrete $W_2^{1,1/2}$ norm is obtained. The result is based on some nonstandard a priori estimates involving fractional order discrete Sobolev norms.

1. Introduction

For a class of finite difference schemes (FDSs) approximating elliptic boundary-value problems (BVPs) with generalized solutions, convergence rate estimates compatible with the smoothness of the data

$$(1.1) \quad \|u - v\|_{W_p^k(\omega)} \leq Ch^{s-k} \|u\|_{W_p^s(\Omega)}, \quad s \geq k,$$

are of great interest (see [6, 13]). Here $u = u(x)$ denotes the solution of the BVP, v denotes the solution of the corresponding FDS, h is the discretization parameter, $W_p^k(\omega)$ is the Sobolev space of mesh functions, and C is a positive generic constant, independent of h and u . In the parabolic case, instead of (1.1) it is natural to look for error bounds of the form

$$(1.2) \quad \|u - v\|_{W_p^{k,k/2}(Q_{h\tau})} \leq C \left(h^{s-k} + \tau^{\frac{s-k}{2}} \right) \|u\|_{W_p^{s,s/2}(Q)}, \quad s \geq k,$$

where τ is the temporal mesh-size. A standard technique for establishing estimates of such types (see [6, 13, 14]) is based on the Bramble–Hilbert lemma [3, 5].

For BVPs with an oblique derivative boundary condition a loss of one half of an order in the convergence rate (usually $O(h^{3/2})$) is often observed, caused by

2010 *Mathematics Subject Classification*: 65M15.

Key words and phrases: parabolic initial-boundary-value problem, oblique derivative boundary condition, finite differences, Sobolev spaces, convergence rate estimates.

The research was supported by Ministry of Education and Science of Republic of Serbia under project 174015.

the approximation of the boundary condition. Nevertheless, improved results are obtained in some cases, mainly for elliptic problems (see [4, 7]).

In the present paper, for the FDS approximating initial-boundary value problem (IBVP) for a parabolic equation with variable coefficients and an oblique derivative boundary condition a second order error bound in the discrete $W_2^{1,1/2}$ norm is obtained under minimal smoothness assumptions on the input data. The result is based on some nonstandard a priori estimates involving fractional order discrete Sobolev norms.

2. Formulation of the Problem

As the model problem we consider, in $Q = \Omega \times (0, T) = (0, 1)^2 \times (0, T)$, the following initial-boundary value problem for a parabolic equation with variable coefficients:

$$(2.1) \quad \frac{\partial u}{\partial t} + Lu = f, \quad (x, t) = (x_1, x_2, t) \in Q,$$

$$(2.2) \quad lu = 0, \quad (x, t) \in \Gamma \times (0, T) = \partial\Omega \times (0, T),$$

$$(2.3) \quad u(x, 0) = u_0(x), \quad x \in \Omega.$$

where

$$(2.4) \quad Lu := - \sum_{i,j=1}^2 \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial u}{\partial x_j} \right), \quad lu := \sum_{i,j=1}^2 a_{ij} \frac{\partial u}{\partial x_j} \cos(\nu, x_i) + \alpha u$$

and ν is the unit outward normal to Γ . We assume that the conditions of strong ellipticity are satisfied:

$$(2.5) \quad \begin{aligned} a_{ij} &= a_{ij}(x) = a_{ji}, & \alpha &= \alpha(x), \\ c_0 \sum_{i=1}^2 \xi_i^2 &\leq \sum_{i,j=1}^2 a_{ij} \xi_i \xi_j \leq c_1 \sum_{i=1}^2 \xi_i^2, & \forall x \in \bar{\Omega}, \forall \xi \in \mathbb{R}^2, & c_i = \text{const.} > 0, \\ \alpha_0 &\leq \alpha(x) \leq \alpha_1, & \alpha_i &= \text{const.} > 0. \end{aligned}$$

Let us denote $\Gamma = \bigcup_{i=1}^2 \bigcup_{k=0}^1 \Gamma_{ik}$, where $\Gamma_{1k} = \{k\} \times [0, 1]$, $\Gamma_{2k} = [0, 1] \times \{k\}$, and $\Sigma_{ik} = \Gamma_{ik} \times [0, T]$.

We also assume that the generalized solution of the problem (2.1)–(2.3) belongs to the Sobolev space $W_2^{s,s/2}(Q)$, $2 < s \leq 3$, while the data satisfy the following smoothness conditions: $a_{ij} \in W_2^{s-1}(\Omega)$, $\alpha \in W_2^{s-3/2}(\Gamma_{ik})$, $\alpha \in C(\Gamma)$, $f \in W_2^{s-2,s/2-1}(Q)$ and $u_0 \in W_2^{s-1}(\Omega)$.

3. Finite Difference Approximation

Let $n, m \in \mathbb{N}$, $n \geq 2$, $m \geq 1$, $h = 1/n$ and $\tau = T/m$. We consider the uniform spatial mesh $\bar{\omega}$ with mesh size h on $\bar{\Omega}$ and the uniform temporal mesh $\bar{\omega}_\tau$ with mesh size τ on $[0, T]$. We also denote $\omega = \bar{\omega} \cap \Omega$, $\omega_\tau = \bar{\omega}_\tau \cap (0, T)$, $\omega_\tau^- = \bar{\omega}_\tau \cap [0, T)$, $\omega_\tau^+ = \bar{\omega}_\tau \cap (0, T]$, $\gamma = \bar{\omega} \cap \Gamma$, $\bar{\gamma}_{ik} = \bar{\omega} \cap \Gamma_{ik}$, $\gamma_{ik} = \{x \in \bar{\gamma}_{ik} : 0 < x_{3-i} < 1\}$, $\gamma_{ik}^- = \{x \in \bar{\gamma}_{ik} : 0 \leq x_{3-i} < 1\}$, $\gamma_{ik}^+ = \{x \in \bar{\gamma}_{ik} : 0 < x_{3-i} \leq 1\}$, $\gamma_{ik}^* = \bar{\gamma}_{ik} \setminus \gamma_{ik}$,

$\gamma^* = \gamma \setminus \{\cup_{i,k} \gamma_{ik}\}$, $\sigma_{ik} = \gamma_{ik} \times \omega_\tau^+$, $\bar{\sigma}_{ik} = \bar{\gamma}_{ik} \times \omega_\tau^+$, $i = 1, 2$, $k = 0, 1$, and $\bar{Q}_{h\tau} = \bar{\omega} \times \bar{\omega}_\tau$.

The finite difference operators are defined in the usual manner [12]:

$$v_{x_i} = (v^{+i} - v)/h, \quad v_{\bar{x}_i} = (v - v^{-i})/h, \quad v_t = (\hat{v} - v)/\tau, \quad v_{\bar{t}} = (v - \check{v})/\tau,$$

where $v^{\pm i}(x, t) = v(x \pm h e_i, t)$, e_i is the unit vector of the axis x_i , $\hat{v}(x, t) = v(x, t + \tau)$ and $\check{v}(x, t) = v(x, t - \tau)$.

We also define the Steklov smoothing operators with the step sizes h and τ [13]:

$$T_i^+ f(x, t) = \int_0^1 f(x + h x' e_i, t) dx' = T_i^- f(x + h e_i, t) = T_i f(x + \frac{h}{2} e_i, t),$$

$$T_i^{2\pm} f(x, t) = 2 \int_0^1 (1 - x') f(x \pm h x' e_i) dx', \quad i = 1, 2,$$

$$T_t^+ f(x, t) = \int_0^1 f(x, t + \tau t') dt' = T_t^- f(x, t + \tau) = T_t f(x, t + \frac{\tau}{2}).$$

These operators commute and transform derivatives into differences, for example:

$$\begin{aligned} T_i^+ \left(\frac{\partial u}{\partial x_i} \right) &= u_{x_i}, & T_i^- \left(\frac{\partial u}{\partial x_i} \right) &= u_{\bar{x}_i}, & T_i^2 \left(\frac{\partial^2 u}{\partial x_i^2} \right) &= u_{\bar{x}_i x_i}, & i = 1, 2, \\ T_t^+ \left(\frac{\partial u}{\partial t} \right) &= u_t, & T_t^- \left(\frac{\partial u}{\partial t} \right) &= u_{\bar{t}}. \end{aligned}$$

We approximate the IBVP (2.1)–(2.3) with the following implicit FDS:

$$(3.1) \quad \begin{aligned} v_{\bar{t}} + L_h v &= \tilde{f}, & x &\in \bar{\omega}, & t &\in \omega_\tau^+, \\ v(x, 0) &= u_0(x), & x &\in \bar{\omega}, \end{aligned}$$

where

$$L_h v = \begin{cases} -\frac{1}{2} \sum_{i,j=1}^2 \left[(a_{ij} v_{x_j})_{\bar{x}_i} + (a_{ij} v_{\bar{x}_j})_{x_i} \right], & x \in \omega \\ \frac{2}{h} \left(-\frac{a_{11} + a_{11}^+}{2} v_{x_1} - a_{12} \frac{v_{x_2} + v_{\bar{x}_2}}{2} + \tilde{\alpha} v \right) - (a_{12} v_{\bar{x}_2})_{x_1} \\ \quad - (a_{21} v_{x_1})_{\bar{x}_2} - \frac{1}{2} (a_{22} v_{x_2})_{\bar{x}_2} - \frac{1}{2} (a_{22} v_{\bar{x}_2})_{x_2}, & x \in \gamma_{10} \\ \frac{2}{h} \left[-\frac{a_{11} + a_{11}^+}{2} v_{x_1} - a_{12} v_{x_2} - a_{21} v_{x_1} - \frac{a_{22} + a_{22}^+}{2} v_{x_2} \right. \\ \quad \left. + (\tilde{\alpha}_1 + \tilde{\alpha}_2) v \right], & x = (0, 0) \\ \frac{2}{h} \left[-\frac{a_{11} + a_{11}^+}{2} v_{x_1} - a_{12} v_{\bar{x}_2} + a_{21} v_{x_1} + \frac{a_{22} + a_{22}^-}{2} v_{\bar{x}_2} \right. \\ \quad \left. + (\tilde{\alpha}_1 + \tilde{\alpha}_2) v \right] - 2 (a_{12} v_{\bar{x}_2})_{x_1} - 2 (a_{21} v_{x_1})_{\bar{x}_2}, & x = (0, 1) \\ \text{and analogously at the other boundary nodes,} & x \in \gamma \setminus \bar{\gamma}_{10} \end{cases}$$

$$\tilde{f} = \begin{cases} T_1^2 T_2^2 T_t^- f, & x \in \omega \\ T_i^{2\pm} T_{3-i}^2 T_t^- f, & x \in \gamma_{i, 0.5 \mp 0.5} \\ T_1^{2\pm} T_2^{2\pm} T_t^- f, & x = (0.5 \mp 0.5, 0.5 \mp 0.5) \in \gamma^* \end{cases}$$

$$\tilde{\alpha} = T_{3-i}^2 \alpha, \quad x \in \gamma_{i0} \cup \gamma_{i1}, \quad i = 1, 2$$

and

$$\tilde{\alpha}_i = T_i^{2\pm} \alpha, \quad x \in \gamma^*, \quad x_i = 0.5 \mp 0.5, \quad i = 1, 2.$$

4. Error Analysis

Let u be the solution of the IBVP (2.1)-(2.3), and let v denote the solution of the FDS (3.1). The error $z = u - v$ is defined on $\bar{Q}_{h\tau}$ and satisfies the following conditions

$$(4.1) \quad \begin{aligned} z_{\bar{t}} + L_h z &= \psi, & x \in \bar{\omega}, \quad t \in \omega_{\tau}^+, \\ z(x, 0) &= 0, & x \in \bar{\omega}, \end{aligned}$$

where

$$\psi = \begin{cases} \xi_{\bar{t}} + \sum_{i,j=1}^2 \eta_{ij, \bar{x}_i}, & x \in \omega \\ \tilde{\xi}_{\bar{t}} + \frac{2}{h} \eta_{11} + \frac{2}{h} \eta_{12} + \tilde{\eta}_{21, \bar{x}_2} + \tilde{\eta}_{22, \bar{x}_2} + \frac{2}{h} \zeta, & x \in \gamma_{10} \\ \tilde{\tilde{\xi}}_{\bar{t}} + \frac{2}{h} \tilde{\eta}_{11} + \frac{2}{h} \tilde{\eta}_{12} + \frac{2}{h} \tilde{\eta}_{21} + \frac{2}{h} \tilde{\eta}_{22} + \frac{2}{h} (\zeta_1 + \zeta_2), & x = (0, 0) \\ \text{and analogously at the other boundary nodes,} & x \in \gamma \setminus \gamma_{10}^- \end{cases}$$

$$\xi = u - T_1^2 T_2^2 u, \quad x \in \omega$$

$$\tilde{\xi} = u - T_i^2 T_{3-i}^{2\pm} u, \quad x \in \gamma_{3-i, 0.5 \mp 0.5}$$

$$\tilde{\tilde{\xi}} = u - T_1^{2\pm} T_2^{2\pm} u, \quad x = (0.5 \mp 0.5, 0.5 \mp 0.5) \in \gamma^*$$

$$\eta_{ij} = T_i^+ T_{3-i}^2 T_t^- \left(a_{ij} \frac{\partial u}{\partial x_j} \right) - \frac{1}{2} (a_{ij} u_{x_j} + a_{ij}^{+i} u_{\bar{x}_j}^+), \quad x \in \omega$$

$$\tilde{\eta}_{ii} = T_i^+ T_{3-i}^{2\pm} T_t^- \left(a_{ii} \frac{\partial u}{\partial x_i} \right) - \frac{a_{ii} + a_{ii}^{+i}}{2} u_{x_i}, \quad x \in \gamma_{3-i, 0.5 \mp 0.5}^-$$

$$\tilde{\eta}_{i,3-i} = \begin{cases} T_i^+ T_{3-i}^{2+} T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) - a_{i,3-i} u_{x_{3-i}}, & x \in \gamma_{3-i,0}^- \\ T_i^+ T_{3-i}^{2-} T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) - a_{i,3-i}^{+i} u_{\bar{x}_{3-i}}^+, & x \in \gamma_{3-i,1}^- \end{cases}$$

$$\zeta = (T_i^2 \alpha)u - T_i^2 T_t^-(\alpha u), \quad x \in \gamma_{3-i,0} \cup \gamma_{3-i,1}$$

$$\zeta_i = (T_i^{2\pm} \alpha)u - T_i^{2\pm} T_t^-(\alpha u), \quad x \in \gamma^*, \quad x_i = 0.5 \mp 0.5.$$

We define the following discrete inner products and norms:

$$[v, w] = h^2 \sum_{x \in \omega} v(x)w(x) + \frac{h^2}{2} \sum_{x \in \gamma \setminus \gamma^*} v(x)w(x) + \frac{h^2}{4} \sum_{x \in \gamma^*} v(x)w(x), \quad |[v]|^2 = [v, v],$$

$$[v, w]_i = h^2 \sum_{x \in \omega \cup \gamma_{i0}} v(x)w(x) + \frac{h^2}{2} \sum_{x \in \gamma_{3-i,0}^- \cup \gamma_{3-i,1}^-} v(x)w(x), \quad |[v]_i|^2 = [v, v]_i,$$

$$(v, w)_i = h^2 \sum_{x \in \omega \cup \gamma_{i1}} v(x)w(x) + \frac{h^2}{2} \sum_{x \in \gamma_{3-i,0}^+ \cup \gamma_{3-i,1}^+} v(x)w(x), \quad \|(v, v)_i\|^2 = (v, v)_i,$$

$$[v, w] = h^2 \sum_{x \in \omega \cup \gamma_{10}^- \cup \gamma_{20}^-} v(x)w(x), \quad |[v]|^2 = [v, v], \quad (v, w) = h^2 \sum_{x \in \omega \cup \gamma_{11}^+ \cup \gamma_{21}^+} v(x)w(x),$$

$$\|[v]\|^2 = (v, v), \quad \|[v]\|_{W_2^1(\bar{\omega})}^2 = |[v]|^2 + \|[v_{x_1}]\|_1^2 + \|[v_{x_2}]\|_2^2, \quad \|[v]\|_{C(\bar{\omega})} = \max_{x \in \bar{\omega}} |v(x)|,$$

$$[v, w]_{\bar{\gamma}_{ik}} = h \sum_{x \in \gamma_{ik}} v(x)w(x) + \frac{h}{2} \sum_{x \in \gamma_{ik}^*} v(x)w(x), \quad \|[v]\|_{\bar{\gamma}_{ik}}^2 = [v, v]_{\bar{\gamma}_{ik}},$$

$$[v, w]_{\gamma_{ik}^-} = h \sum_{x \in \gamma_{ik}^-} v(x)w(x), \quad \|[v]\|_{\gamma_{ik}^-}^2 = [v, v]_{\gamma_{ik}^-}, \quad \|v\|_{\gamma_{ik}}^2 = h \sum_{x \in \gamma_{ik}} v^2(x),$$

$$|v|_{W_2^{1/2}(\gamma_{ik}^-)}^2 = h^2 \sum_{x, x' \in \gamma_{ik}^-, x' \neq x} \left[\frac{v(x) - v(x')}{x_{3-i} - x'_{3-i}} \right]^2, \quad \|v\|_{W_2^{1/2}(\gamma_{ik}^-)}^2 = |v|_{W_2^{1/2}(\gamma_{ik}^-)}^2 + \|v\|_{\gamma_{ik}^-}^2,$$

$$\|[v]\|_{\tilde{W}_2^{1/2}(\gamma_{ik}^-)}^2 = |v|_{W_2^{1/2}(\gamma_{ik}^-)}^2 + h \sum_{x \in \gamma_{ik}^-} \left(\frac{1}{x_{3-i} + h/2} + \frac{1}{1 - x_{3-i} - h/2} \right) v^2(x),$$

$$\|v\|_{\tau}^2 = \tau \sum_{t \in \omega_{\tau}^+} v^2(t), \quad \|[v]\|_{i, h\tau}^2 = \tau \sum_{t \in \omega_{\tau}^+} \|[v(\cdot, t)]\|_i^2,$$

$$\|v\|_{\sigma_{ik}}^2 = \tau \sum_{t \in \omega_{\tau}^+} \|v(\cdot, t)\|_{\gamma_{ik}}^2, \quad \|[v]\|_{\sigma_{ik}}^2 = \tau \sum_{t \in \omega_{\tau}^+} \|[v(\cdot, t)]\|_{\bar{\gamma}_{ik}}^2,$$

$$\|[v]\|_{L_2(\omega_{\tau}^+, \tilde{W}_2^{1/2}(\gamma_{ik}^-))}^2 = \tau \sum_{t \in \omega_{\tau}^+} \|[v(\cdot, t)]\|_{\tilde{W}_2^{1/2}(\gamma_{ik}^-)}^2,$$

$$|v|_{W_2^{1/2}(\bar{\omega}_{\tau}, L_2(\bar{\omega}))}^2 = \tau^2 \sum_{t, t' \in \bar{\omega}_{\tau}, t' \neq t} \frac{|v(\cdot, t) - v(\cdot, t')|^2}{(t - t')^2},$$

$$\|[v]\|_{\tilde{W}_2^{1/2}(\bar{\omega}_{\tau}, L_2(\bar{\omega}))}^2 = |v|_{W_2^{1/2}(\bar{\omega}_{\tau}, L_2(\bar{\omega}))}^2 + \tau \sum_{t \in \bar{\omega}_{\tau}} \left(\frac{1}{t + \tau/2} + \frac{1}{T - t + \tau/2} \right) \|[v(\cdot, t)]\|^2,$$

$$\|[v]\|_{L_2(\omega_{\tau}^+, W_2^1(\bar{\omega}))}^2 = \tau \sum_{t \in \omega_{\tau}^+} \|[v(\cdot, t)]\|_{W_2^1(\bar{\omega})}^2,$$

$$\|[v]\|_{W_2^{1,1/2}(Q_{h\tau})}^2 = \|[v]\|_{L_2(\omega_{\tau}^+, W_2^1(\bar{\omega}))}^2 + |v|_{W_2^{1/2}(\bar{\omega}_{\tau}, L_2(\bar{\omega}))}^2.$$

We shall prove a suitable a priori estimate for the FDS (4.1) which will be used to estimate its convergence rate.

LEMMA 4.1. *Let a_{ij} and α satisfy the assumptions from Section 2. Then, for a sufficiently small mesh step h , there exist positive constants C_1 and C_2 such that*

$$C_1 \|v\|_{W_2^1(\bar{\omega})}^2 \leq [L_h v, v] \leq C_2 \|v\|_{W_2^1(\bar{\omega})}^2.$$

PROOF. The proof immediately follows from

$$\begin{aligned} [L_h v, v] &= \frac{1}{2} \sum_{i=1}^2 \left\{ [a_{ii} v_{x_i}, v_{x_i}]_i + (a_{ii} v_{\bar{x}_i}, v_{\bar{x}_i}]_i + [a_{i,3-i} v_{x_{3-i}}, v_{x_i}] \right. \\ &\quad \left. + (a_{i,3-i} v_{\bar{x}_{3-i}}, v_{\bar{x}_i}] \right\} + h \sum_{x \in \gamma \setminus \gamma^*} \tilde{\alpha} v^2 + \frac{h}{2} \sum_{x \in \gamma^*} (\tilde{\alpha}_1 + \tilde{\alpha}_2) v^2 \\ &= \frac{1}{2} \sum_{i,j=1}^2 \left\{ [a_{ij} v_{x_j}, v_{x_i}] + (a_{ij} v_{\bar{x}_j}, v_{\bar{x}_i}] \right\} + \frac{h}{4} \sum_{i=1}^2 \sum_{k=0}^1 (-1)^k [a_{ii}^{+i} - a_{ii}, v_{x_i}^2]_{\gamma_{3-i,k}^-} \\ &\quad + \sum_{x \in \gamma \setminus \gamma^*} \tilde{\alpha} v^2 + \frac{h}{2} \sum_{x \in \gamma^*} (\tilde{\alpha}_1 + \tilde{\alpha}_2) v^2 \end{aligned}$$

and a discrete imbedding theorem [12]. \square

LEMMA 4.2. [2, 7] *The following inequality holds true:*

$$|[v, w_{x_{3-i}}]_{\gamma_{ik}^-}| \leq C \|v\|_{\tilde{W}_2^{1/2}(\gamma_{ik}^-)} \|w\|_{W_2^1(\bar{\omega})}.$$

LEMMA 4.3. [7] *Let v be a mesh function on $\bar{\omega}$, then*

$$\|v\|_{C(\bar{\omega})} \leq C \sqrt{\log \frac{1}{h}} \|v\|_{W_2^1(\bar{\omega})}.$$

LEMMA 4.4. [8] *The solution of the FDS*

$$v_{\bar{t}} + L_h v = \varphi, \quad (x, t) \in \bar{\omega} \times \omega_{\tau}^+; \quad v(x, 0) = 0, \quad x \in \bar{\omega}.$$

satisfies the a priori estimate

$$\|v\|_{W_2^{1,1/2}(Q_{h\tau})} \leq C \left(\tau \sum_{t \in \omega_{\tau}^+} \|[\varphi(\cdot, t)]\|_{-1}^2 \right)^{1/2}$$

where

$$\|[\varphi(\cdot, t)]\|_{-1} := \sup_w \frac{[\varphi(\cdot, t), w]}{\|w\|_{W_2^1(\bar{\omega})}}.$$

LEMMA 4.5. [8] *The solution of the FDS*

$$v_{\bar{t}} + L_h v = \phi_{\bar{t}}, \quad (x, t) \in \bar{\omega} \times \omega_{\tau}^+; \quad v(x, 0) = 0, \quad x \in \bar{\omega}.$$

satisfies the a priori estimate

$$\|v\|_{W_2^{1,1/2}(Q_{h\tau})} \leq C \|\phi\|_{\tilde{W}_2^{1/2}(\bar{\omega}_{\tau}, L_2(\bar{\omega}))}.$$

LEMMA 4.6. *Let $w \in W_2^r(\Gamma_{ik})$, $0 < r \leq 0.5$. Then*

$$|T_{3-i}^+ w|_{W_2^{1/2}(\gamma_{ik}^-)} \leq C(r) h^{r-1/2} |w|_{W_2^r(\Gamma_{ik})}.$$

PROOF. Without loss of generality let us set $i = 1$ and $k = 0$. Hence

$$\begin{aligned} |T_2^+ w|_{W_2^{1/2}(\gamma_{10}^-)}^2 &= h^2 \sum_{i=1}^{n-1} \sum_{j=1, j \neq i}^{n-1} \frac{[T_2^+ w(0, ih) - T_2^+ w(0, jh)]^2}{(ih - jh)^2} \\ &= 2h^2 \sum_{i=1}^{n-1} \sum_{j=1}^{i-1} \frac{[T_2^+ w(0, ih) - T_2^+ w(0, jh)]^2}{(ih - jh)^2} \\ &= 2h^2 \sum_{i=1}^{n-1} \sum_{j=1}^{i-1} \left\{ \frac{1}{h^2} \int_{ih}^{ih+h} \int_{jh}^{jh+h} [w(0, x) - w(0, x')] dx' dx \right\}^2 (ih - jh)^{-2} \\ &= 2h^2 \sum_{i=1}^{n-1} \sum_{j=1}^{i-1} \frac{1}{h^4} \left\{ \int_{ih}^{ih+h} \int_{jh}^{jh+h} \frac{[w(0, x) - w(0, x')]^2}{(x - x')^{1+2r}} dx' dx \right\} \\ &\quad \times \left\{ \int_{ih}^{ih+h} \int_{jh}^{jh+h} \frac{(x - x')^{1+2r}}{(ih - jh)^2} dx' dx \right\} \\ &\leq 2h^{-2} 2^{1+2r} h^2 h^{2r-1} \sum_{i=1}^{n-1} \sum_{j=1}^{i-1} \left\{ \int_{ih}^{ih+h} \int_{jh}^{jh+h} \frac{[w(0, x) - w(0, x')]^2}{(x - x')^{1+2r}} dx' dx \right\} \\ &= 2^{1+2r} h^{2r-1} 2 \int_0^1 \int_0^x \frac{[w(0, x) - w(0, x')]^2}{(x - x')^{1+2r}} dx' dx = 2^{1+2r} h^{2r-1} |w|_{W_2^r(\Gamma_{10})}^2. \end{aligned}$$

□

Let us rearrange the summands in truncation error ψ in the following manner:

$$\tilde{\eta}_{ij} = \eta_{ij} + \eta'_{ij}, \quad \tilde{\xi} = \xi + \xi', \quad \tilde{\xi}^* = \xi + \xi^*,$$

where

$$\begin{aligned} \eta'_{ii} &= \pm \frac{h}{3} T_i^+ T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right), \quad x \in \gamma_{3-i, 0.5 \mp 0.5}, \\ \eta'_{i, 3-i} &= \pm \frac{h}{3} T_i^+ T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{i, 3-i} \frac{\partial u}{\partial x_{3-i}} \right) \right) \mp \frac{h}{2} T_i^+ T_t^- \left(a_{i, 3-i} \frac{\partial^2 u}{\partial x_{3-i}^2} \right) \\ &\quad + \frac{h}{2} T_i^+ T_t^- \left(\frac{\partial}{\partial x_i} \left(a_{i, 3-i} \frac{\partial u}{\partial x_{3-i}} \right) \right), \quad x \in \gamma_{3-i, 0.5 \mp 0.5}, \\ \xi' &= \mp \frac{h}{3} T_{3-i}^2 \left(\frac{\partial u}{\partial x_i} \right), \quad x \in \gamma_i, 0.5 \mp 0.5, \\ \xi^* &= \mp \frac{h}{3} T_2^{2\pm} \left(\frac{\partial u}{\partial x_1} \right) \mp \frac{h}{3} T_1^{2\pm} \left(\frac{\partial u}{\partial x_2} \right), \quad x = (0.5 \mp 0.5, 0.5 \mp 0.5) \in \gamma^*. \end{aligned}$$

Using the boundary condition (2.2) we further obtain

$$\xi_{\tilde{i}}^l = \lambda_i, \bar{x}_{3-i} + \mu_i + \nu_i, \quad x \in \gamma_i, 0.5 \mp 0.5,$$

where

$$\begin{aligned}\lambda_i &= \pm \frac{h}{3} T_{3-i}^+ T_t^- \left(\frac{a_{i,3-i}}{a_{ii}} \frac{\partial u}{\partial t} \right), \\ \mu_i &= \mp \frac{h}{3} T_{3-i}^2 T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(\frac{a_{i,3-i}}{a_{ii}} \right) \frac{\partial u}{\partial t} \right), \\ \nu_i &= -\frac{h}{3} T_{3-i}^2 T_t^- \left(\frac{\alpha}{a_{ii}} \frac{\partial u}{\partial t} \right).\end{aligned}$$

Similarly, for $x = (0, 0)$, we obtain

$$\xi_t^* = \frac{2}{h} \lambda_1 - \frac{2}{h} \lambda_1^* + \mu_1 + \nu_1 + \frac{2}{h} \lambda_2 - \frac{2}{h} \lambda_2^* + \mu_2 + \nu_2,$$

where λ_1 and λ_2 are the same as before and

$$\begin{aligned}\lambda_i^* &= \pm \frac{h}{3} T_t^- \left(\frac{a_{i,3-i}}{a_{ii}} \frac{\partial u}{\partial t} \right), \\ \mu_i &= -\frac{h}{3} T_{3-i}^{2+} T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(\frac{a_{i,3-i}}{a_{ii}} \right) \frac{\partial u}{\partial t} \right), \\ \nu_i &= -\frac{h}{3} T_{3-i}^{2+} T_t^- \left(\frac{\alpha}{a_{ii}} \frac{\partial u}{\partial t} \right),\end{aligned}$$

with an analogous representation at other nodes from γ^* .

Using Lemmas 4.1–4.5, we obtain the following assertion.

THEOREM 4.1. *The finite difference scheme (4.1) is stable in the sense of the a priori estimate*

$$\begin{aligned}(4.2) \quad & \| [z] \|_{W_2^{1,1/2}(Q_{h\tau})} \leq C \left\{ \| [\xi] \|_{\bar{W}_2^{1/2}(\bar{\omega}_\tau, L_2(\bar{\omega}))} + \sum_{i,j=1}^2 \| [\eta_{ij}] \|_{i,h\tau} + \sum_{k=0}^1 \sum_{i=1}^2 \| \zeta \|_{\sigma_{ik}} \right. \\ & + h \sum_{k=0}^1 \sum_{i,j=1}^2 \| [\eta'_{ij}] \|_{L_2(\omega_\tau^+, \bar{W}_2^{1/2}(\gamma_{3-i,k}^-))} + h \sum_{k=0}^1 \sum_{i=1}^2 \| [\lambda_i] \|_{L_2(\omega_\tau^+, \bar{W}_2^{1/2}(\gamma_{ik}^-))} \\ & \left. + h \sum_{k=0}^1 \sum_{i=1}^2 \left(\| [\mu_i] \|_{\bar{\sigma}_{ik}} + \| [\nu_i] \|_{\bar{\sigma}_{ik}} \right) + h \sqrt{\log \frac{1}{h}} \sum_{i=1}^2 \sum_{x \in \gamma^*} \left(\| \zeta_i(x, \cdot) \|_\tau + \| \lambda_i^*(x, \cdot) \|_\tau \right) \right\}.\end{aligned}$$

In accordance with Theorem 4.1, the problem of deriving the convergence rate estimate for the FDS (3.1) is reduced to estimating the right-hand side terms in the inequality (4.2).

Let us assume that $\tau \asymp h^2$, i.e., $c_2 h^2 \leq \tau \leq c_3 h^2$ for some positive constants c_2 and c_3 .

The term η_{ij} at the internal mesh nodes can be estimated in the same manner as in the case of the Dirichlet IBVP (see [6]):

$$(4.3) \quad \tau \sum_{t \in \omega_\tau^+} h^2 \sum_{x \in \omega \cup \gamma_{i0}} \eta_{ij}^2 \leq C h^{2s-2} \| a_{ij} \|_{W_2^{s-1}(\Omega)}^2 \| u \|_{W_2^{s,s/2}(Q)}^2, \quad 2 < s \leq 3.$$

In boundary nodes η_{ii} can be decomposed in the following manner

$$\eta_{ii} = \eta_{ii,1} + \eta_{ii,2} + \eta_{ii,3} + \eta_{ii,4}, \quad x \in \gamma_{3-i, 0.5 \mp 0.5}^-,$$

where

$$\begin{aligned}\eta_{ii,1} &= T_i^+ T_t^- \left(a_{ii} \frac{\partial u}{\partial x_i} \right) - (T_i^+ a_{ii}) \left(T_i^+ T_t^- \frac{\partial u}{\partial x_i} \right), \\ \eta_{ii,2} &= \left[(T_i^+ a_{ii}) - \frac{a_{ii} + a_{ii}^{+i}}{2} \right] \left(T_i^+ T_t^- \frac{\partial u}{\partial x_i} \right), \\ \eta_{ii,3} &= \frac{a_{ii} + a_{ii}^{+i}}{2} \left[\left(T_i^+ T_t^- \frac{\partial u}{\partial x_i} \right) - \left(T_i^+ \frac{\partial u}{\partial x_i} \right) \right], \\ \eta_{ii,4} &= T_i^+ T_{3-i}^{2\pm} T_t^- \left(a_{ii} \frac{\partial u}{\partial x_i} \right) - T_i^+ T_t^- \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \mp \frac{h}{3} T_i^+ T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right).\end{aligned}$$

The terms $\eta_{ii,l}$ for $l = 1, 2, 3$ satisfy the same conditions as analogous terms in [6] whereby it follows that

$$(4.4) \quad \tau \sum_{t \in \omega_\tau^+} h^2 \sum_{x \in \gamma_{3-i,0}^- \cup \gamma_{3-i,1}^-} \eta_{ii,l}^2 \leq Ch^{2s-2} \|a_{ii}\|_{W_2^{s-1}(\Omega)}^2 \|u\|_{W_2^{s,s/2}(Q)}^2, \quad 2 < s \leq 3.$$

For $s > 2.5$ the term $\eta_{ii,4}$ is a bounded linear functional of $w = a_{ii} \frac{\partial u}{\partial x_i} \in W_2^{s-1,(s-1)/2}$ which vanishes when $w = 1, x_1, x_2, t$. Using the Bramble–Hilbert lemma [3, 5] and properties of multipliers in Sobolev spaces [10] we obtain the following result

$$(4.5) \quad \tau \sum_{t \in \omega_\tau^+} h^2 \sum_{x \in \gamma_{3-i,0}^- \cup \gamma_{3-i,1}^-} \eta_{ii,4}^2 \leq Ch^{2s-2} \left\| a_{ii} \frac{\partial u}{\partial x_i} \right\|_{W_2^{s-1,(s-1)/2}(Q)}^2 \\ \leq Ch^{2s-2} \|a_{ii}\|_{W_2^{s-1}(\Omega)}^2 \|u\|_{W_2^{s,s/2}(Q)}^2, \quad 2.5 < s \leq 3.$$

Similarly, at the boundary nodes $\eta_{i,3-i}$ can be decomposed in the following manner

$$\eta_{i,3-i} = \eta_{i,3-i,1} + \eta_{i,3-i,2} + \eta_{i,3-i,3} + \eta_{i,3-i,4}, \quad x \in \gamma_{3-i,0.5\mp 0.5}^-$$

where

$$\begin{aligned}\eta_{i,3-i,1} &= T_i^+ T_{3-i}^{2\pm} T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) - T_i^+ T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) \\ &\quad \mp \frac{h}{3} T_i^+ T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) \right), \\ \eta_{i,3-i,2} &= T_i^+ T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) - T_t^- \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) \\ &\quad - \frac{h}{2} T_i^+ T_t^- \left(\frac{\partial}{\partial x_i} \left(a_{i,3-i} \frac{\partial u}{\partial x_{3-i}} \right) \right), \\ \eta_{i,3-i,3} &= a_{i,3-i} \left[T_t^- \left(\frac{\partial u}{\partial x_{3-i}} \right) - T_{3-i}^\pm \left(\frac{\partial u}{\partial x_{3-i}} \right) \pm \frac{h}{2} T_i^+ T_t^- \left(\frac{\partial^2 u}{\partial x_{3-i}^2} \right) \right], \\ \eta_{i,3-i,4} &= \pm \frac{h}{2} \left[T_i^+ T_t^- \left(a_{i,3-i} \frac{\partial^2 u}{\partial x_{3-i}^2} \right) - a_{i,3-i} T_i^+ T_t^- \left(\frac{\partial^2 u}{\partial x_{3-i}^2} \right) \right].\end{aligned}$$

The terms $\eta_{i,3-i,1}$ and $\eta_{i,3-i,2}$ satisfy estimates analogous to (4.5). For $s > 2.5$ and $a_{i,3-i} \in C(\bar{\Omega})$ the term $\eta_{i,3-i,3}$ is a bounded linear functional of $u \in W_2^{s,s/2}$ which

vanishes when $w = 1, x_1, x_2, t, x_1^2, x_1x_2, x_2^2$. Using the Bramble–Hilbert lemma and the Sobolev imbedding theorem [1] we obtain the following result

$$(4.6) \quad \tau \sum_{t \in \omega_\tau^+} h^2 \sum_{x \in \gamma_{3-i,0}^- \cup \gamma_{3-i,1}^-} \eta_{i,3-i,3}^2 \leq Ch^{2s-2} \|a_{i,3-i}\|_{C(\bar{\Omega})}^2 \|u\|_{W_2^{s,s/2}(Q)}^2 \\ \leq Ch^{2s-2} \|a_{i,3-i}\|_{W_2^{s-1}(\Omega)}^2 \|u\|_{W_2^{s,s/2}(Q)}^2, \quad 2.5 < s \leq 3.$$

Term $\eta_{i,3-i,4}$ can be estimated directly. Let us set $i = 2$ and $x = (0, x_2) \in \gamma_{10}^-$. Then

$$\eta_{2,1,4}(0, x_2, t) = \frac{h}{2} \frac{2}{h\tau} \int_{x_2}^{x_2+h} \left(1 - \frac{x_2'}{h}\right) \int_{t-\tau}^t \int_{x_2}^{x_2'} \frac{\partial a_{21}}{\partial x_2}(0, x_2'') \frac{\partial^2 u}{\partial x_1^2}(0, x_2', t') dx_2'' dt' dx_2',$$

while for other boundary nodes, and also for $i = 1$, we have analogous integral representations. Hence

$$(4.7) \quad \tau \sum_{t \in \omega_\tau^+} h^2 \sum_{x \in \gamma_{3-i,k}^-} \eta_{i,3-i,4}^2 \leq Ch^4 \left\| \frac{\partial a_{i,3-i}}{\partial x_i} \right\|_{L_2(\Gamma_{ik})}^2 \left\| \frac{\partial^2 u}{\partial x_1^2} \right\|_{L_2(\Sigma_{ik})}^2 \\ \leq Ch^4 \|a_{i,3-i}\|_{W_2^{s-1}(\Omega)}^2 \|u\|_{W_2^{s,s/2}(Q)}^2, \quad s > 2.5.$$

In such a way, from (4.3)-(4.7) one obtains

$$(4.8) \quad \|\eta_{ij}\|_{i,h\tau} \leq Ch^{s-1} \|a_{ij}\|_{W_2^{s-1}(\Omega)} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2.5 < s \leq 3.$$

The term ξ at the internal mesh nodes is estimated in [6]. At the boundary nodes ξ admits an analogous integral representation as for $x \in \omega$. Hence, one immediately obtains

$$(4.9) \quad \|\xi\|_{\bar{W}_2^{1/2}(\bar{\omega}_\tau, L_2(\bar{\omega}))} \leq Ch^{s-1} \sqrt{\log \frac{1}{h}} \|a_{ij}\|_{W_2^{s-1}(\Omega)} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2 < s \leq 3.$$

The term ζ on $\sigma_{3-i,k}$ can be represented in the following manner:

$$\zeta = (T_i^2 \alpha) (u - T_i^2 T_t^- u) + [(T_i^2 \alpha) (T_i^2 T_t^- u) - T_i^2 (\alpha T_t^- u)] = \zeta_{01} + \zeta_{02}.$$

For $s > 2.5$ and $\alpha \in C(\Gamma_{3-i,k})$ the term ζ_{01} is a bounded linear functional of $u \in W_2^{s-1, (s-1)/2}(\Sigma_{3-i,k})$ which vanishes when $u = 1$ and $u = x_{3-i}$. Using the Bramble–Hilbert lemma and the Sobolev imbedding theorem we obtain the following result

$$(4.10) \quad \|\zeta_{01}\|_{\sigma_{3-i,k}} \leq Ch^{s-1} \|\alpha\|_{C(\Gamma_{3-i,k})} \|u\|_{W_2^{s-1, (s-1)/2}(\Sigma_{3-i,k})} \\ \leq Ch^{s-1} \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2.5 < s \leq 3.$$

The term ζ_{02} is a bounded linear functional of $(\alpha, T_t^- u) \in W_q^r(\Gamma_{3-i,k}) \times W_{\frac{2q}{q-2}}^p(\Gamma_{3-i,k})$, $q > 2$, which vanishes when $u = 1$ or $T_t^- u = 1$. Using the bilinear version of the Bramble–Hilbert lemma, the Sobolev imbedding theorem and the Hölder inequality,

after summation over the mesh $\gamma_{3-i,k}$ we obtain the following result

$$\begin{aligned} \|\zeta_{02}\|_{\gamma_{3-i,k}} &\leq Ch^{r+p} \|\alpha\|_{W_q^r(\Gamma_{3-i,k})} \|T_t^- u\|_{W_{2q/(q-2)}^p(\Gamma_{3-i,k})} \\ &\leq Ch^{r+p} \|\alpha\|_{W_2^{r+\frac{1}{2}-\frac{1}{q}}(\Gamma_{3-i,k})} \|T_t^- u\|_{W_2^{p+\frac{1}{q}}(\Gamma_{3-i,k})}, \quad 0 \leq r \leq 1, \quad 1 - \frac{1}{q} < p \leq 1. \end{aligned}$$

Summing over the mesh ω_τ^+ and using imbedding and trace theorems we obtain

$$\begin{aligned} \|\zeta_{02}\|_{\sigma_{3-i,k}} &\leq Ch^{r+p} \|\alpha\|_{W_2^{r+\frac{1}{2}-\frac{1}{q}}(\Gamma_{3-i,k})} \|u\|_{W_2^{p+\frac{1}{q}, \frac{1}{2}(p+\frac{1}{q})}(\Sigma_{3-i,k})} \\ &\leq Ch^{r+p} \|\alpha\|_{W_2^{r+\frac{1}{2}-\frac{1}{q}}(\Gamma_{3-i,k})} \|u\|_{W_2^{p+\frac{1}{2}+\frac{1}{q}, \frac{1}{2}(p+\frac{1}{2}+\frac{1}{q})}(Q)} \\ &\leq Ch^{r+p} \|\alpha\|_{W_2^{r+p-\frac{1}{2}}(\Gamma_{3-i,k})} \|u\|_{W_2^{r+p+1, \frac{1}{2}(r+p+1)}(Q)}. \end{aligned}$$

Finally, setting $r+p=s-1$, we get

$$(4.11) \quad \|\zeta_{02}\|_{\sigma_{3-i,k}} \leq Ch^{s-1} \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2 < s \leq 3.$$

From (4.10) and (4.11) it follows that

$$(4.12) \quad \|\zeta\|_{\sigma_{3-i,k}} \leq Ch^{s-1} \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2.5 < s \leq 3.$$

For $x \in \gamma_{3-i,k}^*$ we set

$$\zeta_i = (T_i^{2\pm} \alpha) (u - T_t^- u) + [(T_i^{2\pm} \alpha) (T_t^- u) - T_i^{2\pm} (\alpha T_t^- u)] = \zeta_{i1} + \zeta_{i2}.$$

For $r > 0.5$ and $\alpha \in C(\Gamma_{3-i,k})$ the term ζ_{i1} is a bounded linear functional of $u \in W_2^r(0, T)$ which vanishes when $u = 1$. Using the Bramble–Hilbert lemma and the Sobolev imbedding theorem, after summation over the mesh ω_τ^+ we obtain

$$\begin{aligned} \|\zeta_{i1}\|_\tau &\leq C\tau^r \|\alpha\|_{C(\Gamma_{3-i,k})} \|u(x, \cdot)\|_{W_2^r(0, T)} \\ &\leq Ch^{2r} \|\alpha\|_{W_2^{2r-1/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{2r+1, r+1/2}(Q)}, \quad 0.5 < r \leq 1, \end{aligned}$$

whereby, setting $2r+1=s$,

$$(4.13) \quad \|\zeta_{i1}\|_\tau \leq Ch^{s-1} \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2 < s \leq 3.$$

For $i=1$ and $x=(0,0)$ we have

$$\begin{aligned} \zeta_{12}(0,0,t) &= \frac{2}{h} \int_0^h \left(1 - \frac{x'_1}{h}\right) \alpha(x'_1, 0) [T_t^- u(0,0,t) - T_t^- u(x'_1,0,t)] dx'_1 \\ &= \frac{2}{h} \int_0^h \int_0^{x'_1} \left(1 - \frac{x'_1}{h}\right) \alpha(x'_1, 0) \frac{\partial(T_t^- u)}{\partial x_1}(x'_1, 0, t) dx''_1 dx'_1. \end{aligned}$$

Analogous representations hold for other nodes $x \in \gamma^*$ and also for $i=2$. Hence

$$|\zeta_{i2}| \leq Ch \|\alpha\|_{C(\Gamma_{3-i,k})} \left\| \frac{\partial(T_t^- u)}{\partial x_i} \right\|_{C(\bar{\Omega})} \leq Ch \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|T_t^- u\|_{W_2^s(\Omega)}, \quad s > 2,$$

whereby, after summation over the mesh ω_τ^+ , one obtains

$$(4.14) \quad \|\zeta_{i2}\|_\tau \leq Ch \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad s > 2.$$

In such a way, for $x \in \gamma_{3-i,k}^*$, from (4.13) and (4.14) we get

$$(4.15) \quad \|\zeta_i\|_\tau \leq Ch \|\alpha\|_{W_2^{s-3/2}(\Gamma_{3-i,k})} \|u\|_{W_2^{s,s/2}(Q)}, \quad s > 2.$$

Let us now estimate $\|\eta'_{ij}\|_{L_2(\omega_\tau^+, \dot{W}_2^{1/2}(\gamma_{3-i,k}^-))}$. Using Lemma 4.6 we immediately obtain

$$\begin{aligned} |\eta'_{ii}|_{W_2^{1/2}(\gamma_{3-i,k}^-)} &\leq Ch^{r+1/2} \left| T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right) \right|_{W_2^r(\Gamma_{3-i,k})} \\ &\leq Ch^{r+1/2} \left\| T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right) \right\|_{W_2^{r+1/2}(\Omega)}, \quad 0 < r \leq 0.5. \end{aligned}$$

Using the inequality [11]

$$\|F\|_{L_2(0,\varepsilon)} \leq C \begin{cases} \varepsilon^r \|F\|_{W_2^r(0,1)}, & 0 < r < 0.5, \\ \varepsilon^{1/2} \log \frac{1}{\varepsilon} \|F\|_{W_2^{1/2}(0,1)}, & r = 0.5, \\ \varepsilon^{1/2} \|F\|_{W_2^{1/2}(0,1)}, & r > 0.5, \end{cases}$$

where $0 < \varepsilon < 1$, we obtain

$$\begin{aligned} h \sum_{x \in \gamma_{3-i,k}^-} \left(\frac{1}{x_i + h/2} + \frac{1}{1 - x_i - h/2} \right) (\eta'_{ii})^2 \\ \leq Ch^{2r+1} \log \frac{1}{h} \left\| T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right) \right\|_{W_2^r(\Gamma_{3-i,k}^-)}^2 \\ \leq Ch^{2r+1} \log \frac{1}{h} \left\| T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right) \right\|_{W_2^{r+1/2}(\Omega)}^2, \quad 0 < r < 0.5 \end{aligned}$$

and

$$\begin{aligned} h \sum_{x \in \gamma_{3-i,k}^-} \left(\frac{1}{x_i + h/2} + \frac{1}{1 - x_i - h/2} \right) (\eta'_{ii})^2 \\ \leq Ch^2 \log^3 \frac{1}{h} \left\| T_t^- \left(\frac{\partial}{\partial x_{3-i}} \left(a_{ii} \frac{\partial u}{\partial x_i} \right) \right) \right\|_{W_2^1(\Omega)}^2. \end{aligned}$$

From the obtained inequalities, summing over the mesh ω_τ^+ , denoting $r + 2.5 = s$ and using properties of multipliers in Sobolev spaces, we immediately obtain

$$(4.16) \quad \|\eta'_{ii}\|_{L_2(\omega_\tau^+, \dot{W}_2^{1/2}(\gamma_{3-i,k}^-))} \leq Ch^{s-2} \sqrt{\log \frac{1}{h}} \|a_{ii}\|_{W_2^{s-1}(\Omega)} \|u\|_{W_2^{s,s/2}(Q)},$$

for $2.5 < s < 3$ and

$$(4.17) \quad \|\eta'_{ii}\|_{L_2(\omega_\tau^+, \dot{W}_2^{1/2}(\gamma_{3-i,k}^-))} \leq Ch \left(\log \frac{1}{h}\right)^{3/2} \|a_{ii}\|_{W_2^2(\Omega)} \|u\|_{W_2^{3,3/2}(Q)}.$$

Analogous estimates hold for $\eta'_{i,3-i}$ substituting a_{ii} with $a_{i,3-i}$.

The term λ_i has a similar structure as η'_{ij} . Handling it in the same manner we obtain for $2.5 < s < 3$

$$\|\lambda_i\|_{L_2(\omega_\tau^+, \dot{W}_2^{1/2}(\gamma_{ik}^-))} \leq Ch^{s-2} \sqrt{\log \frac{1}{h}} \left\| \frac{a_{i,3-i}}{a_{ii}} \right\|_{M(W_2^{s-2}(\Omega))} \|u\|_{W_2^{s,s/2}(Q)},$$

where $\|\cdot\|_{M(W_2^{s-2}(\Omega))}$ is the norm in the space of multipliers in $W_2^{s-2}(\Omega)$ [10]. Further, for $2.5 < s < 3$ and $q > 2$, using the properties of multipliers in Sobolev spaces and the assumptions (2.4), we obtain

$$\begin{aligned} \left\| \frac{a_{i,3-i}}{a_{ii}} \right\|_{M(W_2^{s-2}(\Omega))} &\leq C \left\| \frac{a_{i,3-i}}{a_{ii}} \right\|_{W_q^1(\Omega)} \leq C \left(\|a_{i,3-i}\|_{W_q^1(\Omega)} + \|a_{ii}\|_{W_q^1(\Omega)} \right) \\ &\leq C \left(\|a_{i,3-i}\|_{W_2^{s-1}(\Omega)} + \|a_{ii}\|_{W_2^{s-1}(\Omega)} \right). \end{aligned}$$

In such a way, we finally obtain

$$(4.18) \quad \|\lambda_i\|_{L_2(\omega_\tau^+, \bar{W}_2^{1/2}(\gamma_{ik}^-))} \leq Ch^{s-2} \sqrt{\log \frac{1}{h}} \\ \times \left(\|a_{ii}\|_{W_2^{s-1}(\Omega)} + \|a_{i,3-i}\|_{W_2^{s-1}(\Omega)} \right) \|u\|_{W_2^{s,s/2}(Q)}, \quad 2.5 < s < 3$$

and analogously

$$(4.19) \quad \|\lambda_i\|_{L_2(\omega_\tau^+, \bar{W}_2^{1/2}(\gamma_{ik}^-))} \leq Ch \left(\log \frac{1}{h} \right)^{3/2} \\ \times \left(\|a_{ii}\|_{W_2^2(\Omega)} + \|a_{i,3-i}\|_{W_2^2(\Omega)} \right) \|u\|_{W_2^{3,3/2}(Q)}, \quad s = 3.$$

For $r > 0.5$ and $a_{ii}, a_{i,3-i} \in C(\bar{\Omega})$ the term λ_i^* is a bounded linear functional of $u \in W_2^r(0, T)$ which vanishes when $u = 1$. Using the Bramble–Hilbert lemma and handling it in the same manner as in the case of ζ_{i1} we obtain

$$(4.20) \quad \|\lambda_i^*\|_\tau \leq Ch^{s-2} \|a_{i,3-i}\|_{W_2^{s-1}(\Omega)} \|u\|_{W_2^{s,s/2}(Q)}, \quad 2 < s \leq 3.$$

The terms ν_i and μ_i can be estimated directly:

$$(4.21) \quad \begin{aligned} |[\nu_i]_{\bar{\sigma}_{ik}}| &\leq Ch \left\| \frac{\alpha}{a_{ii}} \frac{\partial u}{\partial t} \right\|_{L_2(\Sigma_{ik})} \leq Ch \left\| \frac{\alpha}{a_{ii}} \right\|_{C(\Gamma_{ik})} \|u\|_{W_2^{s,s/2}(Q)} \\ &\leq Ch \|u\|_{W_2^{s,s/2}(Q)}, \quad s > 2.5 \end{aligned}$$

and

$$(4.22) \quad |[\mu_i]_{\bar{\sigma}_{ik}}| \leq Ch \left(\|a_{ii}\|_{W_2^{s-1}(\Omega)} + \|a_{i,3-i}\|_{W_2^{s-1}(\Omega)} \right) \|u\|_{W_2^{s,s/2}(Q)}, \quad s > 2.5$$

From (4.2), (4.8), (4.9), (4.12) and (4.15)–(4.22) we obtain the following result.

THEOREM 4.2. *Let the assumptions from Section 2 hold and let $\tau \asymp h^2$. Then the FDS (3.1) converges in the norm $W_2^{1,1/2}(Q_{h\tau})$ and the convergence rate estimates*

$$\begin{aligned} |[z]|_{W_2^{1,1/2}(Q_{h\tau})} &\leq Ch^{s-1} \sqrt{\log \frac{1}{h}} \\ &\times \left(1 + \max_{i,j} \|a_{ij}\|_{W_2^{s-1}(\Omega)} + \max_{i,k} \|\alpha\|_{W_2^{s-3/2}(\Gamma_{ik})} \right) \|u\|_{W_2^{s,s/2}(Q)}, \quad 2.5 < s < 3 \end{aligned}$$

and

$$|[z]|_{W_2^{1,1/2}(Q_{h\tau})} \leq Ch^2 \left(\log \frac{1}{h} \right)^{\frac{3}{2}} \left(1 + \max_{i,j} \|a_{ij}\|_{W_2^2(\Omega)} + \max_{i,k} \|\alpha\|_{W_2^{3/2}(\Gamma_{ik})} \right) \|u\|_{W_2^{3,3/2}(Q)}$$

hold.

REMARK 4.1. Note that the obtained error bounds are ‘almost’ compatible with the smoothness of the data (up to additional logarithmic factors).

REMARK 4.2. The obtained error bounds have been derived under the assumption $\tau \asymp h^2$, which links the temporal mesh-size τ to the spatial mesh-size h , despite the fact that the implicit FDS is unconditionally stable, and therefore from the point of view of stability there should be no limitation on the choice of mesh-sizes. This limitation can be avoided in certain cases by careful study of truncation error functionals (see e.g. [9]).

References

- [1] R. A. Adams, J. J. F. Fournier, *Sobolev Spaces*, 2nd ed., Pure Appl. Math. Ser. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [2] V. B. Andreev, *Stability of difference schemes for elliptic equations with respect to the Dirichlet boundary conditions*, Zh. Vychisl. Mat. Mat. Fiz. **12** (1972), 598–611 (in Russian).
- [3] J. H. Bramble, S. R. Hilbert, *Bounds for a class of linear functionals with application to the Hermite interpolation*, Numer. Math. **16** (1971), 362–369.
- [4] M. D. Chkhartishvili, G. K. Berikelashvili, *On the convergence in W_2^1 of difference solution of elliptic equation with mixed boundary conditions*, Bull. Acad. Sci. Georgia **148**:2 (1993), 180–184 (in Russian).
- [5] T. Dupont, R. Scott, *Polynomial approximation of functions in the Sobolev spaces*, Math. Comput. **34** (1980), 441–463.
- [6] B. S. Jovanović, *The Finite Difference Method for Boundary-Value Problems with Weak Solutions*, Posebna izdanja 16, Matematički institut, Beograd, 1993.
- [7] B. S. Jovanović, B. Z. Popović, *Convergence of a finite difference scheme for the third boundary value problem for elliptic equation with variable coefficients*, Comput. Methods Appl. Math. **1**:4 (2001), 356–366.
- [8] B. S. Jovanović, L. G. Vulkov, *On the convergence of finite difference schemes for the heat equation with concentrated capacity*, Numer. Math. **89**:4 (2001), 715–734.
- [9] ———, *Finite difference approximation of strong solutions of a parabolic interface problem on disconnected domains*, Publ. Inst. Math. **84(98)** (2008), 37–48.
- [10] V. G. Maz’ya, T. O. Shaposhnikova, *Theory of Multipliers in Spaces of Differentiable Functions*, Monographs and Studies in Mathematics 23, Pitman, Boston, Mass., 1985.
- [11] L. A. Oganessian, L. A. Rukhovets, *Variational-Difference Methods for Solution of Elliptic Equations*, AS Arm., Erevan, 1979 (in Russian).
- [12] A. A. Samarskiĭ, *The Theory of Difference Schemes*, Marcel Dekker, New York, 2001.
- [13] A. A. Samarskiĭ, R. D. Lazarov, V. L. Makarov, *Difference Schemes for Differential Equations with Generalized Solutions*, Vyshaya Shkola, Moscow, 1987 (in Russian).
- [14] E. E. Süli, B. S. Jovanović, L. D. Ivanović, *On the construction of finite difference schemes approximating generalized solutions*, Publ. Inst. Math. **37(51)** (1985), 123–128.

University of Belgrade
 Faculty of Mathematics
 11000 Belgrade, Serbia
 bosko@matf.bg.ac.rs
 zorica.milovanovic@gmail.com

(Received 23 03 2013)
 (Revised 01 04 2013)