

## EXPLICIT AND ASYMPTOTIC FORMULAE FOR VASYUNIN–COTANGENT SUMS

Mouloud Goubi, Abdelmejid Bayad,  
and Mohand Ouamar Hernane

ABSTRACT. For coprime numbers  $p$  and  $q$ , we consider the Vasyunin–cotangent sum

$$V(q, p) = \sum_{k=1}^{p-1} \left\{ \frac{kq}{p} \right\} \cot \left( \frac{\pi k}{p} \right).$$

First, we prove explicit formula for the symmetric sum  $V(p, q) + V(q, p)$  which is a new reciprocity law for the sums above. This formula can be seen as a complement to the Bettin–Conrey result [13, Theorem 1]. Second, we establish an asymptotic formula for  $V(p, q)$ . Finally, by use of continued fraction theory, we give a formula for  $V(p, q)$  in terms of continued fraction of  $\frac{p}{q}$ .

### 1. Introduction and statement of results

**1.1. Introduction.** Let  $H = L^2([0, \infty); t^{-2}dt)$  be the Hilbert space with the inner product

$$(1.1) \quad \langle f, g \rangle = \int_0^\infty f(t) \overline{g(t)} t^{-2} dt, \quad f, g \in H.$$

For any real number  $x$ ,  $[x]$  is the integer part of  $x$ , and  $\{x\} = x - [x]$  is the fractional part of  $x$ . Let  $p$  be a positive integer. Denote by  $e_p$  the function in  $H$  given by  $e_p(t) = \{t/p\}$ . The properties of the subspace  $H_n = \text{Vect}(e_1, \dots, e_n)$  spanned by the functions  $e_1, \dots, e_n$  is studied in [6, 8]. Consider the characteristic function

$$\chi(t) := \begin{cases} 1, & \text{if } t \in [1, \infty), \\ 0, & \text{otherwise.} \end{cases}$$

---

2010 *Mathematics Subject Classification*: Primary: 11B99, 11F67, 11E45; Secondary: 11M26, 11B68.

*Key words and phrases*: Vasyunin-cotangent sum, Estermann zeta function, fractional part function, Riemann Hypothesis.

Supported by Université d'Evry Val d'Essonne and PHC-Tassili program 14MDU914.

Communicated by Gradimir Milovanović.

The distance  $d_n$  from  $\chi$  to the subspace  $H_n$  is given by  $d_n = \text{dist}(\chi, H_n) = \inf_{h \in H_n} \|\chi - h\|$ . In [7] it conjectured that

$$(1.2) \quad d_n^2 \sim \frac{2 + \gamma - \log 4\pi + o(1)}{\log n}, \quad n \rightarrow +\infty.$$

Balazard and Roton proved in [9] that

$$d_n^2 \geq \frac{2 + \gamma - \log 4\pi}{\log n}, \quad n \rightarrow +\infty.$$

It is well known that conjecture (1.2) implies the Riemann hypothesis [2-6, 8]. For computation aspects of  $d_n$ , we refer to Landreau et al [18].

For fixed  $n \geq 1$ , from [10], we quote the formula

$$d_n^2 = \frac{\text{Gram}(\chi, e_1, \dots, e_n)}{\text{Gram}(e_1, \dots, e_n)}.$$

To compute this quantity, we need to evaluate two types of inner products, namely,  $\langle \chi, e_p \rangle$ ,  $\langle e_p, e_q \rangle$ . The first one is given in [6, §1] by

$$\langle \chi, e_p \rangle = \frac{\log p + 1 - \gamma}{p}.$$

On the other hand, for two coprime numbers  $p, q$ , the second inner product is given by the Vasyunin formula [8, 28]

$$(1.3) \quad \langle e_p, e_q \rangle = \frac{\log 2\pi - \gamma}{2} \left( \frac{1}{p} + \frac{1}{q} \right) + \frac{p-q}{2pq} \log \frac{q}{p} - \frac{\pi}{2pq} [V(q, p) + V(p, q)]$$

where

$$(1.4) \quad V(q, p) = \sum_{k=1}^{p-1} \left\{ \frac{kq}{p} \right\} \cot \left( \frac{\pi k}{p} \right).$$

The Vasyunin-cotangent sum  $V(p, q)$  is still curious. Recently, Bettin and Conrey [13, Theorem 1] proved a formula for

$$\frac{\bar{q}}{p} V(q, p) + V(\bar{p}, \bar{q})$$

where  $p\bar{p} \equiv 1(q)$  and  $q\bar{q} \equiv 1(p)$  with  $1 \leq \bar{p} \leq q$ ,  $1 \leq \bar{q} \leq p$ .

For  $q = 1$ , the cotangent sum (1.4) is first studied by Vasyunin [28]. He proved the asymptotic formula (for large  $p$ )

$$V(1, p) = -\frac{p \log p}{\pi} + \frac{p}{\pi} (\log 2\pi - \gamma) + O(\log p).$$

This formula is improved by Rassias [26] and Maier and Rassias [20] as follows

$$V(1, p) = -\frac{p \log p}{\pi} + \frac{p}{\pi} (\log 2\pi - \gamma) + O(1).$$

Let  $p, n \in \mathbb{N}$ ,  $p > 6N$ , with  $N = \lfloor \frac{n}{2} \rfloor + 1$ . Maier and Rassias [21, Theorem 1.7] proved that there exist absolute real constants  $A_1, A_2 \geq 1$  and absolute real

constants  $E_l, l \in \mathbb{N}$  with  $|E_l| \leq (A_1 l)^{2l}$ , such that for each  $n$ , we have

$$(1.5) \quad V(1, p) = -\frac{p \log p}{\pi} + \frac{p}{\pi}(\log 2\pi - \gamma) + \frac{1}{\pi} - \sum_{l=1}^n E_l p^{-l} - R_n^*(p)$$

where  $|R_n^*(p)| \leq (n A_2)^{4n} p^{-(n+1)}$ .

The sum  $V(p, q)$  can be interpreted as the value of the Estermann zeta function [13, 17] at  $s = 0$

$$E_0\left(s, \frac{p}{q}\right) = \sum_{k \geq 1} \frac{\tau(k)}{n^s} \exp\left(\frac{2\pi i k p}{q}\right).$$

Recently, this sum is studied in [12, 13] and it is proved that  $V(p, q)$  satisfies the reciprocity formula for all positive coprime numbers  $p$  and  $q$

$$(1.6) \quad \frac{\bar{q}}{p} V(q, p) + V(\bar{p}, \bar{q}) = -\frac{1}{\pi p} - g\left(\frac{\bar{q}}{p}\right)$$

where  $g$  is an analytic function in  $\mathbb{C} \setminus \mathbb{R}_-$ , which has the following asymptotic expansion of order  $N \geq 2, x \rightarrow 0$

$$g(x) = -\frac{\log 2\pi x - \gamma}{\pi} + \frac{2}{\pi} \sum_{k=2}^N \frac{\zeta(k) B_k}{k} x^k + O(x^{N+1}),$$

$B_k$  is the  $k^{\text{th}}$  Bernoulli number. Since  $\zeta(2k) = (-1)^{k-1} 2^{2k-1} \pi^{2k} B_{2k} / (2k)!$ , we have

$$g(x) = -\frac{\log 2\pi x - \gamma}{\pi} - \frac{1}{2} \sum_{k=1}^{\lfloor N/2 \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} x^k + O(x^{N+1}).$$

**1.2. Statement of main results.** Now consider the digamma function

$$\psi(z) = -\gamma - \frac{1}{z} + \sum_{k=1}^{\infty} \left(\frac{1}{k} - \frac{1}{k+z}\right)$$

and the symmetric function

$$(1.7) \quad G(p, q) = \sum_{r=1}^{pq-1} \left(\psi\left(\frac{r+1}{pq}\right) - \psi\left(\frac{r}{pq}\right)\right) \left\{\frac{r}{p}\right\} \left\{\frac{r}{q}\right\}.$$

We state a new reciprocity formula  $V(p, q)$ .

**THEOREM 1.1.** *For  $p, q$  positive coprime numbers, we have*

$$(1.8) \quad V(p, q) + V(q, p) = \frac{1}{\pi} \log p^{q-1} q^{p-1} - \frac{2}{\pi} - pg\left(\frac{1}{p}\right) - qg\left(\frac{1}{q}\right) - \frac{2}{\pi} G(p, q).$$

**COROLLARY 1.1.** *For  $p, q$  positive coprime numbers, we have*

$$(1.9) \quad \langle e_p, e_q \rangle = \frac{2 + (\log 2\pi - \gamma)(p+q)}{2pq} - \frac{1}{2pq} \log p^{p-1} q^{q-1} + \frac{\pi}{2} \left(\frac{1}{q} g\left(\frac{1}{p}\right) + \frac{1}{p} g\left(\frac{1}{q}\right)\right) + \frac{1}{pq} G(p, q).$$

Next, we state an asymptotic formula for the sum  $V(\bar{a}, pa + r)$ .

**THEOREM 1.2.** *Let  $a > r$  and  $p$  be integers such that  $(a, pa + r) = 1$ ,  $\bar{a}a \equiv 1 \pmod{pa + r}$  and  $\bar{r}r \equiv 1 \pmod{a}$ . Then for large  $p$ , we have*

$$(1.10) \quad V(\bar{a}, pa + r) = -\left(p + \frac{r}{a}\right)V(\bar{r}, a) - \frac{1}{\pi a} + \frac{1}{\pi}\left(p + \frac{r}{a}\right)\left(\log \frac{2\pi a}{pa + r} - \gamma\right) \\ + \frac{1}{2} \sum_{k=1}^{\lfloor N/2 \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{a}{pa + r}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

We get the following corollary.

**COROLLARY 1.2.** *Let  $a \geq 1$ . For large  $p$ , we have*

$$(1.11) \quad V(p+1, ap + a - 1) = \\ -\left(p+1 - \frac{1}{a}\right)V(1, a) - \frac{1}{\pi a} + \frac{1}{\pi}\left(p+1 - \frac{1}{a}\right)\left(\log \frac{2\pi a}{(p+1)a - 1} - \gamma\right) \\ + \frac{1}{2} \sum_{k=1}^{\lfloor \frac{N}{2} \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{a}{(p+1)a - 1}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

In the particular case  $a = 1$ , we have

$$(1.12) \quad V(1, p) = \frac{1}{\pi} \left(\log \frac{2\pi}{p} - \gamma\right)p - \frac{1}{\pi} - \frac{\pi}{144p} \\ + \frac{1}{2} \sum_{k=2}^{\lfloor \frac{N}{2} \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{1}{p}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

Relation (1.12) improves asymptotic formula (1.5) proved by Maier and Rassias in [21, Theorem 1.7].

In what follows, we give a few examples.

**EXAMPLES 1.1.** For large  $p$  we have

$$V(p+1, 2p+1) = -\frac{1}{2\pi} + \frac{1}{\pi}\left(p + \frac{1}{2}\right)\left(\log \frac{4\pi}{2p+1} - \gamma\right) \\ + \frac{1}{2} \sum_{k=1}^{\lfloor N/2 \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{2}{2p+1}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

$$V(2p+1, 3p+1) = -\frac{p + \frac{1}{3}}{3\sqrt{3}} - \frac{1}{3\pi} + \frac{1}{\pi}\left(p + \frac{1}{3}\right)\left(\log \frac{3\pi}{2} - \gamma\right) \\ + \frac{1}{2} \sum_{k=1}^{\lfloor N/2 \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{3}{3p+1}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

$$V(p+1, 3p+2) = \frac{p + \frac{2}{3}}{3\sqrt{3}} - \frac{1}{3\pi} + \frac{1}{\pi}\left(p + \frac{2}{3}\right)\left(\log \frac{6\pi}{5} - \gamma\right) \\ + \frac{1}{2} \sum_{k=1}^{\lfloor N/2 \rfloor} (-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{3}{3p+2}\right)^{2k-1} + O\left(\frac{1}{p^N}\right).$$

We can consider  $V(q, p)$  as a function of a single rational argument, by defining  $V(q/p) = V(q, p)$ . That this function is well defined is clear from the conditions on  $q$  and  $p$ . By use of continued fractions [25, §7] and the reciprocity law [13, Theorem 1], we obtain

**THEOREM 1.3.** *Let  $1 < q < p$  be coprime positive integers and  $\bar{p}$  denote the inverse of  $p$  modulo  $q$ . Write  $\bar{p}/q = [a_0, a_1, \dots, a_n]$  for a simple finite continued fraction of  $\bar{p}/q$  and  $(p_k)_{0 \leq k \leq n+1}$  for the finite sequence given by:  $p_{k-1} = a_{k-1}p_k + p_{k+1}$  with  $p_0 = \bar{p}$ ,  $p_1 = q$  and  $p_{n+1} = 0$ . Then we have*

$$V(p, q) = \frac{1}{\pi(q[a_0, a_1, \dots, a_{n-2}] - \bar{p})} - \sum_{k=2}^n (-1)^k g([0, a_{k-1}, \dots, a_n]) \prod_{j=2}^k [a_{j-1}, \dots, a_n].$$

**COROLLARY 1.3.** *Let  $\frac{q}{p} = [b_0, \dots, b_n]$  and  $(q_k)_k$  the associated sequence:  $q_{k-1} = b_{k-1}q_k + q_{k+1}$ ,  $q_0 = q$ ,  $q_1 = p$ ,  $q_n = 1$ . Then we have*

$$(1.13) \quad V(q, p) = -\frac{1}{\pi} - pg\left(\frac{1}{p}\right) - \frac{1}{\pi} \log pq + \frac{1}{\pi} (\log q + (-1)^n \log q_{n-1}) - \frac{1}{\pi} \sum_{k=1}^{n-2} (-1)^k (q_k \log q_{k+1} + q_{k+1} \log q_k - 2G(q_k, q_{k+1})).$$

Moreover, we have

$$(1.14) \quad V(q, p) = -\frac{1}{\pi} (1 + \log f_1) - q \left( f_1 g\left(\frac{1}{qf_1}\right) - \frac{1}{\pi} \sum_{k=1}^{n-2} (-1)^k (f_k \log f_{k+1}) + f_{k+1} \log f_k \right) + \frac{(-1)^n}{\pi} \log f_{n-1} + \frac{(-1)^n - 1}{\pi} \log q + \frac{((-1)^n f_{n-1} - f_1) q \log q}{\pi} + \frac{2}{\pi} \sum_{k=1}^{n-2} (-1)^k G(qf_k, qf_{k+1}), \quad \text{where } f_k = \prod_{j=1}^k [0, b_{j-1}, \dots, b_n].$$

**1.3. General case:  $p, q$  arbitrary.** Let  $\omega = \gcd(p, q) \geq 1$ . Vasyunin formula in [28] is given by

$$\langle e_p, e_q \rangle = \frac{\log 2\pi - \gamma}{2} \left( \frac{1}{p} + \frac{1}{q} \right) + \frac{p-q}{2pq} \log \frac{q}{p} - \frac{\pi\omega}{2pq} \left( V\left(\frac{p}{\omega}, \frac{q}{\omega}\right) + V\left(\frac{q}{\omega}, \frac{p}{\omega}\right) \right).$$

As a consequence of Theorem 1.1, we find a new expression for  $\langle e_p, e_q \rangle$ :

$$\begin{aligned} \langle e_p, e_q \rangle &= \frac{2\omega + (\log 2\pi - \gamma)(p+q)}{2pq} \\ &\quad - \frac{1}{2pq} ((p-\omega) \log p + (q-\omega) \log q - (p+q-2\omega) \log \omega) \\ &\quad + \frac{\pi}{2} \left( \frac{1}{p} g\left(\frac{\omega}{q}\right) + \frac{1}{q} g\left(\frac{\omega}{p}\right) \right) + \frac{\omega}{pq} G\left(\frac{p}{\omega}, \frac{q}{\omega}\right). \end{aligned}$$

## 2. Proof of Theorem 1.1 and Corollary 1.1

We start this section with some useful preliminary results.

**2.1. Computation of  $\langle v_p, v_q \rangle$ .** Let  $p$  be positive integer and  $v_p$  be the function given by  $v_p(t) = \{\lfloor t \rfloor / p\}$ ;  $v_p$  is defined on  $\mathbb{R}_+$  and it can be regarded as a restriction of  $e_p$  to  $\mathbb{N}$ . Since

$$v_p(x) = \begin{cases} 0, & \text{if } x \in [0, 1], \\ \{k/p\}, & x \in [k, k+1] \end{cases}$$

we have the relation

$$(2.1) \quad v_p = e_p - \frac{1}{p}e_1.$$

Then, from definition (1.1), we get

$$\langle v_p, v_q \rangle = \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\} \left\{ \frac{k}{q} \right\}.$$

LEMMA 2.1. *Let  $p, q$  be coprime numbers. Then we have*

$$\frac{2pq}{\pi} \langle v_p, v_q \rangle = V(1, p) + V(1, q) - (V(p, q) + V(q, p)) + \frac{1}{\pi} \log p^{q-1} q^{p-1}.$$

PROOF. Using expression (2.1), we get

$$\langle v_p, v_q \rangle = \langle e_p - \frac{1}{p}e_1, e_q - \frac{1}{q}e_1 \rangle = \langle e_p, e_q \rangle + \frac{1}{pq} \langle e_1, e_1 \rangle - \frac{1}{p} \langle e_1, e_q \rangle - \frac{1}{q} \langle e_1, e_p \rangle$$

On the other hand, from Vasyunin formula (1.3), we obtain

$$\begin{aligned} \langle e_1, e_q \rangle &= \frac{\log 2\pi - \gamma}{2} \left(1 + \frac{1}{q}\right) + \frac{1-q}{2q} \log q - \frac{\pi}{2q} V(1, q), \\ \langle e_1, e_p \rangle &= \frac{\log 2\pi - \gamma}{2} \left(1 + \frac{1}{p}\right) + \frac{1-p}{2p} \log p - \frac{\pi}{2p} V(1, p), \\ \langle e_p, e_q \rangle &= \frac{\log 2\pi - \gamma}{2} \left(\frac{1}{p} + \frac{1}{q}\right) + \frac{p-q}{2pq} \log \frac{q}{p} - \frac{\pi}{2pq} [V(q, p) + V(p, q)]. \end{aligned}$$

Therefore we deduce

$$\frac{2pq}{\pi} \langle v_p, v_q \rangle = V(1, p) + V(1, q) - V(p, q) - V(q, p) + \frac{1}{\pi} (q \log p + p \log q - \log pq). \quad \square$$

The series

$$(2.2) \quad \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\} \left\{ \frac{k}{q} \right\}$$

is convergent. Let us rewrite it in another form. For integer  $k$  we set  $k \equiv r(pq)$ ,  $1 \leq r \leq pq-1$  and then  $\left\{ \frac{k}{p} \right\} \left\{ \frac{k}{q} \right\} = \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\}$ . Series (2.2) is equal to

$$(2.3) \quad \sum_{r=1}^{pq-1} \sum_{i \geq 0} \frac{1}{(ipq+r)(ipq+r+1)} \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\}.$$

Finally, to estimate  $\langle v_p, v_q \rangle$  we need to compute the sums

$$\sum_{i \geq 0} \frac{1}{(ipq + r)(ipq + r + 1)}.$$

LEMMA 2.2. *For  $a, b$  two distinct positive numbers, we have*

$$\sum_{k=0}^{\infty} \frac{1}{(k+a)(k+b)} = \frac{\psi(a) - \psi(b)}{a-b}.$$

PROOF. We write

$$\psi(a) - \psi(b) = \frac{1}{b} - \frac{1}{a} + \sum_{k \geq 1} \left( \frac{1}{k+b} - \frac{1}{k+a} \right) = (a-b) \sum_{k=0}^{\infty} \frac{1}{(k+a)(k+b)}. \quad \square$$

COROLLARY 2.1. *Let  $\alpha > 0$ . We have*

$$(2.4) \quad \sum_{k=0}^{\infty} \frac{1}{(\alpha k + a)(\alpha k + b)} = \frac{\psi(a/\alpha) - \psi(b/\alpha)}{\alpha(a-b)}.$$

From [19], we have the integral representation

$$\psi(a) - \psi(b) = \int_0^{\infty} \frac{e^{-bt} - e^{-at}}{1 - e^{-t}} dt.$$

Thus, we can get

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{a-b}{(k+a)(k+b)} &= \sum_{k=0}^{\infty} \left( \frac{1}{k+b} - \frac{1}{k+a} \right) = \sum_{k=0}^{\infty} \int_0^1 (x^{k+b-1} - x^{k+a-1}) dx \\ &= \int_0^1 \frac{x^{b-1} - x^{a-1}}{1-x} dx = \int_0^{\infty} \frac{e^{-bt} - e^{-at}}{1 - e^{-t}} dt. \end{aligned}$$

Hence, the special case  $\alpha = pq$  of relations (1.7), (2.3), and (2.4) gives the result.

COROLLARY 2.2. *For  $p, q$  arbitrary, we have*

$$(2.5) \quad \sum_{k \geq 1} \frac{pq}{k(k+1)} \left\{ \frac{k}{p} \right\} \left\{ \frac{k}{q} \right\} = G(p, q).$$

**2.2. Proof of Theorem 1.1 and Corollary 1.1.** From Lemma 2.1 and Corollary 2.2, we deduce that

$$2G(p, q) = V(1, p) + V(1, q) - (V(p, q) + V(q, p)) + \frac{1}{\pi} \log p^{q-1} q^{p-1}.$$

Therefore, we get (1.8) from the expression

$$(2.6) \quad V(1, p) = -\frac{1}{\pi} - pg \left( \frac{1}{p} \right).$$

We can obtain formula (1.9) by use of reciprocity formula (1.8) and Vasyunin formula (1.3).

**3. Proof of Theorem 1.2 and Corollary 1.2**

From reciprocity law (1.6), we have

$$\frac{a}{pa+r}V(\bar{a}, pa+r) + V(\bar{r}, a) = -\frac{1}{\pi(pa+r)} - g\left(\frac{a}{pa+r}\right)$$

and then we obtain

$$V(\bar{a}, pa+r) = -\frac{pa+r}{a}V(\bar{r}, a) - \frac{1}{\pi a} - \frac{pa+r}{a}g\left(\frac{a}{pa+r}\right).$$

Moreover, for large  $p$  we have

$$g\left(\frac{a}{pa+r}\right) = -\frac{1}{\pi}\left(\log\frac{2\pi a}{pa+r} - \gamma\right) - \frac{1}{2}\sum_{k=1}^{\lfloor N/2 \rfloor}(-1)^k \frac{4^k \pi^{2k-1} B_{2k}^2}{k(2k)!} \left(\frac{a}{pa+r}\right)^k + O\left(\frac{1}{p^{N+1}}\right)$$

Taking  $r = a - 1$ , one remarks that for  $a \geq 1$   $(p+1)a \equiv 1 \pmod{pa+a-1}$  and  $(a-1)^2 \equiv 1(a)$ . Then  $\bar{a} = p+1$  and  $\bar{r} = r = a-1$ . Since  $V(a-1, a) = V(1, a)$  and thanks to relation (1.10) of Theorem 1.2, we obtain relation (1.11). Finally from relation (1.11) we can get (1.12).

**4. Proof of Theorem 1.3 and Corollary 1.3**

**4.1. Continued fractions – an overview.** In this subsection we quote some facts about continued fractions, that will be useful later. For more details we refer to [25, §7]. For real number  $x$  let

$$x = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}} = [a_0, a_1, a_3, \dots]$$

be its continued fraction expansion with partial quotients  $a_0 \in \mathbb{Z}$ ,  $a_k \in \mathbb{N} \setminus \{0\}$ ,  $k \geq 1$ . In fact, we can determine  $a_0, a_1, \dots, a_n$  via the *algorithm*:

- $x = [x] + \{x\}$ ,  $a_0 = [x]$ ,  $\xi_0 = \{x\}$ , if  $\xi_0 = 0$ , then  $x$  is represented by  $x = [a_0]$ .
- if  $\xi_0 \neq 0$  then  $\lfloor \frac{1}{\xi_0} \rfloor > 1$ ,  $r_1 = \frac{1}{\xi_0}$  we obtain  $x = [a_0, r_1]$ ,  $a_1 = [r_1]$  and  $\xi_1 = r_1 - a_1$ ; if  $\xi_1 = 0$  then  $x = a_0 + \frac{1}{r_1} = a_0 + \frac{1}{a_1} = [a_0, a_1]$ .
- Otherwise we take  $r_2 = \frac{1}{\xi_1}$  and iterate the process.

We then get the sequence  $a_0, a_1, a_2, \dots$ . This sequence is finite if and only if  $x$  is a rational number. In the rational case, this algorithm is the *Euclidian algorithm*. Let us express  $a/b$  as a continued fraction of the form  $a/b = [a_0, a_1, \dots, a_n]$ , with  $a_{n+1} = 0$ . We can determine  $a_0, a_1, \dots, a_n$  by the Euclidean algorithm

$$(4.1) \quad p_{k-1} = a_{k-1}p_k + p_{k+1}, \quad p_0 = a, \quad p_1 = b, \quad p_{n+1} = 0.$$

Then  $p_n = \gcd(p, q) = 1$ . We quote from [25, §7] the following elementary properties of continued fractions which are needed in this paper. We omit their proofs.



LEMMA 4.1. *Let  $[a_0, a_1, \dots, a_n]$  be continued fraction. Then*

$$[a_0, a_1, \dots, a_n] = \left[ a_0, a_1, \dots, a_{n-1} + \frac{1}{a_n} \right],$$

$$[a_0, a_1, \dots, a_n] = a_0 + \frac{1}{[a_1, \dots, a_n]}.$$

We get this lemma from the continued fraction definition. Let  $s_k, t_k$  be two sequences of integers

$$s_k = \begin{cases} 0, & \text{if } k = -2 \\ 1, & \text{if } k = -1 \\ a_k s_{k-1} + s_{k-2}, & \text{if } k \geq 0 \end{cases} \quad t_k = \begin{cases} 1, & \text{if } k = -2 \\ 0, & \text{if } k = -1 \\ a_k t_{k-1} + t_{k-2}, & \text{if } k \geq 0 \end{cases}$$

The sequences  $p_k, s_k$  and  $t_k$  have the following properties.

LEMMA 4.2.

$$(4.2) \quad [a_0, a_1, \dots, a_k] = \frac{s_k}{t_k}, \quad k \geq 0,$$

$$[a_k, a_{k-1}, \dots, a_1] = \frac{t_k}{t_{k-1}}, \quad k \geq 1,$$

$$(4.3) \quad [0, a_{k-1}, \dots, a_n] = \frac{p_k}{p_{k-1}}, \quad k \geq 0,$$

$$[a_k, a_{k-1}, \dots, a_0] = \frac{s_k}{s_{k-1}}, \quad k \geq 0.$$

This lemma represents the classical properties of the continued fractions. The quotient  $\frac{s_k}{t_k}$  is called the  $k^{\text{th}}$ -convergent of the continued fraction of  $\frac{a}{b}$ .

COROLLARY 4.1. *Let  $\frac{a}{b} = [a_0, a_1, \dots, a_n]$ ; then*

$$(4.4) \quad p_k = b \prod_{j=2}^k [0, a_{j-1}, \dots, a_n], \quad k \geq 2,$$

$$b = p_k t_{k-1} + p_{k+1} t_{k-2}, \quad k \geq 1.$$

LEMMA 4.3. *The sequences  $s_k, t_k$  and  $p_k$  satisfy*

$$(4.5) \quad t_k s_{k-1} - t_{k-1} s_k = (-1)^k,$$

$$p_k = (-1)^k (a t_{k-2} - b s_{k-2}).$$

PROPOSITION 4.1. *Under the hypothesis of Corollary 4.1, we have*

$$(4.6) \quad \sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} = \frac{1}{b^2 [a_0, a_1, \dots, a_{n-2}] - ab}.$$

PROOF. We have from Lemma 4.3

$$\sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} = \frac{1}{b} \sum_{k=1}^{n-1} \left( \frac{(-1)^k t_{k-1}}{p_{k+1}} + \frac{(-1)^k t_{k-2}}{p_k} \right)$$

$$\begin{aligned}
&= \frac{1}{b} \sum_{k=1}^{n-1} \frac{(-1)^k t_{k-1}}{p_{k+1}} + \frac{1}{b} \sum_{k=1}^{n-1} \frac{(-1)^k t_{k-2}}{p_k} \\
&= \frac{1}{b} \sum_{k=2}^n \frac{(-1)^{k-1} t_{k-2}}{p_k} + \frac{1}{b} \sum_{k=1}^{n-1} \frac{(-1)^k t_{k-2}}{p_k} \\
&= \frac{(-1)^{n-1} t_{n-2}}{p_n b} - \frac{t_{-1}}{b p_1} = \frac{(-1)^{n-1} t_{n-2}}{p_n b}.
\end{aligned}$$

From (4.5) we deduce  $p_n = (-1)^n (at_{n-2} - bs_{n-2})$  and then

$$(4.7) \quad \frac{(-1)^{n-1} p_n}{t_{n-2}} = \left( b \frac{s_{n-2}}{t_{n-2}} - a \right).$$

Relation (4.2) implies that

$$(4.8) \quad \frac{s_{n-2}}{t_{n-2}} = [a_0, a_1, \dots, a_{n-2}]$$

From relations (4.7) and (4.8) we have

$$\frac{(-1)^{n-1} t_{n-2}}{p_n b} = \frac{1}{b^2 [a_0, a_1, \dots, a_{n-2}] - ab}.$$

Therefore we obtain

$$\sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} = \frac{1}{b^2 [a_0, a_1, \dots, a_{n-2}] - ab}. \quad \square$$

**4.2. Application: Computation of  $V(p, q)$ .** Using reciprocity formula (1.6) and continued fractions properties, we can get an explicit formula for  $V(p, q)$ .

LEMMA 4.4. *Let  $p$  and  $q$  be coprime positive numbers. We have*

$$(4.9) \quad V(p, q) = \frac{q}{\pi} \sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} - q \sum_{k=2}^n \frac{(-1)^k}{p_k} g\left(\frac{p_k}{p_{k-1}}\right).$$

PROOF. From the reciprocity law (1.6), with  $\bar{p} = a$ ,  $q = b$  and the sequence  $(p_k)_k$  defined in relation (4.1), we have

$$(4.10) \quad \frac{1}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{1}{p_k} V(\bar{p}_{k-1}, p_k) = \frac{1}{p_k} \left( \frac{1}{\pi p_{k-1}} + g\left(\frac{p_k}{p_{k-1}}\right) \right).$$

Observe that

$$p_{k-1} = a_{k-1} p_k + p_{k+1}, \quad \bar{p}_{k-1} = \bar{p}_{k+1}, \quad V(\bar{p}_{k-1}, p_k) = V(\bar{p}_{k+1}, p_k),$$

then relation (4.10) becomes

$$\frac{1}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{1}{p_k} V(\bar{p}_{k+1}, p_k) = -\frac{1}{p_k} \left( \frac{1}{\pi p_{k-1}} + g\left(\frac{p_k}{p_{k-1}}\right) \right).$$

We have the computation

$$\sum_{k=1}^n \left[ \frac{(-1)^k}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}/p_k) \right]$$

$$\begin{aligned} &= \sum_{k=1}^n \frac{(-1)^k}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \sum_{k=1}^n \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}, p_k) \\ &= \sum_{k=0}^{n-1} \frac{(-1)^{k+1}}{p_k} V(\bar{p}_{k+1}, p_k) + \sum_{k=1}^n \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}, p_k) \\ &= -\frac{1}{p_0} V(\bar{p}_1, p_0) + \frac{(-1)^n}{p_n} V(\bar{p}_{n+1}, p_n). \end{aligned}$$

Since  $p_{n+1} = 0$ , the relation  $V(\bar{p}_{n+1}, p_n) = 0$  implies

$$\begin{aligned} &\sum_{k=1}^n \left[ \frac{(-1)^k}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}, p_k) \right] = -\frac{1}{\bar{p}} V(\bar{q}, \bar{p}) \\ &= -\frac{1}{\bar{p}} V(\bar{q}, \bar{p}) - \frac{1}{q} V(p, q) + \sum_{k=2}^n \left[ \frac{(-1)^k}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}, p_k) \right]. \end{aligned}$$

Moreover,

$$\frac{1}{q} V(p, q) = \sum_{k=2}^n \left[ \frac{(-1)^k}{p_{k-1}} V(\bar{p}_k, p_{k-1}) + \frac{(-1)^k}{p_k} V(\bar{p}_{k+1}, p_k) \right]$$

then we deduce

$$\frac{1}{q} V(p, q) = -\sum_{k=2}^n \frac{(-1)^k}{p_k} \left( \frac{1}{\pi p_{k-1}} + g\left(\frac{p_k}{p_{k-1}}\right) \right) = \frac{1}{\pi} \sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} - \sum_{k=2}^n \frac{(-1)^k}{p_k} g\left(\frac{p_k}{p_{k-1}}\right). \quad \square$$

**4.3. Proof of Theorem 1.3 and Corollary 1.3.** First, we prove Theorem 1.3. Taking  $a = \bar{p}$  and  $b = q$ , by virtue of relation (4.6) we obtain

$$(4.11) \quad \frac{q}{\pi} \sum_{k=1}^{n-1} \frac{(-1)^k}{p_k p_{k+1}} = \frac{1}{q[a_0, a_1, \dots, a_{n-2}] - \bar{p}}.$$

From (4.4) we get

$$\frac{1}{p_k} = \frac{1}{q} \prod_{j=2}^k [a_{j-1}, \dots, a_n], \quad k \geq 2,$$

and from (4.3) we obtain

$$(4.12) \quad \sum_{k=2}^n \frac{(-1)^k}{p_k} g\left(\frac{p_k}{p_{k-1}}\right) = \sum_{k=2}^n \frac{1}{q} g([a_{k-1}, \dots, a_n]) \prod_{j=2}^k [a_{j-1}, \dots, a_n].$$

Substituting quantities (4.11) and (4.12) into (4.9) we get Theorem 1.3.

To prove Corollary 1.3, note that for  $p \equiv r(q)$  we have

$$(4.13) \quad V(p, q) = \begin{cases} 0, & \text{if } r = 0 \\ V(r, q), & \text{otherwise.} \end{cases}$$

Applying several times reciprocity formula (1.8) to the sequence  $q_k$ , we obtain

$$\sum_{k=1}^{n-1} (-1)^k \theta(q_{k-1}, q_k) = \sum_{k=1}^{n-1} (-1)^k [V(q_{k-1}, q_k) + V(q_k, q_{k-1})]$$

where

$$\begin{aligned} \theta(q_{k-1}, q_k) &= \frac{1}{\pi} (q_{k-1} \log q_k + q_k \log q_{k-1} - \log q_{k-1} q_k) \\ &\quad - \frac{2}{\pi} - q_{k-1} g\left(\frac{1}{q_{k-1}}\right) - q_k g\left(\frac{1}{q_k}\right) - \frac{2}{\pi} G(q_{k-1}, q_k). \end{aligned}$$

From (4.13) we have  $V(q_{k+1}, q_k) = V(q_{k-1}, q_k)$  and  $V(q_{n-1}, q_n) = 0$ , and then

$$\begin{aligned} \sum_{k=1}^{n-1} (-1)^k \theta(q_{k-1}, q_k) &= \sum_{k=1}^{n-1} (-1)^k V(q_{k-1}, q_k) + \sum_{k=1}^{n-1} (-1)^k V(q_k, q_{k-1}) \\ &= \sum_{k=1}^{n-1} (-1)^k V(q_{k+1}, q_k) + \sum_{k=0}^{n-2} (-1)^{k+1} V(q_{k+1}, q_k) \\ &= -V(p, q) - (-1)^n V(q_n, q_{n-1}). \end{aligned}$$

In addition we have

$$\begin{aligned} \sum_{k=1}^{n-1} (-1)^k \theta(q_{k-1}, q_k) &= \frac{1}{\pi} \sum_{k=1}^{n-1} (-1)^k [(q_{k-1} \log q_k + q_k \log q_{k-1} - \log q_{k-1} q_k)] \\ &= -\frac{2}{\pi} \sum_{k=1}^{n-1} (-1)^k [1 + G(q_{k-1}, q_k)] \\ &\quad - \sum_{k=1}^{n-1} (-1)^k \left[ q_{k-1} g\left(\frac{1}{q_{k-1}}\right) + q_k g\left(\frac{1}{q_k}\right) \right] \end{aligned}$$

and we push this computation we arrive to

$$\begin{aligned} \sum_{k=1}^{n-1} (-1)^k \left[ q_{k-1} g\left(\frac{1}{q_{k-1}}\right) + q_k g\left(\frac{1}{q_k}\right) \right] &= \sum_{k=1}^{n-1} (-1)^k q_{k-1} g\left(\frac{1}{q_{k-1}}\right) + \sum_{k=1}^{n-1} (-1)^k q_k g\left(\frac{1}{q_k}\right) \\ &= \sum_{k=0}^{n-2} (-1)^{k+1} q_k g\left(\frac{1}{q_k}\right) + \sum_{k=1}^{n-1} (-1)^k q_k g\left(\frac{1}{q_k}\right) \\ &= -q g\left(\frac{1}{q}\right) + (-1)^{n-1} q_{n-1} g\left(\frac{1}{q_{n-1}}\right). \end{aligned}$$

We have also

$$\begin{aligned} \sum_{k=1}^{n-1} (-1)^k \log q_{k-1} q_k &= \sum_{k=1}^{n-1} (-1)^k (\log q_{k-1} + \log q_k) \\ &= \sum_{k=1}^{n-1} (-1)^k \log q_{k-1} + \sum_{k=1}^{n-1} (-1)^k \log q_k \end{aligned}$$

$$\begin{aligned}
 &= \sum_{k=0}^{n-2} (-1)^{k+1} \log q_k + \sum_{k=1}^{n-1} (-1)^k \log q_k \\
 &= -\log q - (-1)^n \log q_{n-1}.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 -V(p, q) - (-1)^n V(q_n, q_{n-1}) &= \frac{1}{\pi} \sum_{k=1}^{n-1} (-1)^k (q_{k-1} \log q_k + q_k \log q_{k-1}) + qg\left(\frac{1}{q}\right) \\
 &\quad + (-1)^n q_{n-1} g\left(\frac{1}{q_{n-1}}\right) - \frac{2}{\pi} \sum_{k=1}^{n-1} (-1)^k [1 + G(q_{k-1}, q_k)] \\
 &\quad + \frac{1}{\pi} (\log q + (-1)^n \log q_{n-1})
 \end{aligned}$$

and

$$\begin{aligned}
 -V(p, q) &= (-1)^n V(q_n, q_{n-1}) + (-1)^n q_{n-1} g\left(\frac{1}{q_{n-1}}\right) - \frac{1}{\pi} (q \log p + p \log q) \\
 &\quad + qg\left(\frac{1}{q}\right) + \frac{2}{\pi} (1 + G(q, p)) + \frac{1}{\pi} (\log q + (-1)^n \log q_{n-1}) \\
 &\quad - \frac{1}{\pi} \sum_{k=1}^{n-2} (-1)^k [q_k \log q_{k+1} + q_{k+1} \log q_k - 2G(q_k, q_{k+1}) - 2] \\
 &= \frac{(-1)^{n+1}}{\pi} - V(p, q) - V(q, p) - \frac{1}{\pi} \log pq - pg\left(\frac{1}{p}\right) \\
 &\quad + \frac{1}{\pi} (\log q + (-1)^n \log q_{n-1}) \\
 &\quad - \frac{1}{\pi} \sum_{k=1}^{n-2} (-1)^k [q_k \log q_{k+1} + q_{k+1} \log q_k - 2G(q_k, q_{k+1}) - 2].
 \end{aligned}$$

Then we complete the proof of relation (1.13)

$$\begin{aligned}
 V(q, p) &= -\frac{1}{\pi} - pg\left(\frac{1}{p}\right) - \frac{1}{\pi} \log pq + \frac{1}{\pi} (\log q + (-1)^n \log q_{n-1}) \\
 &\quad - \frac{1}{\pi} \sum_{k=1}^{n-2} (-1)^k [q_k \log q_{k+1} + q_{k+1} \log q_k - 2G(q_k, q_{k+1})].
 \end{aligned}$$

We observe that for any integer  $1 \leq k \leq n$  we have  $q_k = q \prod_{j=1}^k [0, b_{j-1}, \dots, b_n]$ , so that we can get (1.14) from (1.13).

### 5. Further identities on $V(p, q)$ and computation

In this section we relate the sums  $V(p, q)$  to some interesting and well-known convergent series. The functions  $\psi$  and cotangent are related by the reflection formula [1, §6.3.7]

$$(5.1) \quad \psi(1-z) - \psi(z) = \pi \cot(\pi z).$$

Moreover,  $\psi$  can be written in terms of harmonic function  $H_n(z) = \sum_{k=0}^n \frac{1}{(k+z)}$  and the  $n$ -th harmonic number  $H_n = \sum_{k=1}^n \frac{1}{k}$  as follows.

LEMMA 5.1. *Let  $z \in \mathbb{C}$  such that  $\operatorname{Re}(z) > 0$ . Then we have*

$$\psi(z) = \lim_{n \rightarrow \infty} (\log n - H_n(z))$$

at  $z = n$  positive integer we have

$$(5.2) \quad \psi(n+1) = -\gamma + H_n.$$

PROOF. It is easy to see that

$$\psi(z) = -\gamma - \frac{1}{z} + \sum_{k=1}^{\infty} \left( \frac{1}{k} - \frac{1}{k+z} \right) = \lim_{n \rightarrow \infty} \left( \log n - \sum_{k=0}^n \frac{1}{k+z} \right).$$

Then

$$\begin{aligned} \psi(n+1) &= \lim_{m \rightarrow \infty} \left( \log m - \sum_{k=0}^m \frac{1}{k+n+1} \right) = \lim_{m \rightarrow \infty} \left( \log m - \sum_{k=n}^{m+n} \frac{1}{k+1} \right) \\ &= \lim_{m \rightarrow \infty} \left( \log m - \left( \sum_{k=0}^{m+n} \frac{1}{k+1} - \sum_{k=0}^{n-1} \frac{1}{k+1} \right) \right) \\ &= \lim_{m \rightarrow \infty} \left( \log(m+n) - \sum_{k=0}^{m+n} \frac{1}{k+1} \right) + H_n = -\gamma + H_n. \quad \square \end{aligned}$$

PROPOSITION 5.1. *For  $x = \bar{p}/q$  with  $(p, q) = 1$ , we have*

$$(5.3) \quad V(p, q) = \frac{1}{\pi q} \sum_{k \geq 0} \sum_{r=1}^{q-1} \frac{r(1-2rx)}{(k+1-rx)(k+rx)}.$$

PROOF. We have

$$V(p, q) = \sum_{r=1}^{q-1} \left\{ \frac{rp}{q} \right\} \cot \left( \frac{\pi r}{q} \right) = \sum_{r=1}^{q-1} \frac{r}{q} \cot \left( \frac{\pi r \bar{p}}{q} \right).$$

By reflection formula (5.1) we have

$$\cot \left( \frac{\pi r \bar{p}}{q} \right) = \frac{1}{\pi} \left( \psi \left( \frac{q-r\bar{p}}{q} \right) - \psi \left( \frac{r\bar{p}}{q} \right) \right).$$

From relations (1.7) and (2.5) we have

$$\psi \left( \frac{q-r\bar{p}}{q} \right) - \psi \left( \frac{r\bar{p}}{q} \right) = q \sum_{k \geq 0} \frac{q-2r\bar{p}}{(q(k+1)-r\bar{p})(qk+r\bar{p})}.$$

Then

$$V(p, q) = \frac{1}{\pi} \sum_{r=1}^{q-1} \sum_{k \geq 0} \frac{r(q-2r\bar{p})}{(q(k+1)-r\bar{p})(qk+r\bar{p})},$$

and we obtain

$$V(p, q) = \frac{1}{\pi q} \sum_{r=1}^{q-1} \sum_{k \geq 0} \frac{r(1 - 2rx)}{(k + 1 - rx)(k + rx)}. \quad \square$$

As an immediate consequence we derive

COROLLARY 5.1. *Let  $q$  be a positive integer; we have*

$$(5.4) \quad V(1, q) = -\frac{q(\psi(q) + \gamma - 2) + 2}{\pi} + \frac{1}{\pi} \sum_{k \geq 1} \sum_{r=1}^{q-1} \frac{r(q - 2r)}{(q(k + 1) - r)(qk + r)},$$

$$V(1, q) = -\frac{q(\psi(q) + \gamma - 2) + 2}{\pi} - \frac{1}{\pi} \sum_{k \geq 1} \sum_{r=1}^{\lfloor q/2 \rfloor} \frac{(q - 2r)^2}{(q(k + 1) - r)(qk + r)}.$$

PROOF. From (5.3), special case  $p = 1$ , we can get

$$V(1, q) = \frac{1}{\pi q} \sum_{k \geq 0} \sum_{r=1}^{q-1} \frac{r(1 - 2r/q)}{(k + 1 - r/q)(k + r/q)}$$

then we have

$$V(1, q) = \frac{1}{\pi} \sum_{k \geq 0} \sum_{r=1}^{q-1} \frac{r(q - 2r)}{(q(k + 1) - r)(qk + r)}$$

and

$$V(1, q) = \frac{1}{\pi} \sum_{r=1}^{q-1} \frac{q - 2r}{q - r} - \frac{1}{\pi} \sum_{k \geq 1} \sum_{r=1}^{q-1} \frac{r(q - 2r)}{(q(k + 1) - r)(qk + r)}$$

$$= \frac{1}{\pi} (q(2 - H_{q-1}) - 2) + \frac{1}{\pi} \sum_{k \geq 1} \sum_{r=1}^{q-1} \frac{r(q - 2r)}{(q(k + 1) - r)(qk + r)}.$$

From (5.2) we obtain (5.4). To end the proof we use that for  $t + r = q$ , we have

$$\frac{1}{(q(k + 1) - r)(qk + r)} = \frac{1}{(q(k + 1) - t)(qk + t)},$$

$$r(q - 2r) + t(q - 2t) = (q - 2r)^2. \quad \square$$

PROPOSITION 5.2. *The sum  $V(1, q)$  has the following integral representation.*

$$(5.5) \quad V(1, q) = -\frac{1}{\pi} \int_0^1 \frac{(q - 2)x^q - qx^{q-1} + qx - q + 2}{(x - 1)^2(x^q - 1)} dx.$$

REMARK 5.1. Note that from (5.5) and (2.6) we deduce that

$$g\left(\frac{1}{q}\right) = \frac{1}{\pi q} \int_0^1 \left( \frac{(q - 2)x^q - qx^{q-1} + qx - q + 2}{(x - 1)^2(x^q - 1)} - 1 \right) dx.$$

PROOF. From the relation (5.4) we have

$$V(1, q) = \frac{1}{\pi} \sum_{r=1}^{q-1} r \sum_{k \geq 0} \frac{q-2r}{(q(k+1)-r)(qk+r)},$$

$$\frac{q-2r}{(q(k+1)-r)(qk+r)} = \frac{1}{qk+r} - \frac{1}{qk+q-r} = \int_0^1 (x^{qk+r-1} - x^{qk+q-r-1}) dx.$$

Then

$$\sum_{k \geq 0} \frac{q-2r}{(q(k+1)-r)(qk+r)} = \int_0^1 \frac{x^{r-1} - x^{q-r-1}}{1-x^q} dx$$

which gives

$$V(1, q) = -\frac{1}{\pi} \int_0^1 \frac{\sum_{r=1}^{q-1} r(x^{r-1} - x^{q-r-1})}{x^q - 1} dx.$$

On the other hand

$$\sum_{r=1}^{q-1} r x^{r-1} = \left( \frac{x^q - 1}{x - 1} \right)' = \frac{(q-1)x^q - qx^{q-1} + 1}{(x-1)^2}. \quad \square$$

Thanks to Proposition 5.2 we give a few values of  $V(1, p)$

EXAMPLE 5.1.  $V(1, 2) = 0$ ,

$$V(1, 3) = -\frac{1}{\pi} \int_0^1 \frac{dx}{x^2 + x + 1} = -\frac{1}{3\sqrt{3}}, \quad V(1, 4) = -\frac{2}{\pi} \int_0^1 \frac{dx}{x^2 + 1} = -\frac{1}{2},$$

$$V(1, 6) = -\frac{1}{\pi} \int_0^1 \left( \frac{3}{x^2 - x + 1} + \frac{1}{x^2 + x + 1} \right) dx = -\frac{7}{3\sqrt{3}}.$$

## 6. Another estimation of $V(p, q) + V(q, p)$

In this section we establish another asymptotic formula for the sum  $V(p, q) + V(q, p)$ . For this we prove the lemma

LEMMA 6.1. *Let  $p$  be a positive integer. We have*

$$(6.1) \quad \frac{\log p}{p} = \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\},$$

$$\sum_{r=1}^{pq-1} \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\} = \frac{(p-1)(q-1)}{4}.$$

PROOF. We have

$$\sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\} = \sum_{k \geq 1} \left\{ \frac{k}{p} \right\} \int_0^1 (x^{k-1} - x^k) dx$$

$$= \sum_{r=1}^{p-1} \sum_{i \geq 0} \left\{ \frac{r}{p} \right\} \int_0^1 (x^{ip+r-1} - x^{ip+r}) dx$$



$$= \frac{1}{p} \int_0^1 \frac{\sum_{r=1}^{p-1} r x^{r-1}}{1+x+\dots+x^{p-1}} dx = \frac{\log p}{p}.$$

Thus we obtain (6.1). Let us write

$$\sum_{r=1}^{pq-1} \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\} = \sum_{i=0}^{p-1} \sum_{t=1}^{q-1} \left\{ \frac{iq+t}{p} \right\} \left\{ \frac{t}{q} \right\} = \frac{1}{pq} \sum_{t=1}^{q-1} t \left( \sum_{i=0}^{p-1} \left\{ \frac{iq+t}{p} \right\} \right)$$

and we take  $r = p \left\{ \frac{iq+t}{p} \right\}$ , observe that  $r$  is integer and runs from 1 to  $p-1$ . Then, we have

$$\sum_{r=1}^{pq-1} \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\} = \frac{1}{pq} \sum_{t=0}^{q-1} \sum_{r=1}^{p-1} tr = \frac{(p-1)(q-1)}{4}. \quad \square$$

REMARK 6.1. Another proof of relation (6.1) is given in [10].

The geometry of  $\psi\left(\frac{r}{p}\right)$  is studied in [14, 24]. In the next corollary we write  $\log p$  as a linear combination of  $\psi\left(\frac{r}{p}\right)$  for  $1 \leq r \leq p-1$ .

COROLLARY 6.1. *Let  $p$  be a positive integer. Then we have*

$$\log p = \frac{1}{p} \sum_{r=1}^{p-1} r \left( \psi\left(\frac{r+1}{p}\right) - \psi\left(\frac{r}{p}\right) \right).$$

PROOF. We start with the equalities

$$\frac{\log p}{p} = \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\} = \sum_{r=1}^{p-1} \left( \sum_{i \geq 0} \frac{1}{(ip+r)(ip+r+1)} \right) \left\{ \frac{r}{p} \right\}$$

and from (2.4) we have

$$\sum_{i=0}^{\infty} \frac{1}{(pi+r)(pi+r+1)} = \frac{\psi(r+1)/p - \psi(r/p)}{p}$$

thus the result follows.  $\square$

PROPOSITION 6.1. *Let  $p, q$  be two coprime integers and put*

$$\Delta(p, q) = \frac{2}{\pi} \left( 1 + \max \left\{ \frac{(p-1)(q-1)}{4} \psi\left(\frac{1}{pq}\right), \min\{q - \gamma - \psi(q), p - \gamma - \psi(p)\} \right\} \right).$$

*Then we have*

$$(6.2) \quad \Delta(p, q) < \frac{1}{\pi} \log p^{q-1} q^{p-1} - pg\left(\frac{1}{p}\right) - qg\left(\frac{1}{q}\right) - V(p, q) - V(q, p) \\ \leq \frac{2 + 2\sqrt{pq \log p \log q}}{\pi}.$$

PROOF. From equality (6.1) we remark that

$$\langle v_p, v_q \rangle \leq \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{p} \right\} = \frac{\log p}{p} \quad \text{and} \quad \langle v_p, v_q \rangle \leq \sum_{k \geq 1} \frac{1}{k(k+1)} \left\{ \frac{k}{q} \right\} = \frac{\log q}{q}$$

we deduce that  $\langle v_p, v_q \rangle \leq \sqrt{\log p \log q / pq}$ . For  $q < p$  we have

$$\langle v_p, v_q \rangle > \frac{1}{pq} \sum_{k=1}^{q-1} \frac{k}{k+1}.$$

Similarly if  $p < q$  we have  $\langle v_p, v_q \rangle > \frac{1}{pq} \sum_{k=1}^{p-1} \frac{k}{k+1}$ . Therefore we have

$$\langle v_p, v_q \rangle > \frac{1}{pq} \left( q - \sum_{k=1}^{q-1} \frac{1}{k+1} \right) \quad \text{or} \quad \langle v_p, v_q \rangle > \frac{1}{pq} \left( p - \sum_{k=1}^{p-1} \frac{1}{k+1} \right).$$

From (5.2) we have

$$\langle v_p, v_q \rangle > \frac{1}{pq} (q - \gamma - \psi(q)) \quad \text{or} \quad \langle v_p, v_q \rangle > \frac{1}{pq} (p - \gamma - \psi(p))$$

and then  $\langle v_p, v_q \rangle > \frac{1}{pq} \min\{q - \gamma - \psi(q), p - \gamma - \psi(p)\}$ . Finally we obtain

$$\frac{1}{\pi pq} \min\{q - \gamma - \psi(q), p - \gamma - \psi(p)\} < \frac{1}{\pi} \langle v_p, v_q \rangle \leq \sqrt{\log p \log q / pq}.$$

In addition we have

$$pq \langle v_p, v_q \rangle = G(p, q) = \sum_{r=1}^{pq-1} \left( \psi\left(\frac{r+1}{pq}\right) - \psi\left(\frac{r}{pq}\right) \right) \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\}$$

and for  $x > 0$  and  $0 < y < 1$ , from Sulaiman [27, Theorem 2.2] we have the inequality  $\psi(x+y) - \psi(x) \geq \psi(y)$ . Taking  $x = \frac{r}{pq}$  and  $y = \frac{1}{pq}$ , from the above inequalities we obtain

$$G(p, q) \geq \psi\left(\frac{1}{pq}\right) \sum_{r=1}^{pq-1} \left\{ \frac{r}{p} \right\} \left\{ \frac{r}{q} \right\} \geq \frac{(p-1)(q-1)}{4} \psi\left(\frac{1}{pq}\right).$$

Furthermore

$$\frac{1}{pq} \max \left\{ \frac{(p-1)(q-1)}{4} \psi\left(\frac{1}{pq}\right), \min\{q - \gamma - \psi(q), p - \gamma - \psi(p)\} \right\} < \langle v_p, v_q \rangle \leq \sqrt{\log p \log q / pq}.$$

Then we have

$$\frac{2}{\pi} \max \left\{ \frac{(p-1)(q-1)}{4} \psi\left(\frac{1}{pq}\right), \min\{q - \gamma - \psi(q), p - \gamma - \psi(p)\} \right\} < \frac{2}{\pi} G(p, q) \leq \frac{2}{\pi} \sqrt{pq \log p \log q}.$$

From (1.8) we obtain

$$\frac{2}{\pi} G(p, q) + \frac{2}{\pi} = \frac{1}{\pi} \log p^{q-1} q^{p-1} - (V(p, q) + V(q, p)) - pg\left(\frac{1}{p}\right) - qg\left(\frac{1}{q}\right).$$

This implies relation (6.2). □

## References

1. M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Function with Formula, Graphs and Mathematical Tables*, tenth printing with corrections, US Department of Commerce, New York, 1972.
2. L. Baez-Duarte, *On Beurling real variable reformulation of the Riemann hypothesis*, Adv. Math. **101** (1993), 10–30.
3. ———, *News versions of the Nyman-Beurling criterion for the Riemann hypothesis*, Int. J. Math. Math. Sci. **31**(7) (2002), 387–406.
4. ———, *A strengthening of the Nyman-Beurling criterion for the a Riemann hypothesis*, Rend. Mat. Acc. Lincei (s9) **14** (2003), 5–11.
5. ———, *A general strong Nyman-Beurling criterion for the Riemann hypothesis*, Publ. Inst. Math., Nouv. Sér. **78**(92) (2005), 117–125.
6. L. Baéz-Duarte, M. Balazard, B. Landreau, E. Saias, *Etude de l'autocorrélation multiplicative de la fonction partie fractionnaire*, Ramanujan J. **9**(1–2) (2005), 215–240.
7. ———, *Notes sur la fonction  $\zeta$  de Riemann, 3*, Adv. Math. **149**(1) (2000), 130–144.
8. M. Balazard, *Sur les dilatations entières de la fonction partie fractionnaire*, Funct. Approximatio, Comment. Math. **35** (2006), 37–49.
9. M. Balazard, A. de Roton, *Sur un critère de Baez-Duarte pour l'hypothèse de Riemann*, Int. J. Number Theory **6**(4) (2010), 883–903.
10. A. Bayad, M. Goubi, *Proof of the Möbius conjecture revisited*, Proc. Jangjeon Math. Soc. **16**(2) (2013), 237–243.
11. P. Barrucand, M. Deboué, *Fraction continues, sommes de Dedekind et formes quadratiques*, Rend. Circ. Mat. Palermo (2) **33**(1) (1984), 62–84.
12. S. Bettin, J. B. Conrey, *Period functions and cotangent sums*, Algebra Number Theory **7**(1) (2013), 215–242.
13. ———, *A reciprocity formula for a cotangent sum*, Int. Math. Res. Not. **2013**(24) (2013), 5709–5726.
14. S. Gun, M. Ram Murty, P. Rath, *Linear independence of digamma function and a variant of a conjecture of Rohrlich*, J. Number Theory **129**(8) (2009), 1858–1873.
15. D. Hickerson, *Continued fractions and density results for Dedekind sums*, J. Reine Angew. Math. **290** (1977), 113–116.
16. K. Girstmair, *Continued fractions and Dedekind sums: three term relations and distribution*, J. Number Theory **119**(1) (2006), 66–85.
17. M. Ishibashi, *The value of the Estermann zeta function at  $s = 0$* , Acta Arith. **73**(4) (1995), 357–361.
18. B. Landreau, F. Richard, *Le critère de Beurling et Nyman pour l'hypothèse de Riemann: aspects numériques*, Exp. Math. **11**(3) (2002), 349–360.
19. L. A. Medina, V. H. Moll, *The integrals in Gradshteyn and Ryzhik. Part 10: The digamma function*, Sci., Ser. A, Math. Sci. (N.S.) **17** (2009), 45–66.
20. H. Maier, M. Th. Rassias, *The order of magnitude for moments for certain cotangent sums*, J. Math. Anal. Appl. **429**(1) (2015), 576–590.
21. ———, *Generalizations of a cotangent sum associated to the Estermann zeta function*, Commun. Contemp. Math. **18** (2016), 1550078, 89 pp.
22. ———, *The rate of growth of moments of certain cotangent sums*, Aequationes Math. (2015), DOI 10.1007/s00010-015-0361-3.
23. ———, *Asymptotics and equidistribution of cotangent sums associated to the Estermann and Riemann zeta functions*, in: J. Sander, J. Steuding, R. Steuding (Eds.), *From Arithmetic to Zeta-Functions*, Springer, Cham., 2016.
24. M. Ram Murty, N. Saradha, *Transcendental values of the digamma function*, J. Number Theory **125**(2) (2007), 298–318.
25. I. Niven, H. S. Zuckerman, *An Introduction to the Theory of Numbers*, 3rd ed., New York, 1972

26. M. Th. Rassias, *A cotangent sum related to the zeros of the Estermann zeta function*, Appl. Math. Comput. **240** (2014), 161–167.
27. W. T. Sulaiman, *Turan inequalities for the digamma and polygamma functions*, South Asian J. Math. **1**(2) (2011), 49–55.
28. V. I. Vasyunin, *On a biorthogonal system associated with the Riemann hypothesis*, Algebra Anal. **7**(3) (1995), 118–135.

Laboratoire d'Algèbre et Théorie des Nombres  
Department of Mathematics  
University of UMMTO  
Tizi-ouzou  
Algeria  
mouloud.ummto@hotmail.fr

(Received 29 12 2015)

(Revised 06 02 2016)

LAMME, Univ. Evry  
Université Paris-Saclay  
Evry  
France  
abayad@maths.univ-evry.fr

Département d'Algèbre et Théorie des Nombres  
Faculté de Mathématiques  
Université des Sciences et de la technologie Houari-Boumediène (USTHB)  
Alger  
Algérie  
mhernane@usthb.dz