

INTEGRAL REPRESENTATIONS OF GRAPHS *

FERNANDO C. SILVA

Presented by J.A. Dias da Silva

Abstract: Following the definition of graph representation modulo an integer given by Erdős and Evans in [1], we call degree of a representation to the number of prime factors in the prime factorization of its modulo. Here we study the smallest possible degree for a representation of a graph.

The starting point for this research is the concept of representation introduced in [1], and the proposed study of relations between properties of graphs and properties of their representations.

Let $G = (V, E)$ be a graph with n vertices v_1, \dots, v_n . The graph G is said to be *representable* modulo a positive integer b if there exist distinct integers a_1, \dots, a_n such that $0 \leq a_i < b$, and $\text{g.c.d.}\{a_i - a_j, b\} = 1$ if and only if v_i and v_j are adjacent. We say that $\{a_1, \dots, a_n\}$ is a *representation* of G modulo b . We call *degree* of the representation to the number of prime factors, counting multiplicities, in the prime factorization of b . The concept of degree was not mentioned in [1] explicitly. However we can see in the proof of the theorem of [1] that there always exists a representation of degree equal to the number of edges of the complement of a graph that results from G by adjoining an isolated vertex. We shall see that there exist representations of smaller degree. We call *representation degree* of G , $d_r(G)$, to the smallest possible degree for a representation of G .

We say that a function $\phi: E \rightarrow X$ is *transitive* if, for every $(v_i, v_j), (v_j, v_k) \in E$ such that $\phi(v_i, v_j) = \phi(v_j, v_k) = x$, we have $(v_i, v_k) \in E$ and $\phi(v_i, v_k) = x$. For example, if $\phi: E \rightarrow X$ is one-to-one, then ϕ is transitive. Given a set Y , $\#Y$

Received: March 31, 1995.

* This work was done within the activities of the Centro de Álgebra da Universidade de Lisboa.

denotes its cardinal number. We call *degree* of a transitive function $\phi: E \rightarrow X$ to $\#\phi(E)$ and we call *transitive degree* of G , $d_t(G)$, to the smallest $\#\phi(E)$, when ϕ runs over the transitive functions defined in E . It is not difficult to prove some properties of $d_t(G)$. For example:

Proposition 1. $d_t(G) \leq \#E$.

Proposition 2. $d_t(G) = \max_H d_t(H)$, where H runs over the maximal connected subgraphs of G .

Proposition 3. Suppose that G is connected. Then

- a) $d_t(G) = 0$ if and only if $\#V = 1$.
- b) $d_t(G) = 1$ if and only if $\#V \geq 2$ and G is complete.
- c) $d_t(G) = \#E$ if and only if there exists a vertex incident with all the edges of G .

Let $G' = (V', E')$ be the complement of G . The following theorems are our main results. We shall prove them later.

Theorem 4. Let ϕ be a transitive function defined in E' of degree $d \geq 2$. Then there exists a representation of G of degree d .

Corollary 5. If $d_t(G') \geq 2$, then $d_r(G) \leq d_t(G') \leq \#E'$.

Corollary 5 is not always true when $d_t(G') \leq 1$. The following proposition shows this and is easy to prove.

Proposition 6.

- a) $d_r(G) = 0$ if and only if $\#V = 1$.
- b) $d_r(G) = 1$ if and only if $\#V \geq 2$ and G is complete.
- c) $d_r(G) \leq 1$ if and only if $d_t(G') = 0$.
- d) If $d_t(G') = 1$, then $d_r(G) = 2$.

Theorem 7. Suppose that G' does not have any subgraph isomorphic to K_3 . If G has a representation of degree d , then there exists a transitive function defined in E' of degree $\leq d$.

Counter-example. If G' has subgraphs isomorphic to K_3 , then Theorem 7 is not always true, as the following example shows. Suppose that G is a graph with 5 vertices and only one edge. Then $R = \{0, 3, 5, 15, 30\}$ is a representation of G modulo $b = 3 \times 5 \times 7 = 105$. It is not difficult to see that any transitive function defined in E' has degree greater than 3.

Corollary 8. *If $d_t(G') \geq 2$ and G' does not have any subgraph isomorphic to K_3 , then $d_r(G) = d_t(G')$.*

Let $M(G')$ be the maximum number of edges incident with one vertex in G' .

Theorem 9. *Suppose that G' has no cycles. Then*

a) $d_t(G') = M(G')$.

b) *If at least one of the maximal connected subgraphs of G' has at least 3 vertices, then*

$$(1) \quad d_r(G) = d_t(G') = M(G') .$$

Corollary 10. *If G' is a tree and $n \neq 2$, then (1) holds.*

Now we are going to prove the theorems above. We split the proof of Theorem 4 into several lemmas.

Lemma 11. *Suppose that $\phi: E' \rightarrow X$ is a transitive function with $\#\phi(E')=1$. Let δ be a positive integer. Then there exists a positive prime $p > \delta$ and there exist distinct nonnegative integers a_1, \dots, a_n such that $(v_i, v_j) \in E'$ if and only if p divides $a_i - a_j$, $i, j \in \{1, \dots, n\}$, $i \neq j$.*

Proof: Let H_1, \dots, H_t be the maximal connected subgraphs of G' . Without loss of generality, suppose that $H_s = \{v_{k_1+\dots+k_{s-1}+1}, \dots, v_{k_1+\dots+k_s}\}$, $k_s = \#H_s$, $s \in \{1, \dots, t\}$. Let p be a prime $> \max\{t, \delta\}$. If $i = k_1 + \dots + k_{s-1} + j$, $1 \leq j \leq k_s$, let $a_i = s + jp$. Since $\#\phi(E') = 1$, the graphs H_i are complete. It is easy to conclude that the lemma is satisfied. ■

Lemma 12. *Let α and β be integers with $\text{g.c.d.}\{\alpha, \beta\} = 1$. Let p be a prime. Then there exists at most one $\epsilon \in \{0, \dots, p-1\}$ such that $\epsilon\beta + \alpha \in (p)$, where (p) denotes the principal ideal, of the ring of the integers, generated by p .*

Proof: Firstly, suppose that p divides β . Then p does not divide α and, therefore, $\epsilon\beta + \alpha \notin (p)$, for every integer ϵ . Now suppose that p does not divide β

and that there exist $\epsilon_1, \epsilon_2 \in \{1, \dots, p-1\}$ such that $\epsilon_1 \neq \epsilon_2$ and $\epsilon_1\beta + \alpha, \epsilon_2\beta + \alpha \in (p)$. Then $(\epsilon_1 - \epsilon_2)\beta \in (p)$. As p is prime, p divides $\epsilon_1 - \epsilon_2$ or p divides β , what is impossible. ■

Lemma 13. *Let $\alpha_1, \dots, \alpha_s, \beta_1, \dots, \beta_s$ be integers such that $\text{g.c.d.}\{\alpha_j, \beta_j\} = 1$, $j \in \{1, \dots, s\}$. Let $b = p_1 \cdots p_r$, where p_1, \dots, p_r are positive primes. If $\min\{p_i : 1 \leq i \leq r\} > sr$, then there exists an integer γ such that*

$$(2) \quad \text{g.c.d.}\{\gamma\beta_j + \alpha_j, b\} = 1, \quad j \in \{1, \dots, s\}.$$

Proof: Let $m = \min\{p_i\}$. From the previous lemma, it can easily be deduced that there exists $\gamma \in \{0, \dots, m-1\}$ such that $\gamma\beta_j + \alpha_j \notin (p_i)$, $j \in \{1, \dots, s\}$, $i \in \{1, \dots, r\}$. That is, γ satisfies (2). ■

Lemma 14. *Let $\phi : E' \rightarrow X$ be a transitive function. Suppose that $d = \#\phi(E') \geq 2$ and $\phi(E') = \{x_1, \dots, x_d\}$. Let δ be a positive integer. Then there exist distinct positive primes p_1, \dots, p_d and there exist distinct integers a_1, \dots, a_n such that:*

- i) $0 \leq a_i < p_1 \cdots p_d$, $i \in \{1, \dots, n\}$.
- ii) $\text{g.c.d.}\{a_i - a_j, p_1 \cdots p_d\} = 1$ if and only if $(v_i, v_j) \notin E'$, $i, j \in \{1, \dots, n\}$, $i \neq j$.
- iii) $\text{g.c.d.}\{a_i - a_j, p_1 \cdots p_d\} = p_u$ if and only if $(v_i, v_j) \in E'$, and $\phi(v_i, v_j) = x_u$, $i, j \in \{1, \dots, n\}$, $i \neq j$, $u \in \{1, \dots, d\}$.
- iv) $\min\{p_1, \dots, p_d\} > \delta$.

Proof: By induction on n . As $d \geq 2$, we have $n \geq 3$. Let $G_0 = (V_0, E_0)$ be the subgraph that we obtain from G' deleting v_n and all the edges incident with v_n . Without loss of generality, we assume that $E_0 \neq E'$ and $\phi(E_0) = \{x_1, \dots, x_e\}$. We choose p_1, \dots, p_e and a_1, \dots, a_{n-1} as follows. Note that $e \leq 1$ when $n = 3$.

If $e \geq 2$, then, by the induction assumption, there exist distinct primes p_1, \dots, p_e and there exist distinct integers a_1, \dots, a_{n-1} such that:

- i₀) $0 \leq a_i < p_1 \cdots p_e$, $i \in \{1, \dots, n-1\}$.
- ii₀) $\text{g.c.d.}\{a_i - a_j, p_1 \cdots p_e\} = 1$ if and only if $(v_i, v_j) \notin E_0$, $i, j \in \{1, \dots, n-1\}$, $i \neq j$.
- iii₀) $\text{g.c.d.}\{a_i - a_j, p_1 \cdots p_e\} = p_u$ if and only if $(v_i, v_j) \in E_0$, and $\phi(v_i, v_j) = x_u$, $i, j \in \{1, \dots, n-1\}$, $i \neq j$, $u \in \{1, \dots, e\}$.
- iv₀) $\min\{p_1, \dots, p_e\} > \max\{\delta, (n-1)d\}$.

If $e = 1$, then, according to Lemma 11, there exists a prime p_1 and there exist distinct nonnegative integers a_1, \dots, a_{n-1} satisfying ii₀), iii₀) and iv₀).

If $e = 0$, take $a_i = i - 1$, $i \in \{1, \dots, n - 1\}$.

In any case $e \geq 0$, we choose primes p_{e+1}, \dots, p_d such that:

- I) p_1, \dots, p_d are distinct.
- II) None of the primes p_{e+1}, \dots, p_d divide $a_i - a_j$, $i, j \in \{1, \dots, n - 1\}$, $i \neq j$.
- III) $\min\{p_1, \dots, p_d\} > \max\{\delta, (n - 1)d\}$.

Without loss of generality, suppose that v_1, \dots, v_t are the vertices of G' incident with v_n . Let $x_{k_i} = \phi(v_i, v_n)$, $i \in \{1, \dots, t\}$. Without loss of generality, suppose that k_1, \dots, k_r are pairwise distinct and $k_i \in \{k_1, \dots, k_r\}$ whenever $i \in \{r + 1, \dots, t\}$.

According to the Chinese Remainder Theorem, there exists an integer z such that

$$(3) \quad z - a_j \in (p_{k_j}), \quad j \in \{1, \dots, r\} .$$

Let $i \in \{r + 1, \dots, t\}$ and suppose that $k_i = k_j$, where $j \in \{1, \dots, r\}$. As ϕ is transitive, $(v_i, v_j) \in E'$ and $\phi(v_i, v_j) = x_{k_i}$. Therefore $k_i \in \{1, \dots, e\}$. From iii₀), it follows that $a_i - a_j \in (p_{k_i})$. Thus $z - a_i = (z - a_j) + (a_j - a_i) \in (p_{k_i})$.

Now suppose that $z - a_i \in (p_{k_j})$, with $i \in \{1, \dots, n - 1\}$, $j \in \{1, \dots, r\}$. From (3), $a_i - a_j \in (p_{k_j})$. Bearing in mind II), ii₀) and iii₀), we conclude that $k_j \in \{1, \dots, e\}$, $(v_i, v_j) \in E_0$ and $\phi(v_i, v_j) = x_{k_j}$. From the transitivity of ϕ , $(v_i, v_n) \in E'$ and $\phi(v_i, v_n) = x_{k_j}$. Therefore, $i \in \{1, \dots, t\}$ and $k_i = k_j$.

It is not difficult to prove that

$$(4) \quad \text{g.c.d.}\{p_{k_1} \cdots p_{k_r}, z - a_i\} = p_{k_i}, \quad i \in \{1, \dots, t\} ,$$

$$(5) \quad \text{g.c.d.}\{p_{k_1} \cdots p_{k_r}, z - a_i\} = 1, \quad i \in \{t + 1, \dots, n - 1\} .$$

Using Lemma 13, it follows from (4), (5) and III) that there exists an integer γ such that

$$(6) \quad \text{g.c.d.}\left\{\gamma \frac{p_{k_1} \cdots p_{k_r}}{p_{k_i}} + \frac{z - a_i}{p_{k_i}}, b\right\} = 1, \quad i \in \{1, \dots, t\} ,$$

$$(7) \quad \text{g.c.d.}\{\gamma p_{k_1} \cdots p_{k_r} + z - a_i, b\} = 1, \quad i \in \{t + 1, \dots, n - 1\} ,$$

where $b = p_1 \cdots p_d$. Let $a_n = \gamma p_{k_1} \cdots p_{k_r} + z + wb$, where w is an integer chosen so that $0 \leq a_n < b$. Then (6) and (7) take the forms

$$\begin{aligned} \text{g.c.d.}\{a_n - a_i, b\} &= p_{k_i}, \quad i \in \{1, \dots, t\} , \\ \text{g.c.d.}\{a_n - a_i, b\} &= 1, \quad i \in \{t + 1, \dots, n - 1\} . \end{aligned}$$

Clearly a_n is different from a_i , $i \in \{1, \dots, n-1\}$, and conditions i)–iv) are satisfied. ■

Now Theorem 4 follows immediately from Lemma 14.

Proof of Theorem 7: Let $R = \{a_1, \dots, a_n\}$ be a representation of G modulo $b = p_1 \cdots p_d$, where p_1, \dots, p_d are primes. Suppose that a_1, \dots, a_n are ordered so that $\text{g.c.d.}\{a_i - a_j, b\} = 1$ if and only if $(v_i, v_j) \in E$. For each $(v_i, v_j) \in E'$, let $\phi(v_i, v_j)$ be an element of $\{p_1, \dots, p_d\}$ such that $\phi(v_i, v_j)$ divides $a_i - a_j$. It is easy to see that $\phi: E' \rightarrow \{p_1, \dots, p_d\}$ is transitive. ■

Proof of Theorem 9: a) For each $i \in \{1, \dots, n\}$, we denote by $E_i(G')$ the set of all the edges incident with v_i in G' . Given a transitive function $\phi: E' \rightarrow X$, the restriction of ϕ to $E_i(G')$ is one-to-one. Therefore, $\#\phi(E') \geq \#E_i(G')$. Consequently, $d_t(G') \geq \max\{\#E_i(G')\}$.

Now we prove that $d_t(G') \leq \max\{\#E_i(G')\}$ by induction on $\#E'$. If E' is empty, this is trivial. Suppose that $\#E' \geq 1$. Then there exists $i \in \{1, \dots, n\}$ such that $\#E_i(G') = 1$. Without loss of generality, assume that $E_n(G') = \{(v_{n-1}, v_n)\}$. Let $\tilde{G} = (V, \tilde{E})$, where $\tilde{E} = E' \setminus \{(v_{n-1}, v_n)\}$. By the induction assumption, $d_t(\tilde{G}) \leq \max\{\#E_i(\tilde{G})\}$. Let $\psi: \tilde{E} \rightarrow X$ be a transitive function of degree $d_t(\tilde{G})$. If there exists $i \in \{1, \dots, n-2\}$ such that $\#E_{n-1}(\tilde{G}) < \#E_i(\tilde{G}) (= \#E_i(G'))$, let x be an element of $\psi(E_i(\tilde{G})) \setminus \psi(E_{n-1}(\tilde{G}))$. If $\#E_i(\tilde{G}) \leq \#E_{n-1}(\tilde{G})$, $i \in \{1, \dots, n-2\}$, let x be an element that does not belong to X . Let $\phi: E' \rightarrow X \cup \{x\}$ be the extension function of ψ satisfying $\phi(v_{n-1}, v_n) = x$. It is easy to see that ϕ is transitive and

$$d_t(G') \leq \#\phi(E') \leq \max\{\#E_i(G')\}.$$

b) Since G' is acyclic, the hypothesis of b) is equivalent to $d_t(G') \geq 2$. Thus b) follows from a) and Corollary 8. ■

REFERENCES

- [1] ERDÖS, P. and EVANS, A.B. – Representations of graphs and ortogonal latin square graphs, *J. Graph Theory*, 13 (1989), 593–595.

Fernando C. Silva,
Departamento de Matemática, Faculdade de Ciências, Universidade de Lisboa,
Rua Ernesto de Vasconcelos, 1700 Lisboa – Portugal