

A Quasi-Lie Schemes Approach to Second-Order Gambier Equations*

José F. CARIÑENA [†], Partha GUHA [‡] and Javier DE LUCAS [§]

[†] Department of Theoretical Physics and IUMA, University of Zaragoza,
Pedro Cerbuna 12, 50.009, Zaragoza, Spain
E-mail: jfc@unizar.es

[‡] S.N. Bose National Centre for Basic Sciences, JD Block, Sector III,
Salt Lake, Kolkata - 700.098, India
E-mail: partha@bose.res.in

[§] Faculty of Mathematics and Natural Sciences, Cardinal Stefan Wyszyński University,
Wóycickiego 1/3, 01-938, Warsaw, Poland
E-mail: j.delucasaraujo@uksw.edu.pl

Received September 26, 2012, in final form March 14, 2013; Published online March 26, 2013
<http://dx.doi.org/10.3842/SIGMA.2013.026>

Abstract. A *quasi-Lie scheme* is a geometric structure that provides t -dependent changes of variables transforming members of an associated family of systems of first-order differential equations into members of the same family. In this note we introduce two quasi-Lie schemes for studying second-order Gambier equations in a geometric way. This allows us to study the transformation of these equations into simpler canonical forms, which solves a gap in the previous literature, and other relevant differential equations, which leads to derive new constants of motion for families of second-order Gambier equations. Additionally, we describe general solutions of certain second-order Gambier equations in terms of particular solutions of Riccati equations, linear systems, and t -dependent frequency harmonic oscillators.

Key words: Lie system; Kummer–Schwarz equation; Milne–Pinney equation; quasi-Lie scheme; quasi-Lie system; second-order Gambier equation; second-order Riccati equation; superposition rule

2010 Mathematics Subject Classification: 34A26; 34A05; 34A34; 17B66; 53Z05

1 Introduction

Apart from their inherent mathematical interest, differential equations are important due to their use in all branches of science [41, 50]. This strongly motivates their analysis as a means to study the problems they model. A remarkable approach to differential equations is given by geometric methods [47], which have resulted in powerful techniques such as Lax pairs, Lie symmetries, and others [48, 49].

A particular class of systems of ordinary differential equations that have been drawing some attention in recent years are the so-called *Lie systems* [1, 9, 33, 46, 54, 56]. Lie systems form a class of systems of first-order differential equations possessing a *superposition rule*, i.e. a function that enables us to write the general solution of a first-order system of differential equations in terms of a generic collection of particular solutions and some constants to be related to initial conditions [7, 18].

The theory of Lie systems furnishes many geometric methods for studying these systems [6, 17, 19, 21, 22, 26, 27, 56]. For instance, superposition rules can be employed to simplify the use

*This paper is a contribution to the Special Issue “Symmetries of Differential Equations: Frames, Invariants and Applications”. The full collection is available at <http://www.emis.de/journals/SIGMA/SDE2012.html>

of numerical techniques for solving differential equations [56], and the theory of reduction of Lie systems reduces the integration of Lie systems on Lie groups to solving Lie systems on simple Lie groups [6, 19].

The classification of systems admitting a superposition rule is due to Lie. His result, the nowadays called Lie–Scheffers theorem, states that a system admits a superposition rule if and only if it describes the integral curves of a t -dependent vector field taking values in a finite-dimensional Lie algebra of vector fields. The existence of such Lie algebras on \mathbb{R} and \mathbb{R}^2 was analysed by Lie in his famous work [45]. More recently, the topic was revisited by Olver and coworkers [28], who clarified a number of details that were not properly described in the previous literature.

Despite their interesting properties, Lie systems have a relevant drawback: there exist just a few Lie systems of broad interest [36]. Indeed, the Lie–Scheffers theorem and, more specifically, the classification of finite-dimensional Lie algebras of vector fields on low dimensional manifolds [28, 45] clearly show that that being a Lie system is the exception rather than the rule. This has led to generalise the theory of Lie systems so as to tackle a larger family of remarkable systems [2, 3, 9, 36]. In particular, we henceforth focus on the so-called *quasi-Lie schemes*. These recently devised structures [15, 20] have been found quite successful in investigating transformation and integrability properties of differential equations, e.g. Abel equations, dissipative Milne–Pinney equations, second-order Riccati equations, and others [9]. In addition, the obtained results are useful so as to research on the physical and mathematical problems described through these equations.

In this work, we study the second-order Gambier equations by means of the theory of quasi-Lie schemes. We provide two new quasi-Lie schemes. Their associated groups [15] give rise to groups of t -dependent changes of variables, which are used to transform second-order Gambier equations into another ones. Such groups allow us to explain in a geometric way the existence of certain transformations reducing a quite general subclass of second-order Gambier equations into simpler ones. Our approach provides a better understanding of a result pointed out in [30]. As a byproduct, we show that the procedure given in the latter work does not apply to every second-order Gambier equation, which solves a gap performed in there.

We provide conditions for second-order Gambier equations, written as first-order systems, to be mapped into Lie systems via t -dependent changes of variables induced by our quasi-Lie schemes. This is employed to determine families of Gambier equations which can be transformed into second-order Riccati equations [31], Kummer–Schwarz equations [4, 5] and Milne–Pinney equations [25]. These results are employed to derive, as far as we know, new constants of motion for certain second-order Gambier equations. Moreover, the description of their general solutions in terms of particular solutions of t -dependent frequency harmonic oscillators, linear systems, or Riccati equations is provided [9, 29].

The structure of our paper goes as follows. Section 2 addresses the description of the fundamental notions to be employed throughout our work. Section 3 describes a new quasi-Lie scheme for studying second-order Gambier equations. In Section 4 this quasi-Lie scheme is used to analyse the reduction of second-order Gambier equations to a simpler canonical form [30]. By using the theory of quasi-Lie systems, we determine in Section 5 a family of second-order Gambier equations that can be mapped into second-order Kummer–Schwarz equations. The investigation of constants of motion for some members of the previous family is performed in Section 6. In Section 7 we describe the general solutions of a family of second-order Gambier equations in terms of particular solutions of other differential equations. We present a second quasi-Lie scheme for investigating second-order Gambier equations in Section 8, and conditions are given to be able to transform these equations into second-order Riccati equations. Those second-order Gambier equations that can be transformed into second-order Riccati equations are integrated in Section 9. Finally, Section 10 is devoted to summarising our main results.

2 Fundamentals

Let us survey the fundamental results to be used throughout the work (see [15, 16, 17, 18] for details). In general, we hereafter assume all objects to be smooth, real, and globally defined on linear spaces. This simplifies our exposition and allows us to avoid tackling minor details.

Given the projection $\pi : (t, x) \in \mathbb{R} \times \mathbb{R}^n \mapsto x \in \mathbb{R}^n$ and the tangent bundle projection $\tau : T\mathbb{R}^n \rightarrow \mathbb{R}^n$, a *t-dependent vector field* on \mathbb{R}^n is a mapping $X : \mathbb{R} \times \mathbb{R}^n \rightarrow T\mathbb{R}^n$ such that $\tau \circ X = \pi$. This condition entails that every *t-dependent vector field* X gives rise to a family $\{X_t\}_{t \in \mathbb{R}}$ of vector fields $X_t : x \in \mathbb{R}^n \mapsto X(t, x) \in T\mathbb{R}^n$ and vice versa. We call *minimal Lie algebra* of X the smallest real Lie algebra V^X containing the vector fields $\{X_t\}_{t \in \mathbb{R}}$. Given a finite-dimensional \mathbb{R} -linear space V of vector fields on \mathbb{R}^n , we write $V(C^\infty(\mathbb{R}))$ for the $C^\infty(\mathbb{R})$ -module of *t-dependent vector fields* taking values in V .

An *integral curve* of X is a standard integral curve $\gamma : \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}^n$ of its *suspension*, i.e. the vector field $\bar{X} = \partial/\partial t + X(t, x)$ on $\mathbb{R} \times \mathbb{R}^n$. Note that the integral curves of X of the form $\gamma : t \in \mathbb{R} \rightarrow (t, x(t)) \in \mathbb{R} \times \mathbb{R}^n$ are the solutions of the system

$$\frac{dx^i}{dt} = X^i(t, x), \quad i = 1, \dots, n, \quad (2.1)$$

the referred to as *associated system* of X . Conversely, given such a system, we can define a *t-dependent vector field* on \mathbb{R}^n [16]

$$X(t, x) = \sum_{i=1}^n X^i(t, x) \frac{\partial}{\partial x^i}$$

whose integral curves of the form $(t, x(t))$ are the solutions to (2.1). This justifies to write X for both a *t-dependent vector field* and its associated system.

We call *generalised flow* a map $g : (t, x) \in \mathbb{R} \times \mathbb{R}^n \mapsto g_t(x) \in \mathbb{R}^n$ such that $g_0 = \text{Id}_{\mathbb{R}^n}$. Every *t-dependent vector field* X can be associated with a generalised flow g satisfying that the general solution of X can be written in the form $x(t) = g_t(x_0)$ with $x_0 \in \mathbb{R}^n$. Conversely, every generalised flow defines a vector field by means of the expression [15]

$$X(t, x) = \left. \frac{d}{ds} \right|_{s=t} g_s \circ g_t^{-1}(x).$$

Generalised flows act on *t-dependent vector fields* [18]. More precisely, given a generalised flow g and a *t-dependent vector field* X , we can define a unique *t-dependent vector field*, $g_\star X$, whose associated system has general solution $\bar{x}(t) = g_t(x(t))$, where $x(t)$ is the general solution of X . In other words, every g induces a *t-dependent change of variables* $\bar{x}(t) = g_t(x(t))$ transforming the system X into $g_\star X$. Indeed, g can be viewed as a diffeomorphism $\bar{g} : (t, x) \in \mathbb{R} \times \mathbb{R}^n \mapsto (t, g_t(x)) \in \mathbb{R} \times \mathbb{R}^n$, and it can easily be proved that $g_\star X$ is the only *t-dependent vector field* such that $\overline{g_\star X} = \bar{g}_* \bar{X}$, where \bar{g}_* is the standard action of the diffeomorphism \bar{g} on vector fields (see [18]).

Among all *t-dependent vector fields*, we henceforth focus on those whose associated systems are Lie systems. The characteristic property of Lie systems is to possess a superposition rule [7, 18, 46]. A *superposition rule* for a system X on \mathbb{R}^n is a map $\Phi : (u_{(1)}, \dots, u_{(m)}; k_1, \dots, k_n) \in (\mathbb{R}^n)^m \times \mathbb{R}^n \mapsto \Phi(u_{(1)}, \dots, u_{(m)}; k_1, \dots, k_n) \in \mathbb{R}^n$ allowing us to write its general solution $x(t)$ as

$$x(t) = \Phi(x_{(1)}(t), \dots, x_{(m)}(t); k_1, \dots, k_n),$$

for a generic family of particular solutions $x_{(1)}(t), \dots, x_{(m)}(t)$ and a set of constants k_1, \dots, k_n to be related to initial conditions.

The celebrated Lie–Scheffers theorem [46, Theorem 44] states that a system X possesses a superposition rule if and only if it is a t -dependent vector field taking values in a finite-dimensional real Lie algebra of vector fields, termed *Vessiot–Guldberg Lie algebra* [34, 53]. In other words, X is a Lie system if and only if V^X is finite-dimensional [9]. This is indeed the main reason to define V^X [14].

To illustrate the above notions, let us consider the Riccati equation [35]

$$\frac{dx}{dt} = b_1(t) + b_2(t)x + b_3(t)x^2, \quad (2.2)$$

where $b_1(t), b_2(t), b_3(t)$ are arbitrary functions of time. Its general solution, $x(t)$, can be obtained from an expression [38, 56]

$$x(t) = \Phi(x_{(1)}(t), x_{(2)}(t), x_{(3)}(t); k),$$

where k is a real number to be related to the initial conditions of every particular solution, $x_{(1)}(t), x_{(2)}(t), x_{(3)}(t)$ are three different particular solutions of (2.2) and $\Phi : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$\Phi(u_{(1)}, u_{(2)}, u_{(3)}; k) = \frac{u_{(1)}(u_{(2)} - u_{(3)}) - ku_{(2)}(u_{(3)} - u_{(1)})}{(u_{(2)} - u_{(3)}) - k(u_{(3)} - u_{(1)})}.$$

That is, the Riccati equations admit a superposition rule. Therefore, from the Lie–Scheffers Theorem, we infer that the t -dependent vector field X associated to a Riccati equation is such that V^X is finite-dimensional. Indeed,

$$X = (b_1(t) + b_2(t)x + b_3(t)x^2) \frac{\partial}{\partial x}.$$

Taking into account that $X_1 = \partial/\partial x$, $X_2 = x\partial/\partial x$, $X_3 = x^2\partial/\partial x$ span a finite-dimensional real Lie algebra V of vector fields and $X_t = b_1(t)X_1 + b_2(t)X_2 + b_3(t)X_3$, we obtain that $\{X_t\}_{t \in \mathbb{R}} \subset V$ and V^X becomes a (finite-dimensional) Lie subalgebra of V .

The Lie–Scheffers theorem shows that just some few first-order systems are Lie systems [9, 36]. For instance, this theorem implies that all Lie systems on the real line are, up to a change of variables, a particular case of a linear or Riccati equation [55]. Therefore, many other important differential equations cannot be studied through Lie systems (see [9] for examples of this). In order to treat non-Lie systems, new techniques generalising Lie systems need to be developed. We here focus on the theory of quasi-Lie schemes [15, 20].

Definition 2.1. Let W, V be finite-dimensional real vector spaces of vector fields on \mathbb{R}^n . We say that they form a *quasi-Lie scheme* $S(W, V)$ if:

- W is a vector subspace of V .
- W is a Lie algebra of vector fields, i.e. $[W, W] \subset W$.
- W normalises V , i.e. $[W, V] \subset V$.

Associated to each quasi-Lie scheme, we have the $C^\infty(\mathbb{R})$ -modules $W(C^\infty(\mathbb{R}))$ and $V(C^\infty(\mathbb{R}))$ of t -dependent vector fields taking values in W and V , respectively. Now, from the Lie algebra W , we define the group $\mathcal{G}(W)$ of generalised flows of t -dependent vector fields taking values in W , the so-called *group of the scheme*. The relevance of this group is due to the following theorem [15].

Theorem 2.1. *Given a quasi-Lie scheme $S(W, V)$, every generalised flow of $\mathcal{G}(W)$ acts transforming elements of $V(C^\infty(\mathbb{R}))$ into members of $V(C^\infty(\mathbb{R}))$.*

In other words, the elements of the group of a scheme provide t -dependent changes of variables that transform systems of $V(C^\infty(\mathbb{R}))$ into systems of this family. Roughly speaking, we can understand this group as a generalisation of the t -independent symmetry group of a system: apart from transforming the initial system into itself, the transformations of the group of a scheme also may transform the initial system into one of the “same type”. For instance, given a Lie system X associated with a Vessiot–Guldberg Lie algebra V , then $S(V, V)$ becomes a quasi-Lie scheme. The group $\mathcal{G}(V)$ allows us to transform X into a Lie system with a Vessiot–Guldberg Lie algebra V . This can be employed, for example, to transform Riccati equations into Riccati equations that can be easily integrated, giving rise to methods to integrate Riccati equations.

In order to illustrate previous notions, we now turn to proving that quasi-Lie schemes allow us to cope with Abel equations of first-order and first kind [11], i.e.

$$\frac{dx}{dt} = b_1(t) + b_2(t)x + b_3(t)x^2 + b_4(t)x^3, \quad (2.3)$$

with $b_1(t), \dots, b_4(t)$ being arbitrary t -dependent functions. Indeed, if we fix $W = \langle \partial/\partial x, x\partial/\partial x \rangle$, it can be proved that $S(W, V)$ and $V = \langle \partial/\partial x, x\partial/\partial x, x^2\partial/\partial x, x^3\partial/\partial x \rangle$ is a quasi-Lie scheme and $X \in V(C^\infty(\mathbb{R}))$ for every X related to an Abel equation (2.3). The elements of $\mathcal{G}(W)$ transform Abel equations into Abel equations and geometrically recover the usual t -dependent changes of variables used to study these equations. This was employed in [11] to describe integrability properties of Abel equations.

Given a quasi-Lie scheme $S(W, V)$, certain systems in $V(C^\infty(\mathbb{R}))$ can be mapped into Lie systems admitting a Vessiot–Guldberg Lie algebra contained in V . This enables us to study the transformed system through techniques from the theory of Lie systems and, undoing the performed transformation, to obtain properties of the initial system under study [15].

Definition 2.2. Let $S(W, V)$ be a quasi-Lie scheme and X a t -dependent vector field in $V(C^\infty(\mathbb{R}))$, we say that X is a *quasi-Lie system with respect to $S(W, V)$* if there exists a generalised flow $g \in \mathcal{G}(W)$ and a Lie algebra of vector fields $V_0 \subset V$ such that $g_\star X \in V_0(C^\infty(\mathbb{R}))$.

3 A new quasi-Lie scheme for investigating second-order Gambier equations

The *Gambier equation* [30, 32] can be described as the coupling of two Riccati equations in cascade, which can be given in the following form

$$\begin{aligned} \frac{dy}{dt} &= -y^2 + a_1y + a_2, \\ \frac{dx}{dt} &= a_0x^2 + nyx + \sigma, \end{aligned}$$

where n is an integer, σ is a constant, which can be scaled to 1 unless it happens to be 0, and a_0, a_1, a_2 are certain functions depending on time. The precise form of the coefficients of the Gambier equation is determined by singularity analysis, which leads to some constraints on a_0, a_1 and a_2 [30]. For simplicity, we hereafter assume $a_0(0) \neq 0$. Nevertheless, all our results can easily be generalised for the case $a_0(0) = 0$.

If $n \neq 0$, we can eliminate y between the two equations above, which gives rise to the referred to as *second-order Gambier equation* [32, 39, 40, 44], i.e.

$$\begin{aligned} \frac{d^2x}{dt^2} &= \frac{n-1}{xn} \left(\frac{dx}{dt} \right)^2 + a_0 \frac{(n+2)}{n} x \frac{dx}{dt} + a_1 \frac{dx}{dt} - \sigma \frac{(n-2)}{nx} \frac{dx}{dt} \\ &\quad - \frac{a_0^2}{n} x^3 + \left(\frac{da_0}{dt} - a_0 a_1 \right) x^2 + \left(a_2 n - 2a_0 \frac{\sigma}{n} \right) x - a_1 \sigma - \frac{\sigma^2}{nx}. \end{aligned} \quad (3.1)$$

The importance of second-order Gambier equations is due to their relations to remarkable differential equations such as second-order Riccati equations [31, 35], second-order Kummer–Schwarz equations [5, 16] and Milne–Pinney equations [32]. Additionally, by making appropriate limits in their coefficients, Gambier equations describe all the linearisable equations of the Painlevé–Gambier list [32]. Several particular cases of these equations have also been studied in order to analyse discrete systems [40].

Particular instances of (3.1) have already been investigated through the theory of Lie systems and quasi-Lie schemes. For instance, by fixing $n = -2$, $\sigma = a_1 = 0$, and a_0 to be a constant, the second-order Gambier equation (3.1) becomes a second-order Kummer–Schwarz equation (KS2 equation) [5, 16]

$$\frac{d^2x}{dt^2} = \frac{3}{2x} \left(\frac{dx}{dt} \right)^2 - 2c_0x^3 + 2\omega(t)x, \quad (3.2)$$

where we have written $c_0 = -a_0^2/4$, with c_0 a non-positive constant, and $\omega(t) = -a_2(t)$ so as to keep, for simplicity in following procedures, the same notion as used in the literature, e.g. in [16]. The interest of KS2 equations is due to their relations to other differential equations of physical and mathematical interest [16, 24, 32]. For instance, for $x > 0$ the change of variables $y = 1/\sqrt{x}$ transforms KS2 equations into Milne–Pinney equations, which frequently occur in cosmology [32]. Meanwhile, the non-local transformation $dy/dt = x$ maps KS2 equations into a particular type of third-order Kummer–Schwarz equations, which are closely related to Schwarzian derivatives [4, 16, 43]. Additionally, KS2 equations can be related, through the addition of the new variable $dx/dt = v$, to a Lie system associated to a Vessiot–Guldberg Lie algebra isomorphic to $\mathfrak{sl}(2, \mathbb{R})$, which gave rise to various methods to study its properties and related problems [16].

If we now assume $n = 1$ and $\sigma = 0$ in (3.1), it results

$$\frac{d^2x}{dt^2} = (a_1 + 3a_0x) \frac{dx}{dt} - a_0^2x^3 + \left(\frac{da_0}{dt} - a_0a_1 \right) x^2 + a_2x,$$

which is a particular case of second-order Riccati equations [23, 31] that has been treated through the theory of quasi-Lie schemes and Lie systems in several works [10, 13, 29]. Furthermore, equations of this type have been broadly investigated because of its appearance in the study of the Bäcklund transformations for PDEs, their relation to physical problems, and the interest of the algebraic structure of their Lie symmetries [8, 23, 31, 37, 42].

In view of the previous results, it is natural to wonder which kind of second-order Gambier equations can be studied through the theory of quasi-Lie schemes. To this end, let us build up a quasi-Lie scheme for analysing these equations.

As usual, the introduction of the new variable $v \equiv dx/dt$ enables us to relate the second-order Gambier equation (3.1) to the first-order system

$$\begin{aligned} \frac{dx}{dt} &= v, \\ \frac{dv}{dt} &= \frac{(n-1)v^2}{n} \frac{1}{x} + a_0 \frac{(n+2)}{n} xv + a_1v - \sigma \frac{(n-2)v}{n} \frac{1}{x} - \frac{a_0^2}{n} x^3 \\ &\quad + \left(\frac{da_0}{dt} - a_0a_1 \right) x^2 + \left(a_2n - 2a_0 \frac{\sigma}{n} \right) x - a_1\sigma - \frac{\sigma^2}{nx}, \end{aligned}$$

which is associated to the t -dependent vector field on $\mathbb{T}\mathbb{R}_0$, with $\mathbb{R}_0 \equiv \mathbb{R} \setminus \{0\}$, given by

$$X = v \frac{\partial}{\partial x} + \left[\frac{(n-1)v^2}{n} \frac{1}{x} + a_0 \frac{(n+2)}{n} xv + a_1v - \sigma \frac{(n-2)v}{n} \frac{1}{x} - \frac{a_0^2}{n} x^3 \right.$$

$$+ \left(\frac{da_0}{dt} - a_0 a_1 \right) x^2 + \left(a_2 n - 2a_0 \frac{\sigma}{n} \right) x - a_1 \sigma - \frac{\sigma^2}{nx} \Big] \frac{\partial}{\partial v},$$

termed henceforth the *Gambier vector field*. To obtain a quasi-Lie scheme for studying the above equations, we need to find a finite-dimensional \mathbb{R} -linear space V_G such that $X \in V_G(C^\infty(\mathbb{R}))$ for all a_0, a_1, a_2, σ and n . Observe that X can be cast in the form

$$X = \sum_{\alpha=1}^{10} b_\alpha \left(a_0, \frac{da_0}{dt}, a_1, a_2, \sigma, n \right) Y_\alpha, \quad (3.3)$$

with b_1, \dots, b_{10} being certain t -dependent functions whose form depends on the functions $a_0, da_0/dt, a_1, a_2$ and the constants σ and n . More specifically, these functions read

$$\begin{aligned} b_1 &= 1, & b_2 &= \frac{n-1}{n}, & b_3 &= a_0 \frac{n+2}{n}, & b_4 &= a_1, & b_5 &= -\sigma \frac{n-2}{n}, \\ b_6 &= -\frac{a_0^2}{n}, & b_7 &= \frac{da_0}{dt} - a_0 a_1, & b_8 &= a_2 n - 2a_0 \frac{\sigma}{n}, & b_9 &= -a_1 \sigma, & b_{10} &= -\frac{\sigma^2}{n} \end{aligned}$$

and

$$\begin{aligned} Y_1 &= v \frac{\partial}{\partial x}, & Y_2 &= \frac{v^2}{x} \frac{\partial}{\partial v}, & Y_3 &= xv \frac{\partial}{\partial v}, & Y_4 &= v \frac{\partial}{\partial v}, & Y_5 &= \frac{v}{x} \frac{\partial}{\partial v}, \\ Y_6 &= x^3 \frac{\partial}{\partial v}, & Y_7 &= x^2 \frac{\partial}{\partial v}, & Y_8 &= x \frac{\partial}{\partial v}, & Y_9 &= \frac{\partial}{\partial v}, & Y_{10} &= \frac{1}{x} \frac{\partial}{\partial v}. \end{aligned}$$

For convenience, we further define the vector field

$$Y_{11} = x \frac{\partial}{\partial x},$$

which, although does not appear in the decomposition (3.3), will shortly become useful so as to describe the properties of Gambier vector fields.

In view of (3.3), it easily follows that we can choose V_G to be the space spanned by Y_1, \dots, Y_{11} . It is interesting to note that the linear space V_G is not a Lie algebra as $[Y_3, Y_6]$ does not belong to V_G . Moreover, as

$$\text{ad}_{Y_3}^j Y_6 \equiv \overbrace{[Y_3, [Y_3, [\dots, [Y_3, Y_6] \dots]]]}^{j\text{-times}} = (-1)^j x^{j+3} \frac{\partial}{\partial v}, \quad j \in \mathbb{N},$$

there is no finite-dimensional real Lie algebra $\widehat{V} \supset V_G$ such that $X \in \widehat{V}(C^\infty(\mathbb{R}))$. Hence, X is not in general a Lie system, which suggests us to use quasi-Lie schemes to investigate it.

To determine a quasi-Lie scheme involving V_G , we must find a real finite-dimensional Lie algebra $W_G \subset V_G$ such that $[W_G, V_G] \subset V_G$. In view of Table 1, we can do so by setting $W_G = \langle Y_4, Y_8, Y_{11} \rangle$, which is a solvable three-dimensional Lie algebra. In fact,

$$[Y_4, Y_8] = -Y_8, \quad [Y_4, Y_{11}] = 0, \quad [Y_8, Y_{11}] = -Y_8.$$

In other words, we have proved the following proposition providing a new quasi-Lie scheme to study Gambier vector fields and, as shown posteriorly, second-order Gambier equations.

Proposition 3.1. *The spaces $V_G = \langle Y_1, \dots, Y_{11} \rangle$ and $W_G = \langle Y_4, Y_8, Y_{11} \rangle$ form a quasi-Lie scheme $S(W_G, V_G)$ such that $X \in V_G(C^\infty(\mathbb{R}))$ for every Gambier vector field X .*

Recall that Y_{11} is not necessary so that every Gambier vector field takes values in V_G . Hence, why it is convenient to add it to V_G ? One reason can be found in Table 1. If V_G had not contained Y_{11} , then $S(W_G, V_G)$ would have not been a quasi-Lie scheme as $W_G \not\subset V_G$. In addition, Y_8 could not belong to W_G neither, as $[Y_8, Y_1] \in V_G$ provided $Y_{11} - Y_4 \in V_G$. Hence, including Y_{11} in V_G allows us to choose a larger $W_G \subset V_G$. In turn this gives rise to a larger group $\mathcal{G}(W_G)$, which will be of great use in following sections.

Table 1. Lie brackets $[Y_i, Y_j]$ with $i = 4, 8, 11$ and $j = 1, \dots, 11$.

$[\cdot, \cdot]$	Y_1	Y_2	Y_3	Y_4	Y_5	Y_6	Y_7	Y_8	Y_9	Y_{10}	Y_{11}
Y_4	Y_1	Y_2	0	0	0	$-Y_6$	$-Y_7$	$-Y_8$	$-Y_9$	$-Y_{10}$	0
Y_8	$Y_{11} - Y_4$	$2Y_4$	Y_7	Y_8	Y_9	0	0	0	0	0	$-Y_8$
Y_{11}	$-Y_1$	$-Y_2$	Y_3	0	$-Y_5$	$3Y_6$	$2Y_7$	Y_8	0	$-Y_{10}$	0

4 Transformation properties of second-order Gambier equations

Remind that Theorem 2.1 states that the t -dependent changes of variables associated to the elements of the group $\mathcal{G}(W)$ of a quasi-Lie scheme $S(W, V)$ establish bijections among the t -dependent vector fields taking values in V . This may be of great use so as to transform their associated systems into simplified forms, e.g. in the case of Abel equations [11]. We next show how this can be done for studying the transformation of second-order Gambier equations into simpler ones whose corresponding coefficients a_1 vanish [30]. This retrieves known results from a geometrical viewpoint and shows that certain second-order Gambier equations cannot be transformed into simpler ones, solving a small gap performed in [30].

As the vector fields in W_G span a finite-dimensional real Lie algebra of vector fields on $\mathbb{T}\mathbb{R}_0$, there exists a local Lie group action $\varphi : G \times \mathbb{T}\mathbb{R}_0 \rightarrow \mathbb{T}\mathbb{R}_0$ whose fundamental vector fields are the elements of W_G . By integrating the vector fields of W_G (see [12] for details), the action can easily be written as

$$\varphi \left(g, \begin{pmatrix} x \\ v \end{pmatrix} \right) = \begin{pmatrix} \alpha x \\ \gamma x + \delta v \end{pmatrix}, \quad \text{where} \quad g \in T_d \equiv \left\{ \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \mid \alpha, \delta \in \mathbb{R}_+, \gamma \in \mathbb{R} \right\}.$$

The theory of Lie systems [9, 17] states that the solutions of a system associated to a t -dependent vector field taking values in the real Lie algebra W_G are of the form $(x(t), v(t)) = \varphi(h(t), (x_0, v_0))$, with $h(t)$ being a curve in T_d with $h(0) = e$. Therefore, every $g \in \mathcal{G}(W_G)$ can be written as $g_t(\cdot) = \varphi(h(t), \cdot)$ for a certain curve $h(t)$ in G with $h(0) = e$. Conversely, given a curve $h(t)$ in T_d with $h(0) = e$, the curve $(x(t), v(t)) = \varphi(h(t), (x_0, v_0))$ is the general solution of a system of $W_G(C^\infty(\mathbb{R}))$, which leads to a generalised flow $g_t(\cdot) = \varphi(h(t), \cdot)$ of $\mathcal{G}(W_G)$ [9, 17]. Hence, the elements of $\mathcal{G}(W_G)$ are generalised flows of the form

$$g^{h(t)}(t, x, v) = \varphi(h(t), x, v),$$

for $h(t)$ any curve in T_d with $h(0) = e$. Observe that every $h(t)$ is a matrix of the form

$$h(t) = \begin{pmatrix} \alpha(t) & 0 \\ \gamma(t) & \delta(t) \end{pmatrix}$$

where $\alpha(t)$, $\delta(t)$ and $\gamma(t)$ are t -dependent functions such that $\alpha(0) = \delta(0) = 1$ and $\gamma(0) = 0$, because $h(0) = e$, and $\alpha(t) > 0$, $\delta(t) > 0$ as $h(t) \in T_d$ for every $t \in \mathbb{R}$. Hence, every element of $\mathcal{G}(W_G)$ is of the form

$$g^{\alpha(t), \gamma(t), \delta(t)}(t, x, v) \equiv g^{h(t)}(t, x, v) = (t, \alpha(t)x, \gamma(t)x + \delta(t)v). \quad (4.1)$$

Theorem 2.1 implies that for every $g \in \mathcal{G}(W_G)$ and Gambier vector field $X \in V_G(C^\infty(\mathbb{R}))$, we have $g_\star X \in V_G(C^\infty(\mathbb{R}))$. More specifically, a long but straightforward calculation shows that

$$g_\star X = \sum_{\alpha=1}^{11} \bar{b}_\alpha(t) Y_\alpha, \quad (4.2)$$

where the functions $\bar{b}_\alpha = \bar{b}_\alpha(t)$, with $\alpha = 1, \dots, 11$, are

$$\begin{aligned}\bar{b}_1 &= \frac{\alpha}{\delta}, & \bar{b}_2 &= \frac{n-1}{n} \frac{\alpha}{\delta}, & \bar{b}_3 &= \frac{a_0(n+2)}{n\alpha}, & \bar{b}_4 &= a_1 + \frac{1}{\delta} \frac{d\delta}{dt} + \frac{(2-n)\gamma}{n\delta}, \\ \bar{b}_5 &= \frac{(2-n)\sigma\alpha}{n}, & \bar{b}_6 &= -\frac{a_0^2\delta}{n\alpha^3}, & \bar{b}_7 &= \frac{1}{\alpha^2} \left(\delta \frac{da_0}{dt} - a_0 a_1 \delta - \frac{n+2}{n} a_0 \gamma \right), \\ \bar{b}_8 &= \frac{\delta}{\alpha} \left(na_2 - \frac{2\sigma}{n} a_0 - \frac{\gamma}{\delta} a_1 - \frac{\gamma^2}{n\delta^2} - \frac{\gamma}{\delta^2} \frac{d\delta}{dt} + \frac{1}{\delta} \frac{d\gamma}{dt} \right), \\ \bar{b}_9 &= -\sigma \left(a_1 \delta + \frac{(2-n)\gamma}{n} \right), & \bar{b}_{10} &= -\frac{\sigma^2 \alpha \delta}{n}, & \bar{b}_{11} &= \frac{1}{\alpha} \frac{d\alpha}{dt} - \frac{\gamma}{\delta}.\end{aligned}\tag{4.3}$$

Let us use this so as to transform the initial Gambier vector field into another one that is related, up to a t -reparametrisation $\tau = \tau(t)$, to a second-order Gambier equation with τ -dependent coefficients $\bar{a}_0, \bar{a}_1, \bar{a}_2$, a constant $\bar{\sigma}$ and an integer number \bar{n} . Additionally, we impose $\bar{a}_1 = 0$ in order to reduce our initial first-order Gambier equation into a simpler one. In this way, we have

$$\begin{aligned}g_\star X &= \xi(t) \left[Y_1 + \frac{\bar{n}-1}{\bar{n}} Y_2 + \bar{a}_0 \frac{\bar{n}+2}{\bar{n}} Y_3 - \sigma \frac{\bar{n}-2}{\bar{n}} Y_5 \right. \\ &\quad \left. - \frac{\bar{a}_0^2}{\bar{n}} Y_6 + \frac{d\bar{a}_0}{d\tau} Y_7 + \left(\bar{a}_2 \bar{n} - 2\bar{a}_0 \frac{\bar{\sigma}}{\bar{n}} \right) Y_8 - \frac{\bar{\sigma}^2}{\bar{n}} Y_{10} \right],\end{aligned}\tag{4.4}$$

for a certain non-vanishing function $\xi(t) = d\tau/dt$. Therefore, $\bar{b}_4 = \bar{b}_9 = \bar{b}_{11} = 0$, i.e.

$$a_1 + \frac{1}{\delta} \frac{d\delta}{dt} + \frac{(2-n)\gamma}{n\delta} = 0,\tag{4.5}$$

$$\sigma \left(a_1 \delta + \frac{(2-n)\gamma}{n} \right) = 0,\tag{4.6}$$

$$\frac{1}{\alpha} \frac{d\alpha}{dt} - \frac{\gamma}{\delta} = 0.\tag{4.7}$$

As we want our method to work for all values of σ , e.g. $\sigma \neq 0$, equation (4.6) implies

$$a_1 \delta + \frac{(2-n)\gamma}{n} = 0.\tag{4.8}$$

As $\delta > 0$, the above equation involves that $a_1 = 0$ for $n = 2$. In other words, we cannot transform a Gambier vector field with $a_1 \neq 0$ into a new one with $\bar{a}_1 = 0$ through our methods if $n = 2$ and $\sigma \neq 0$. In view of this, let us assume that $n \neq 2$.

From (4.5) and (4.8), and using again that $\delta > 0$, we infer that $d\delta/dt = 0$. As $\delta(0) = 1$, then $\delta = 1$. Plugging the value of δ into (4.7) and (4.8), we obtain

$$\frac{1}{\alpha} \frac{d\alpha}{dt} = \frac{na_1}{n-2} = \gamma \iff \alpha = \exp \left(\int_0^t \frac{na_1}{n-2} dt' \right), \quad \gamma = \frac{na_1}{n-2},$$

which fixes the form of g mapping a system X into (4.4). Bearing previous results in mind, we see that the non-vanishing t -dependent coefficients (4.3) become

$$\begin{aligned}\bar{b}_1 &= \alpha, & \bar{b}_2 &= \frac{n-1}{n} \alpha, & \bar{b}_3 &= \frac{a_0(n+2)}{n\alpha}, & \bar{b}_5 &= \frac{(2-n)\sigma\alpha}{n}, & \bar{b}_6 &= -\frac{a_0^2}{n\alpha^3}, \\ \bar{b}_7 &= \frac{d}{dt} \left(\frac{a_0}{\alpha^2} \right), & \bar{b}_8 &= \frac{1}{\alpha} \left(na_2 - \frac{2\sigma a_0}{n} - \frac{n(n-1)}{(n-2)^2} a_1^2 + \frac{n}{n-2} \frac{da_1}{dt} \right), & \bar{b}_{10} &= -\frac{\sigma^2 \alpha}{n}.\end{aligned}$$

Comparing this and (4.4), we see that to transform the initial first-order Gambier equation into a new one through a t -dependent change of variables (4.1) and a t -reparametrisation $d\tau = \alpha dt$ requires $\xi = \alpha$. The resulting system reads

$$\begin{aligned} \frac{d\bar{x}}{d\tau} &= \bar{v}, \\ \frac{d\bar{v}}{d\tau} &= \frac{(n-1)\bar{v}^2}{n\bar{x}} + a_0 \frac{(n+2)}{\alpha^2 n} \bar{x}\bar{v} - \sigma \frac{(n-2)\bar{v}}{n\bar{x}} - \frac{a_0^2}{\alpha^4 n} \bar{x}^3 \\ &\quad + \frac{d(a_0/\alpha^2)}{d\tau} \bar{x}^2 + \left(\bar{a}_2 n - 2a_0 \alpha^{-2} \frac{\sigma}{n} \right) \bar{x} - \frac{\sigma^2}{n\bar{x}}, \end{aligned}$$

where

$$\bar{a}_2 = \frac{1}{\alpha^2} \left(a_2 - \frac{(n-1)}{(n-2)^2} a_1^2 + \frac{1}{n-2} \frac{da_1}{dt} \right).$$

Therefore, redefining $\bar{n} = n$, $\bar{a}_0 = a_0/\alpha^2$ and $\bar{\sigma} = \sigma$, we obtain that the above system is associated to a second-order Gambier equation with $\bar{a}_1 = 0$. Meanwhile, as g induces a t -dependent change of variables given by

$$\bar{x} = \alpha x, \quad \bar{v} = \frac{d\alpha}{d\tau} x + v,$$

we see that this t -dependent change of variables can be viewed as a t -dependent change of variables $\bar{x} = \alpha x$ along with a t -reparametrisation $t \rightarrow \tau$ such that $\bar{v} = d\bar{x}/d\tau$. Indeed,

$$\bar{x} = \alpha x \quad \implies \quad \frac{d\bar{x}}{d\tau} = \frac{d\alpha}{d\tau} x + \alpha \frac{dx}{d\tau} = \frac{d\alpha}{d\tau} x + v = \bar{v}.$$

Furthermore, these transformations map the initial second-order Gambier equation with $n \neq 2$ into a new one with $\bar{a}_1 = 0$, i.e. depending only on two functions \bar{a}_0 and \bar{a}_2 and the constants σ and n . We can therefore formulate the following result.

Proposition 4.1. *Every second-order Gambier equation (3.1) with $n \neq 2$ can be transformed via a t -dependent change of variables $\bar{x} = \alpha(t)x$ and a t -reparametrisation $\tau = \tau(t)$, with $d\tau = \alpha dt$, $\alpha(0) = 1$ and*

$$\frac{1}{\alpha} \frac{d\alpha}{dt} = \frac{n}{n-2} a_1,$$

into a second-order Gambier equation whose a_1 vanishes and n, σ remain the same.

In the above proposition, we excluded second-order Gambier equations with $n = 2$ as we noticed that in this case the proof of this proposition does not hold: we cannot transform an initial Gambier vector field with $\sigma \neq 0$ and $a_1 \neq 0$ into a new one with $\bar{a}_1 = 0$. Moreover, it is easy to see that the transformation provided in Proposition 4.1 does not exist for $n = 2$ and $a_1 \neq 0$. This was not noticed in [30], where this transformation is wrongly claimed to transform any Gambier equation into a simpler one with $a_1 = 0$. In view of this, we cannot neither ensure, as claimed in [30], that second-order Gambier equations are not given in their simplest form.

5 Quasi-Lie systems and Gambier equations

The theory of quasi-Lie schemes mainly provides information about quasi-Lie systems, which can be mapped to Lie systems through one of the transformations of the group of a quasi-Lie scheme. This allows us to employ the techniques of the theory of Lie systems so as to study the

obtained Lie systems, and, undoing the performed t -dependent change of variables, to describe properties of the initial system [15].

Motivated by the above, we determine and study the Gambier vector fields $X \in V(C^\infty(\mathbb{R}))$ which are quasi-Lie systems relative to $S(W_G, V_G)$. This task relies in finding triples (g, X, V_0) , with $g \in \mathcal{G}(W_G)$ and V_0 being a real Lie algebra included in V_G in such a way that $g_\star X \in V_0(C^\infty(\mathbb{R}))$.

One of the key points to determine quasi-Lie systems is to find a Lie algebra V_0 . In the case of Gambier vector fields, this can readily be obtained by recalling that certain instances of second-order Gambier equations, e.g. (3.2), are particular cases of KS2 equations. By adding a new variable $v \equiv dx/dt$ to (3.2), the resulting first-order system becomes a Lie system (see [16]) related to a three-dimensional Vessiot–Guldberg Lie algebra $V_0 \subset V_G$ of vector fields on TR_0 spanned by

$$X_1 = 2x \frac{\partial}{\partial v}, \quad X_2 = x \frac{\partial}{\partial x} + 2v \frac{\partial}{\partial v}, \quad X_3 = v \frac{\partial}{\partial x} + \left(\frac{3}{2} \frac{v^2}{x} - 2c_0 x^3 \right) \frac{\partial}{\partial v}, \quad (5.1)$$

i.e.

$$X_1 = 2Y_8, \quad X_2 = Y_{11} + 2Y_4, \quad X_3 = Y_1 + \frac{3}{2}Y_2 - 2c_0 Y_6.$$

Consequently, it makes sense to look for Gambier vector fields $X \in V(C^\infty(\mathbb{R}))$ that can be transformed, via an element $g \in \mathcal{G}(W_G)$, into a Lie system $g_\star X \in V_0(C^\infty(\mathbb{R}))$, i.e.

$$g_\star X = 2f(t)Y_8 + g(t)(Y_{11} + 2Y_4) + h(t) \left(Y_1 + \frac{3}{2}Y_2 - 2c_0 Y_6 \right),$$

for certain t -dependent functions f , g and h . Comparing the expression of $g_\star X$ given by (4.3) and the above, we find that $g_\star X \in V_0(C^\infty(\mathbb{R}))$ if and only if

$$\bar{b}_3 = \bar{b}_5 = \bar{b}_7 = \bar{b}_9 = \bar{b}_{10} = 0 \quad (5.2)$$

and

$$\bar{b}_4 = 2\bar{b}_{11}, \quad \bar{b}_2 = \frac{3}{2}\bar{b}_1, \quad \bar{b}_6 = -2c_0\bar{b}_1. \quad (5.3)$$

From expressions (4.3) and remembering that $\alpha > 0$ and $\delta > 0$, we see that condition $\bar{b}_{10} = 0$ implies that $\sigma = 0$. This involves, along with (4.3), that $\bar{b}_5 = \bar{b}_9 = 0$. Meanwhile, from $\bar{b}_2 = 3\bar{b}_1/2$ we obtain $n = -2$, which in turn ensures that $\bar{b}_3 = 0$. Bearing all this in mind, $\bar{b}_7 = 0$ reads

$$a_0 a_1 - \frac{da_0}{dt} = 0. \quad (5.4)$$

Above results impose restrictions on the form of the Gambier vector field X to be able to be transformed into a Lie system possessing a Vessiot–Guldberg Lie algebra V_0 via an element $g \in \mathcal{G}(W_G)$. Let us show that the remaining conditions in (5.2) and (5.3) merely characterise the form of the t -dependent change of variables g and the coefficient c_0 appearing in (3.2).

Conditions $2\bar{b}_{11} = \bar{b}_4$ and $\bar{b}_6 = -2c_0\bar{b}_1$ read

$$\frac{d}{dt} \log \frac{\alpha^2}{\delta} = a_1, \quad a_0^2 = -4c_0 \frac{\alpha^4}{\delta^2}. \quad (5.5)$$

Using the first condition above, the relation (5.4), and taking into account that $\alpha(0) = \delta(0) = 1$ and $\delta, \alpha > 0$, we see that

$$\frac{\frac{d}{dt}(\alpha^2 \delta^{-1})}{\alpha^2 \delta^{-1}} = \frac{1}{a_0} \frac{da_0}{dt} \implies \frac{\alpha^2}{\delta} = \frac{a_0}{a_0(0)}.$$

Using the second equality in (5.5), we obtain

$$4c_0 = -a_0(0)^2,$$

which fixes c_0 in terms of a coefficient of the initial second-order Gambier equation.

Concerning the t -dependent coefficients $\bar{b}_1, \dots, \bar{b}_{11}$, the non-vanishing ones under above conditions, i.e. $\bar{b}_1, \bar{b}_2, \bar{b}_4, \bar{b}_6, \bar{b}_8$ and \bar{b}_{11} , can readily be obtained through relations (5.3) and

$$\bar{b}_1 = \frac{\alpha}{\delta}, \quad \bar{b}_8 = \frac{\delta}{\alpha} \left(-2a_2 - \frac{\gamma}{\delta a_0} \frac{da_0}{dt} + \frac{\gamma^2}{2\delta^2} - \frac{\gamma}{\delta^2} \frac{d\delta}{dt} + \frac{1}{\delta} \frac{d\gamma}{dt} \right), \quad \bar{b}_{11} = \frac{1}{\alpha} \frac{d\alpha}{dt} - \frac{\gamma}{\delta}.$$

In other words, we have proved the following proposition.

Proposition 5.1. *A Gambier vector field X is a quasi-Lie system relative to $S(W_G, V_G)$ that can be transformed into a Lie system taking values in a Lie algebra of the form V_0 if and only if $a_0 a_1 = da_0/dt$, $n = -2$ and $\sigma = 0$. Under these conditions, the constant c_0 appearing in X_3 becomes $c_0 = -a_0(0)^2/4$. Additionally, a transformation $g(\alpha(t), \gamma(t), \delta(t)) \in \mathcal{G}(W_G)$ maps X into $g_\star X \in V_0(C^\infty(\mathbb{R}))$ if and only if $\alpha^2/\delta = a_0/a_0(0)$. More specifically, $g_\star X$ reads*

$$g_\star X = \frac{\alpha}{\delta} X_3 + \left(\frac{1}{\alpha} \frac{d\alpha}{dt} - \frac{\gamma}{\delta} \right) X_2 + \frac{\delta}{2\alpha} \left(-2a_2 - \frac{\gamma}{\delta a_0} \frac{da_0}{dt} + \frac{\gamma^2}{2\delta^2} - \frac{\gamma}{\delta^2} \frac{d\delta}{dt} + \frac{1}{\delta} \frac{d\gamma}{dt} \right) X_1. \quad (5.6)$$

The above proposition allows us to determine the transformations $g \in \mathcal{G}(W_G)$ ensuring that a Gambier vector field and its related second-order Gambier equation can be mapped, maybe up to a t -reparametrisation, into a new Gambier vector field related to a KS2 equation. Indeed, from (3.2), we see that to do so, we need to impose

$$g_\star X = \xi(t)(X_3 + \omega(t)X_1),$$

for a certain function $\omega(t)$ and a function $\xi(t)$ such that $\xi(t) \neq 0$ for all $t \in \mathbb{R}$. Comparing (5.6) with the above expression, we see that

$$\frac{1}{\alpha} \frac{d\alpha}{dt} - \frac{\gamma}{\delta} = 0, \quad (5.7)$$

which, in view of the fact that $g^{\alpha(t), \gamma(t), \delta(t)}$ satisfies

$$\frac{\alpha^2}{\delta} = \frac{a_0}{a_0(0)}, \quad (5.8)$$

enables us to determine the searched transformations. In fact, fixed a non-vanishing t -dependent function α with $\alpha(0) = 1$, the above conditions determine the values of δ and γ of g .

Now, a t -dependent reparametrisation

$$\tau = \int_0^t \frac{\alpha}{\delta} dt' \quad (5.9)$$

transforms the system associated to $g_\star X$ into a new system related to $X_3 + \omega(t(\tau))X_1$, where ω is given by

$$\omega = -\frac{\delta^2}{2\alpha^2} \left(2a_2 + \frac{\gamma}{a_0\delta} \frac{da_0}{dt} - \frac{\gamma^2}{2\delta^2} - \frac{1}{\delta} \frac{d\gamma}{dt} + \frac{\gamma}{\delta^2} \frac{d\delta}{dt} \right). \quad (5.10)$$

Proposition 5.2. *Every Gambier vector field X satisfying $a_0 a_1 = da_0/dt$, $n = -2$ and $\sigma = 0$ is a quasi-Lie system relative to $S(W_G, V_G)$ that can be transformed into a Lie system associated to $X_3 + \omega(t)X_1$, with $c_0 = -a_0(0)^2/4$ and $\omega(t)$ given by (5.10), through a transformation $g^{\alpha(t), \gamma(t), \delta(t)} \in \mathcal{G}(W_G)$, whose coefficients are given by any solution of (5.7) and (5.8), and the t -dependent reparametrisation (5.9).*

Note that the transformation $g^{\alpha(t),\gamma(t),\delta(t)}$ can be viewed as a t -dependent change of variables $\bar{x} = \alpha x$ and a t -reparametrisation $d\tau = \alpha/\delta dt$, i.e.

$$\bar{x} = \alpha x, \quad \bar{v} = \gamma x + \delta v,$$

and, in view of the first condition in (5.7), we see that

$$\frac{d\bar{x}}{d\tau} = \frac{d\alpha}{d\tau}x + \alpha \frac{dx}{d\tau} = \gamma x + \delta v = \bar{v}.$$

From this and Proposition 5.2, we obtain the following proposition about second-order Gambier equations.

Corollary 5.1. *Every second-order Gambier equation of the form*

$$\frac{d^2x}{dt^2} = \frac{3}{2x} \left(\frac{dx}{dt} \right)^2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{dx}{dt} + \frac{a_0^2}{2} x^3 - 2a_2x \quad (5.11)$$

can be mapped into a KS2 equation

$$\frac{d^2\bar{x}}{d\tau^2} = \frac{3}{2\bar{x}} \left(\frac{d\bar{x}}{d\tau} \right)^2 + \frac{1}{2} a_0(0)^2 \bar{x}^3 - \frac{\delta^2}{\alpha^2} \left(2a_2 + \frac{\gamma}{a_0\delta} \frac{da_0}{dt} - \frac{1}{2} \frac{\gamma^2}{\delta^2} - \frac{1}{\delta} \frac{d\gamma}{dt} + \frac{\gamma}{\delta^2} \frac{d\delta}{dt} \right) \bar{x} \quad (5.12)$$

through a t -dependent change of variables $\bar{x}(t) = \alpha(t)x(t)$, where $\alpha(t)$ is any positive function with $\alpha(0) = 1$, and a t -reparametrisation $\tau(t)$ with $d\tau = \alpha/\delta dt$, with δ and γ being determined from α by the relations

$$\delta = \alpha^2 \frac{a_0(0)}{a_0}, \quad \gamma = \frac{\alpha a_0(0)}{a_0} \frac{d\alpha}{dt}. \quad (5.13)$$

6 Constants of motion for second-order Gambier equations

In this section, we obtain constants of motion for second-order Gambier equations. We do so by analysing the existence of t -independent constants of motion for systems $g_{\star}X \in V_0(C^\infty(\mathbb{R}))$, with X being a Gambier vector field. By undoing the t -dependent change of variables g , this leads to determining constants of motion for a Gambier vector field X and its corresponding second-order Gambier equation.

Let $F : \mathbb{T}\mathbb{R}_0 \rightarrow \mathbb{R}$ be a t -independent constant of motion for $g_{\star}X$, we have $(g_{\star}X)_t F = 0$ for all $t \in \mathbb{R}$. This involves that F is a t -independent constant of motion for all successive Lie brackets of elements of $\{(g_{\star}X)_t\}_{t \in \mathbb{R}}$ as well as their linear combinations. In other words, F is a common first-integral for the vector fields $V^{g_{\star}X} \subset V_0$.

When $\omega(t)$ is not a constant, it can be verified that $V^{g_{\star}X} = V_0$. Thus, if F is a first-integral for all these vector fields, it is so for all vector fields contained in the generalised distribution $\mathcal{D}_p = \langle (X_1)_p, (X_2)_p, (X_3)_p \rangle$, with $p \in \mathbb{T}\mathbb{R}_0$. Hence, $dF_p \in \mathcal{D}_p^\circ$, i.e. dF_p is incident to all vectors of \mathcal{D}_p . In this case, $\mathcal{D}_p = \mathbb{T}_p\mathbb{T}\mathbb{R}_0$ for a generic $p \in \mathbb{T}\mathbb{R}_0$, which implies that $dF_p = 0$ at almost every point. Since F is assumed to be differentiable, we have that F is constant on each connected component of $\mathbb{T}\mathbb{R}_0$ and $g_{\star}X$ has no non-trivial t -independent constant of motion.

If $\omega(t) = \lambda$ for a real constant λ , then $\dim \mathcal{D}_p = 1$ at a generic point and it makes sense to look for non-trivial t -dependent constants of motion for $g_{\star}X$. In view of (5.10), this condition implies

$$\lambda = -\frac{a_0(0)^2 \alpha^2}{2a_0^2} \left[2a_2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{d \log \alpha}{dt} - \frac{1}{2} \left(\frac{d \log \alpha}{dt} \right)^2 - \frac{d^2 \log \alpha}{dt^2} \right].$$

The function F can therefore be obtained by integrating

$$v \frac{\partial F}{\partial x} + \left(\frac{3}{2} \frac{v^2}{x} + \frac{a_0(0)^2}{2} x^3 + 2\lambda x \right) \frac{\partial F}{\partial v} = 0.$$

In employing the *characteristics method* [10], we find that F must be constant along the integral curves of the so-called *characteristic system*, namely

$$\frac{dx}{v} = \frac{dv}{\frac{3}{2} \frac{v^2}{x} + \frac{a_0(0)^2}{2} x^3 + 2\lambda x}.$$

Let us focus on the region with $v \neq 0$, i.e. $\mathcal{O} \equiv \{(x, v) \in \mathbb{T}\mathbb{R}_0 \mid v \neq 0\}$. We obtain from the previous equations that

$$\frac{dv}{dx} = \frac{3}{2} \frac{v}{x} + \frac{a_0(0)^2}{2} \frac{x^3}{v} + 2\lambda \frac{x}{v}.$$

Let us focus on the case $x > 0$; the other case can be obtained in a similar way and leads to the same result. Multiplying on right and left by v/x and defining $w \equiv v^2$ and $z \equiv x^2$, we obtain

$$\frac{dw}{dz} = \frac{3}{2} \frac{w}{z} + \frac{a_0(0)^2}{2} z + 2\lambda.$$

As this equation is linear, its general solution can be easily derived to obtain

$$w(z) = (a_0(0)^2 z^{1/2} - 4\lambda z^{-1/2} + \xi) z^{3/2},$$

for an arbitrary real constant ξ . From here, it results

$$-a_0(0)^2 x + \frac{v^2}{x^3} + \frac{4\lambda}{x} = \xi.$$

Consequently, F is any function of the form $F = F(\xi)$, for instance,

$$F = -a_0(0)^2 x + \frac{v^2}{x^3} + \frac{4\lambda}{x}, \quad (x, v) \in \mathcal{O}.$$

In principle, F was defined only on \mathcal{O} . Nevertheless, as this region is an open and dense subset of $\mathbb{T}\mathbb{R}_0$, and in view of the expression for F , we can extend it differentiably to $\mathbb{T}\mathbb{R}_0$ in a unique way. Since F is a constant of motion on \mathcal{O} , it trivially becomes so on the whole $\mathbb{T}\mathbb{R}_0$.

Summarising, we have proved the following.

Theorem 6.1. *A second-order Gambier equation*

$$\frac{d^2 x}{dt^2} = \frac{3}{2x} \left(\frac{dx}{dt} \right)^2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{dx}{dt} + \frac{a_0^2}{2} x^3 - 2a_2 x, \quad (6.1)$$

admits a constant of motion of the form

$$F = -a_0(0)^2 \bar{x} + \frac{\bar{v}^2}{\bar{x}^3} + \frac{4\lambda}{\bar{x}}, \quad (6.2)$$

where λ is a real constant, $\bar{x} = \alpha x$ and $\bar{v} = \delta dx/dt + \gamma x$, with α being a particular positive solution with $\alpha(0) = 1$ of

$$\frac{d^2 \log \alpha}{dt^2} = \frac{2\lambda a_0^2}{a_0(0)^2 \alpha^2} + 2a_2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{d \log \alpha}{dt} - \frac{1}{2} \left(\frac{d \log \alpha}{dt} \right)^2 \quad (6.3)$$

and γ, δ are determined from α by means of the conditions (5.13).

The above proposition can be employed to derive a constant of motion for certain families of second-order Gambier equations. For instance, if we start by a Gambier equation (6.1) with $a_2 = -\lambda a_0^2/a_0(0)^2$, for a certain real constant λ , then $\alpha = 1$ is a solution of (6.3). In view of (5.13), $\gamma = 0$ and $\delta = a_0(0)/a_0$. Therefore, Theorem 6.1 establishes that the second-order Gambier equation (6.1) admits a constant of motion

$$F = -a_0(0)^2 x + \frac{a_0(0)^2}{a_0^2 x^3} \left(\frac{dx}{dt} \right)^2 + \frac{4\lambda}{x}. \quad (6.4)$$

Consider now a general second-order Gambier equation (6.1) and let us search for a constant of motion (6.2) with $\lambda = 0$. In this case, (6.3) can be brought into a Riccati equation

$$\frac{dw}{dt} = 2a_2 + \frac{1}{a_0} \frac{da_0}{dt} w - \frac{1}{2} w^2,$$

where $w \equiv d \log \alpha / dt$, whose solutions can be investigated through many methods [19, 22]. The derivation of a particular solution provides a constant of motion for the second-order Gambier equation (6.1) that can be obtained through the previous theorem. Additionally, this particular solution can be used to obtain the general solution of the Riccati equation [9, 19], which in turns could be used to derive new constants of motion for the second-order Gambier equation (6.1).

Note that all the above procedure depends deeply on the fact that λ is a constant. Recall that if λ is not a constant, then $g_{\star} X$ does not admit any t -independent constant of motion. Nevertheless, other methods can potentially be applied in this case. For instance, using that $S(V_0, V_0)$ is a quasi-Lie scheme, we can derive the group $\mathcal{G}(V_0)$ of this scheme and to use an element $h \in \mathcal{G}(V_0)$ to transform $g_{\star} X$ into other Lie system $h_{\star} g_{\star} X$ of the same type, e.g. another of the form $h_{\star} g_{\star} X = \xi_2(t)(c_1 X_1 + c_2 X_2 + c_3 X_3)$, with $c_1, c_2, c_3 \in \mathbb{R}$, the vector fields X_1, X_2, X_3 are those of (5.1), and $\xi_2(t)$ is any t -dependent nonvanishing function. As this system is, up to a t -parametrization, a t -independent vector field, it admits a local t -independent constant of motion. By inverting the t -dependent changes of variables h and g , it gives rise to a t -dependent constant of motion of X and the corresponding second-order Gambier equation.

7 Second-order Gambier and Milne–Pinney equations

Consider the KS2 equation (5.12) with $x > 0$ (we can proceed analogously for the case $x < 0$). The change of variables $x = 1/y^2$, with $y > 0$, transforms it into a Milne–Pinney equation [52]

$$\frac{d^2 y}{d\tau^2} = -\omega(t(\tau))y - \frac{a_0(0)^2}{4y^3}. \quad (7.1)$$

These equations admit a description in terms of a Lie system related to a Vessiot–Guldberg Lie algebra isomorphic to $\mathfrak{sl}(2, \mathbb{R})$ [12, 56]. This was employed in several works to describe their general solutions in terms of particular solutions of the same or others differential equations, e.g. Riccati equations and t -dependent harmonic oscillators [9].

Previous results allow us to describe the general solution of (5.11) in terms of particular solutions of Riccati equations or t -dependent frequency harmonic oscillators. Indeed, in view of Corollary 5.1, these equations can be transformed into a KS2 equation through a t -dependent change of variables $\bar{x}(t) = \alpha(t)x(t)$ and a t -reparametrisation $d\tau = (\alpha/\delta) dt$. In turn, $y = 1/\sqrt{\bar{x}}$ transforms (5.12) into (7.1), whose general solution $y(t)$ can be written as [12]

$$y(\tau) = \sqrt{k_1 z_1^2(\tau) + k_2 z_2^2(\tau) + 2C(k_1, k_2, W) z_1(\tau) z_2(\tau)},$$

where $C^2(k_1, k_2, W) = k_1 k_2 + a_0(0)^2/(4W^2)$, the functions $z_1(\tau)$, $z_2(\tau)$ are two linearly independent solutions of the system

$$\frac{d^2 z}{d\tau^2} = -\omega(t(\tau))z,$$

and W is the Wronskian related to such solutions. Inverting previous changes of variables, the general solution for any second-order Gambier equation (6.1) reads

$$x(t) = \alpha^{-1} \left[k_1 z_1^2(\tau(t)) + k_2 z_2^2(\tau(t)) \pm 2\sqrt{k_1 k_2 + a_0(0)^2/(4W^2)} z_1(\tau(t)) z_2(\tau(t)) \right]^{-1}.$$

Therefore, we have proved the following proposition:

Proposition 7.1. *The general solution of a second-order Gambier equation*

$$\frac{d^2 x}{dt^2} = \frac{3}{2x} \left(\frac{dx}{dt} \right)^2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{dx}{dt} + \frac{a_0^2}{2} x^3 - 2a_2 x, \quad (7.2)$$

can be brought into the form

$$x(t) = \alpha^{-1} \left[k_1 z_1^2(\tau(t)) + k_2 z_2^2(\tau(t)) \pm 2\sqrt{k_1 k_2 + a_0(0)^2/(4W^2)} z_1(\tau(t)) z_2(\tau(t)) \right]^{-1},$$

where $z_1(\tau)$ and $z_2(\tau)$ are particular solutions of the τ -dependent harmonic oscillator

$$\frac{d^2 z}{d\tau^2} = -\omega(t(\tau))z = -\frac{\delta^2}{2\alpha^2} \left(2a_2 + \frac{\gamma}{a_0 \delta} \frac{da_0}{dt} - \frac{1}{2} \frac{\gamma^2}{\delta^2} - \frac{1}{\delta} \frac{d\gamma}{dt} + \frac{\gamma}{\delta^2} \frac{d\delta}{dt} \right) z,$$

and $W = z_1 dz_2/d\tau - z_2 dz_1/d\tau$, with $d\tau = \alpha/\delta dt$ and α , δ and γ certain t -dependent functions satisfying (5.13).

Many other similar results can be obtained in an analogous manner. For instance, the theory of Lie systems was also used in [12] to prove that the general solution of a Milne–Pinney equation (7.1) can be written as

$$y(\tau) = \sqrt{\frac{[k_1(x_1(\tau) - x_2(\tau)) - k_2(x_1(\tau) - x_3(\tau))]^2 - a_0(0)^2[x_2(\tau) - x_3(\tau)]/4}{(k_2 - k_1)(x_2(\tau) - x_3(\tau))(x_2(\tau) - x_1(\tau))(x_1(\tau) - x_3(\tau))}},$$

where $x_1(\tau)$, $x_2(\tau)$ and $x_3(\tau)$ are three different particular solutions of the Riccati equation

$$\frac{dx}{d\tau} = -\omega(t(\tau)) - x^2.$$

Proceeding as before, we can describe the general solution of a second-order Gambier equation (7.2) in terms of solutions of these Riccati equations. In addition, by applying the theory of Lie systems [19] to Milne–Pinney equations (7.1), we can obtain many other results about subfamilies of second-order Gambier equations. In addition, the relation between second-order Gambier equations and Milne–Pinney equations enables us to obtain several other results in a simple way.

Proposition 7.2. *The second-order Gambier equation (5.11) with $2a_2 = a_0^2 > 0$, i.e.*

$$\frac{d^2 x}{dt^2} = \frac{3}{2x} \left(\frac{dx}{dt} \right)^2 + \frac{1}{a_0} \frac{da_0}{dt} \frac{dx}{dt} + \frac{a_0^2}{2} x^3 - a_0^2 x, \quad (7.3)$$

can be transformed into the integrable Milne–Pinney equation

$$\frac{d^2 y}{d\tau^2} - \frac{y}{2} + \frac{1}{4y^3} = 0 \quad (7.4)$$

under the transformation $x = 1/y^2$ and a t -reparametrisation $d\tau = a_0 dt$.

Note 7.1. For second-order Gambier equations (5.11) with $a_0 = 0$, the transformations given in the above proposition map these equations into harmonic oscillators.

Proposition 7.3. *A constant of motion of (7.4) is given by*

$$I_{MP} = \frac{1}{2} \left[\left(\frac{dy}{d\tau} \right)^2 - \left(\frac{y^2}{2} + \frac{1}{4y^2} \right) \right].$$

Using $y = x^{-1/2}$ transformation, we obtain a constant of motion for the special second-order Gambier equation (7.3) of the form

$$I_{2G} = \frac{1}{4} \frac{1}{x^3 a_0^2} \left(\frac{dx}{dt} \right)^2 - \left(\frac{1}{2x} + \frac{x}{4} \right). \quad (7.5)$$

Thus, we say the special equation (7.3), which yields a constant of motion, is an integrable deformation of the Milne–Pinney equation.

Note 7.2. Recall that Theorem 6.1 can be applied to those equations (5.11) where $a_2 = -\lambda a_0^2/a_0^2(0)$ giving rise to the constant of motion (6.4). Observe that (7.3) is of this type with $\lambda = -a_0(0)^2/2$. Then, the constant of motion (6.4) is, up to a multiplicative constant, the same as (7.5). Despite this, the above illustrates how certain results can readily be obtained through Milne–Pinney equations.

8 A second quasi-Lie schemes approach to second-order Gambier equations

Apart from our first approach, we can provide a second quasi-Lie scheme to study second-order Gambier equations. This one is motivated by the fact that some cases of these equations, namely

$$\frac{d^2x}{dt^2} = - \left(3x \frac{dx}{dt} + x^3 \right) + f(t) + g(t)x + h(t) \left(x^2 + \frac{dx}{dt} \right),$$

for arbitrary t -dependent functions f , g and h , form a particular subclass of second-order Riccati equations that are *SODE Lie systems* [13, 16]. That is, when written as first-order systems

$$\begin{aligned} \frac{dx}{dt} &= v, \\ \frac{dv}{dt} &= - (3xv + x^3) + f(t) + g(t)x + h(t) (x^2 + v), \end{aligned} \quad (8.1)$$

they become Lie systems. Indeed, these systems possess a Vessiot–Guldberg Lie algebra V_0 isomorphic to $\mathfrak{sl}(3, \mathbb{R})$ given by [10, 29]

$$\begin{aligned} X_1 &= v \frac{\partial}{\partial x} - (3xv + x^3) \frac{\partial}{\partial v}, & X_2 &= \frac{\partial}{\partial v}, & X_3 &= -\frac{\partial}{\partial x} + 3x \frac{\partial}{\partial v}, \\ X_4 &= x \frac{\partial}{\partial x} - 2x^2 \frac{\partial}{\partial v}, & X_5 &= (v + 2x^2) \frac{\partial}{\partial x} - x(v + 3x^2) \frac{\partial}{\partial v}, \\ X_6 &= 2x(v + x^2) \frac{\partial}{\partial x} + 2(v^2 - x^4) \frac{\partial}{\partial v}, & X_7 &= \frac{\partial}{\partial x} - x \frac{\partial}{\partial v}, & X_8 &= 2x \frac{\partial}{\partial x} + 4v \frac{\partial}{\partial v}. \end{aligned}$$

In terms of these vector fields, (8.1) is the associated system to the t -dependent vector field

$$X_1 + f(t)X_2 + \frac{1}{2}g(t)(X_3 + X_7) + \frac{h(t)}{4}(-2X_4 + X_8).$$

Table 2. Lie brackets $[Y_i, Y_j]$ with $i = 4, 8, 11$ and $j = 12, \dots, 17$.

$[\cdot, \cdot]$	Y_{12}	Y_{13}	Y_{14}	Y_{15}	Y_{16}	Y_{17}
Y_4	0	0	Y_{14}	0	$-Y_{16}$	Y_{17}
Y_8	$-Y_9$	$-Y_7$	$Y_{13} - Y_3$	$-Y_6$	0	$2Y_3$
Y_{11}	$-Y_{12}$	Y_{13}	0	$2Y_{15}$	$4Y_{16}$	0

Therefore, it is natural to analyse which Gambier vector fields X can be transformed through a quasi-Lie scheme into one of these Lie systems.

Despite the interest of the previous idea, the quasi-Lie scheme provided in the previous section cannot be employed to determine all quasi-Lie systems of this type. The reason is that V_0 is not included in V_G and, therefore, there exists no $g \in \mathcal{G}(W_G)$ such that $g_\star X \in V_0(C^\infty(\mathbb{R})) \subset V_G(C^\infty(\mathbb{R}))$. To solve this drawback, we now determine a new quasi-Lie scheme $S(W_G, V'_G)$ such that the space V'_G contains $V_G + V_0$. This can be done by defining V'_G as the \mathbb{R} -linear space generated by the elements of V_G and the vector fields

$$\begin{aligned} Y_{12} &= \frac{\partial}{\partial x}, & Y_{13} &= x^2 \frac{\partial}{\partial x}, & Y_{14} &= xv \frac{\partial}{\partial x}, \\ Y_{15} &= x^3 \frac{\partial}{\partial x}, & Y_{16} &= x^4 \frac{\partial}{\partial v}, & Y_{17} &= v^2 \frac{\partial}{\partial v}. \end{aligned}$$

Indeed, using Tables 1 and 2, we readily obtain that $[W_G, V'_G] \subset V'_G$. Moreover, as we already know that W_G is a Lie algebra, $S(W_G, V'_G)$ becomes a quasi-Lie scheme.

Since we impose that $g_\star X$, which is given by (4.2), must be of the form (8.1) up to a t -reparametrisation, we obtain

$$g_\star X = \sum_{\alpha=1}^{11} \bar{b}_\alpha(t) Y_\alpha = \xi(t) [(Y_1 - 3Y_3 - Y_6) + f(t)Y_9 + g(t)Y_8 + h(t)(Y_7 + Y_4)], \quad (8.2)$$

for a certain t -dependent non-vanishing function $\xi(t)$. Taking into account that the vector fields Y_1, \dots, Y_{11} are linearly independent over \mathbb{R} , we see that

$$\bar{b}_2 = 0, \quad \bar{b}_5 = 0, \quad \bar{b}_{10} = 0, \quad \bar{b}_{11} = 0.$$

Bearing in mind the form of the coefficients \bar{b}_α given by (4.3) and recalling that $\alpha > 0$, we see that $\bar{b}_2 = 0$ entails $n = 1$. Meanwhile, conditions $\bar{b}_5 = 0$ and $\bar{b}_{11} = 0$ entail $\sigma = 0$ and

$$\frac{1}{\alpha} \frac{d\alpha}{dt} = \frac{\gamma}{\delta}, \quad (8.3)$$

respectively. In turn, $\sigma = 0$ also entails that $\bar{b}_9 = \bar{b}_{10} = 0$. Moreover, from (8.2) we also see that

$$\bar{b}_3 = -3\bar{b}_1, \quad \bar{b}_6 = -\bar{b}_1, \quad \bar{b}_7 = \bar{b}_4.$$

From these conditions and $n = 1, \sigma = 0$, we obtain

$$\frac{a_0}{\alpha} = -\frac{\alpha}{\delta}, \quad \frac{a_0^2 \delta}{\alpha^3} = \frac{\alpha}{\delta}, \quad \frac{1}{\delta} \frac{d\delta}{dt} = \frac{2\gamma}{\delta} - \frac{1}{a_0} \frac{da_0}{dt}. \quad (8.4)$$

Since $\delta, \alpha > 0$, it can readily be seen that the second equation is an immediate consequence of the first one, which in turn implies $a_0 = -\alpha^2/\delta$. Using that $\delta(0) = \alpha(0) = 1$, we obtain $a_0(0) = -1$. Previous results along with (8.3) ensure that the last condition in (8.4) holds. Hence, \bar{b}_1, \bar{b}_4 and \bar{b}_8 become

$$\bar{b}_1 = -\frac{a_0}{\alpha}, \quad \bar{b}_4 = a_1 - \frac{1}{a_0} \frac{da_0}{dt} + \frac{3}{\alpha} \frac{d\alpha}{dt},$$

$$\bar{b}_8 = -\frac{a_2\alpha}{a_0} + \frac{a_1}{a_0} \frac{d\alpha}{dt} + \frac{2}{a_0\alpha} \left(\frac{d\alpha}{dt} \right)^2 - \frac{1}{a_0} \frac{d^2\alpha}{dt^2}, \quad (8.5)$$

and the remaining coefficients \bar{b}_α are either zero or can be obtained from them.

Proposition 8.1. *Every Gambier vector field X is a quasi-Lie system relative to the quasi-Lie scheme $S(W_G, V'_G)$ and the Lie algebra V_0 if and only if $n = 1$, $a_0(0) = -1$ and $\sigma = 0$. In such a case, every t -dependent transformation $g \in \mathcal{G}(W_G)$ given by*

$$\bar{x} = \alpha x, \quad \bar{v} = -\frac{\alpha}{a_0} \left(\frac{d\alpha}{dt} x + \alpha v \right), \quad (8.6)$$

transforms X into a Lie system

$$\begin{aligned} \frac{d\bar{x}}{d\tau} &= \bar{v}, \\ \frac{d\bar{v}}{d\tau} &= -3\bar{x}\bar{v} - \bar{x}^3 + \bar{b}_4\bar{b}_1^{-1}\bar{x} + \bar{b}_8\bar{b}_1^{-1}(\bar{x}^2 + \bar{v}), \end{aligned} \quad (8.7)$$

where $d\tau = -a_0 dt/\alpha$ and \bar{b}_1 , \bar{b}_4 and \bar{b}_8 are given by (8.5).

9 Exact solutions for several second-order Gambier equations

As a result of Proposition 8.1, we can apply to (8.7) the techniques of the theory of Lie systems so as to devise, by inverting the change of variables (8.6), new properties of the Gambier equations related to second-order Riccati equations. For instance, we obtain new exact solutions of some of these families of second-order Gambier equations.

In the study of every Lie system, special relevance takes the algebraic structure of its Vessiot–Guldberg Lie algebras. When a Lie system possesses a solvable Vessiot–Guldberg Lie algebra, we can apply several methods to explicitly integrate it (cf. [9, 13, 19]). Otherwise, the general solution of the Lie system usually needs to be obtained through approximative methods [51] or expressed in terms of solutions of other Lie systems [12].

System (8.7) is a special case of a Lie system related to a Vessiot–Guldberg Lie algebra V_0 isomorphic to $\mathfrak{sl}(3, \mathbb{R})$, which is simple and therefore difficult to integrate explicitly. Nevertheless, we can prove that (8.7) is in general related to a Vessiot–Guldberg Lie algebra that is a proper Lie subalgebra of V_0 . Moreover, we next prove that X is related to a solvable Vessiot–Guldberg Lie algebra in some particular cases that can be explicitly integrated.

System (8.7) describes the integral curves of the t -dependent vector field

$$X_t = Z_1 + \bar{b}_4\bar{b}_1^{-1}Z_2 + \bar{b}_8\bar{b}_1^{-1}Z_3.$$

For generic t -dependent functions \bar{b}_1 , \bar{b}_4 and \bar{b}_8 , the above system admits a Vessiot–Guldberg Lie algebra spanned by $Z_1 = X_1$, $Z_2 = (X_3 + X_7)/2$, $Z_3 = (X_8 - 2X_4)/4$ and their successive Lie brackets, which generates a proper Lie subalgebra V of V_0 . More specifically, we first have

$$[Z_1, Z_2] = Z_3 - Z_4, \quad [Z_1, Z_3] = -(Z_1 + Z_5)/2, \quad [Z_1, Z_5] = Z_6,$$

where $Z_4 = X_4$, $Z_5 = X_5$, $Z_6 = X_6$. This shows that the smallest Lie algebra containing Z_1 , Z_2 and Z_3 must include these vector fields and Z_4 , Z_5 and Z_6 . Since these vector fields additionally satisfy the following commutation relations

$$\begin{aligned} [Z_1, Z_4] &= Z_5, & [Z_3, Z_4] &= 0, & [Z_1, Z_6] &= 0, & [Z_2, Z_3] &= Z_2, & [Z_2, Z_4] &= -Z_2, \\ [Z_2, Z_5] &= Z_4 - Z_3, & [Z_2, Z_6] &= Z_5 - Z_1, & [Z_3, Z_5] &= (Z_5 + Z_1)/2, & [Z_3, Z_6] &= Z_6, \\ [Z_4, Z_5] &= -Z_1, & [Z_4, Z_6] &= 0, & [Z_5, Z_6] &= 0, \end{aligned}$$

we see that they span a six-dimensional proper Lie subalgebra V of V_0 . Moreover, it is easy to show that $V \simeq \langle Z_1 + Z_5, Z_3 - Z_4, Z_2 \rangle \oplus_S \langle Z_6, Z_1 - Z_5, Z_3 + Z_4 \rangle$, where $\langle Z_1 + Z_5, Z_3 - Z_4, Z_2 \rangle \simeq \mathfrak{sl}(2, \mathbb{R})$, namely

$$\begin{aligned} [Z_1 + Z_5, Z_3 - Z_4] &= -2(Z_1 + Z_5), \\ [Z_2, Z_3 - Z_4] &= 2Z_2, \quad [Z_1 + Z_5, Z_2] = 2(Z_3 - Z_4), \end{aligned}$$

the linear space $\langle Z_6, Z_1 - Z_5, Z_3 + Z_4 \rangle$ is a solvable ideal of V and \oplus_S stands for the semidirect sum of $\langle Z_1 + Z_5, Z_3 - Z_4, Z_2 \rangle$ by $\langle Z_6, Z_1 - Z_5, Z_3 + Z_4 \rangle$.

Hence, X is related to a non-solvable Vessiot–Guldberg Lie algebra and it is not known a general method to explicitly integrate X for arbitrary \bar{b}_1 , \bar{b}_4 and \bar{b}_8 . Nevertheless, we can focus on a particular instance of these functions so that X can be related to a solvable Vessiot–Guldberg Lie algebra V_1 . For example, consider the case $\bar{b}_4 = 0$. We then have

$$V_1 = \langle Z_1, Z_3, Z_1 + Z_5, Z_6 \rangle.$$

Note that the derived series of V_1 become zero, i.e. $\mathcal{D}^1 \equiv [V_1, V_1] = \langle Z_1 + Z_5, Z_6 \rangle$, $\mathcal{D}^2 = 0$. In other words, V_1 is solvable and we can expect to integrate it through some method. Let us do so through the so-called *mixed superposition rules*, i.e. a generalisation of superposition rules describing the general solution of a Lie system in terms of several particular solutions of other systems and a set of constants [29]. In our case, it is known (cf. [29, p. 194]) that the general solution of (8.7) can be written in the form

$$\bar{x}(\tau) = \frac{\lambda_1 v_{y_1}(\tau) + \lambda_2 v_{y_2}(\tau) + \lambda_3 v_{y_3}(\tau)}{\lambda_1 y_1(\tau) + \lambda_2 y_2(\tau) + \lambda_3 y_3(\tau)}, \quad \bar{v}(\tau) = \frac{d}{d\tau} \left(\frac{\lambda_1 v_{y_1}(\tau) + \lambda_2 v_{y_2}(\tau) + \lambda_3 v_{y_3}(\tau)}{\lambda_1 y_1(\tau) + \lambda_2 y_2(\tau) + \lambda_3 y_3(\tau)} \right),$$

where $(\lambda_1, \lambda_2, \lambda_3) \in \mathbb{R}^3 - \{(0, 0, 0)\}$ and $(y_i(\tau), v_{y_i}(\tau), a_{y_i}(\tau))$, with $i = 1, 2, 3$, are linearly independent solutions of the linear Lie system

$$\frac{dy}{d\tau} = v_y, \quad \frac{dv_y}{d\tau} = a_y, \quad \frac{da_y}{d\tau} = \bar{b}_8 \bar{b}_1^{-1} a_y.$$

The above Lie system is related to a Vessiot–Guldberg Lie algebra spanned by

$$W_1 = v_y \frac{\partial}{\partial y} + a_y \frac{\partial}{\partial v_y}, \quad W_2 = a_y \frac{\partial}{\partial a_y}, \quad W_3 = 2a_y \frac{\partial}{\partial v_y}, \quad W_4 = -2a_y \frac{\partial}{\partial y},$$

which close the same commutation relations as Z_1 , Z_3 , $Z_1 + Z_5$ and Z_6 . Hence, this system is related to a solvable Vessiot–Guldberg Lie algebra and it can easily be integrated:

$$a_y = \exp \left(\int^\tau \bar{b}_8(\tau') \bar{b}_1^{-1}(\tau') d\tau' \right), \quad v_y = \int^\tau a_y(\tau') d\tau', \quad y = \int^\tau v_y(\tau') d\tau'.$$

Since we can assume

$$\lambda_1 y_1(\tau) + \lambda_2 y_2(\tau) + \lambda_3 y_3(\tau) = \int^\tau \int^{\tau'} \exp \left(\int^{\tau''} \bar{b}_8(\tau''') \bar{b}_1^{-1}(\tau''') d\tau''' \right) d\tau'' d\tau',$$

we obtain the solution of (8.7) for $\bar{b}_4 = 0$, i.e.

$$\bar{x}(\tau) = \frac{\int^\tau \exp \left(\int^{\tau'} \bar{b}_8(\tau'') \bar{b}_1^{-1}(\tau'') d\tau'' \right) d\tau'}{\int^\tau \int^{\tau'} \exp \left(\int^{\tau''} \bar{b}_8(\tau''') \bar{b}_1^{-1}(\tau''') d\tau''' \right) d\tau'' d\tau'}, \quad \bar{v}(\tau) = \frac{d\bar{x}}{d\tau}(\tau).$$

From above results, we have the following proposition.

Proposition 9.1. *Given a second-order Gambier equation with $n = 1$, $a_0(0) = -1$, $\sigma = 0$ and a particular solution $\alpha(t)$ with $\alpha(0) = 1$ of*

$$a_1 = \frac{1}{a_0} \frac{da_0}{dt} - \frac{3}{\alpha} \frac{d\alpha}{dt}, \quad (9.1)$$

its general solution reads

$$x(\tau(t)) = \frac{1}{\alpha(\tau(t))} \frac{\int^{\tau(t)} \exp\left(\int^{\tau'} \bar{b}_8(\tau'') \bar{b}_1^{-1}(\tau'') d\tau''\right) d\tau'}{\int^{\tau(t)} \int^{\tau'} \exp\left(\int^{\tau''} \bar{b}_8(\tau''') \bar{b}_1^{-1}(\tau''') d\tau'''\right) d\tau'' d\tau'},$$

where $d\tau = -a_0 dt/\alpha$ and \bar{b}_1 and \bar{b}_8 are given by (8.5).

Note 9.1. Since equation (9.1) can be easily integrated, the above proposition shows that we can integrate exactly every second-order Gambier equation with $n = 1$, $a_0(0) = -1$ and $\sigma = 0$.

10 Conclusions

Two quasi-Lie schemes have been introduced to analyse the second-order Gambier equations. Our first quasi-Lie scheme has been used to recover previous results concerning the reduction of such equations to reduced canonical forms from a geometric clarifying approach, which allowed us to fill a gap in the previous literature. This quasi-Lie scheme also led to determine some quasi-Lie systems related to certain second-order Gambier equations, which enabled us to transform them into second-order Kummer–Schwarz equations. We have expressed the general solutions of such second-order Gambier equations in terms of particular solutions of t -dependent frequency harmonic oscillators and Riccati equations. Additionally, new constants of motion were derived for some of them.

The introduction of a second quasi-Lie scheme resulted in the description of an integrable family of second-order Gambier equations related to second-order Riccati equations.

Acknowledgements

The research of J.F. Cariñena and J. de Lucas was supported by the Polish National Science Centre under the grant HARMONIA Nr 2012/04/M/ST1/00523. They also acknowledge partial financial support by research projects MTM–2009–11154 (MEC) and E24/1 (DGA). J. de Lucas would like to thank for a research grant FMI40/10 (DGA) to accomplish a research stay in the University of Zaragoza.

References

- [1] Anderson R.L., Harnad J., Winternitz P., Group theoretical approach to superposition rules for systems of Riccati equations, *Lett. Math. Phys.* **5** (1981), 143–148.
- [2] Beekers J., Gagnon L., Hussin V., Winternitz P., Nonlinear differential equations and Lie superalgebras, *Lett. Math. Phys.* **13** (1987), 113–120.
- [3] Beekers J., Gagnon L., Hussin V., Winternitz P., Superposition formulas for nonlinear superequations, *J. Math. Phys.* **31** (1990), 2528–2534.
- [4] Berkovich L.M., Canonical forms of ordinary linear differential equations, *Arch. Math. (Brno)* **24** (1988), 25–42.
- [5] Berkovich L.M., The generalized Emden–Fowler equation, in Proceedings of Second International Conference “Symmetry in Nonlinear Mathematical Physics” (Kyiv, 1997), Editors M.I. Shkil, A.G. Nikitin, V.M. Boyko, Institute of Mathematics, Kyiv, 1997, 155–163.

- [6] Blázquez-Sanz D., Morales-Ruiz J.J., Lie's reduction method and differential Galois theory in the complex analytic context, *Discrete Contin. Dyn. Syst.* **32** (2012), 353–379, [arXiv:0901.4479](#).
- [7] Blázquez-Sanz D., Morales-Ruiz J.J., Local and global aspects of Lie superposition theorem, *J. Lie Theory* **20** (2010), 483–517, [arXiv:0901.4478](#).
- [8] Bouquet S.E., Feix M.R., Leach P.G.L., Properties of second-order ordinary differential equations invariant under time translation and self-similar transformation, *J. Math. Phys.* **32** (1991), 1480–1490.
- [9] Cariñena J.F., de Lucas J., Lie systems: theory, generalisations, and applications, *Dissertationes Math.* **479** (2011), 162 pages, [arXiv:1103.4166](#).
- [10] Cariñena J.F., de Lucas J., Superposition rules and second-order Riccati equations, *J. Geom. Mech.* **3** (2011), 1–22, [arXiv:1007.1309](#).
- [11] Cariñena J.F., de Lucas J., Rañada M.F., A geometric approach to integrability of Abel differential equations, *Internat. J. Theoret. Phys.* **50** (2011), 2114–2124, [arXiv:1012.2257](#).
- [12] Cariñena J.F., de Lucas J., Rañada M.F., Recent applications of the theory of Lie systems in Ermakov systems, *SIGMA* **4** (2008), 031, 18 pages, [arXiv:0803.1824](#).
- [13] Cariñena J.F., de Lucas J., Sardón C., A new Lie-systems approach to second-order Riccati equations, *Int. J. Geom. Methods Mod. Phys.* **9** (2012), 1260007, 8 pages, [arXiv:1110.3298](#).
- [14] Cariñena J.F., de Lucas J., Sardón C., Lie–Hamilton systems: theory and applications, *Int. J. Geom. Methods Mod. Phys.*, to appear, [arXiv:1211.6991](#).
- [15] Cariñena J.F., Grabowski J., de Lucas J., Quasi-Lie schemes: theory and applications, *J. Phys. A: Math. Theor.* **42** (2009), 335206, 20 pages, [arXiv:0810.1160](#).
- [16] Cariñena J.F., Grabowski J., de Lucas J., Superposition rules for higher order systems and their applications, *J. Phys. A: Math. Theor.* **45** (2012), 185202, 26 pages, [arXiv:1111.4070](#).
- [17] Cariñena J.F., Grabowski J., Marmo G., Lie–Scheffers systems: a geometric approach, Bibliopolis, Naples, 2000.
- [18] Cariñena J.F., Grabowski J., Marmo G., Superposition rules, Lie theorem, and partial differential equations, *Rep. Math. Phys.* **60** (2007), 237–258, [math-ph/0610013](#).
- [19] Cariñena J.F., Grabowski J., Ramos A., Reduction of time-dependent systems admitting a superposition principle, *Acta Appl. Math.* **66** (2001), 67–87.
- [20] Cariñena J.F., Leach P.G.L., de Lucas J., Quasi-Lie schemes and Emden–Fowler equations, *J. Math. Phys.* **50** (2009), 103515, 21 pages, [arXiv:0908.2236](#).
- [21] Cariñena J.F., Marmo G., Nasarre J., The nonlinear superposition principle and the Wei–Norman method, *Internat. J. Modern Phys. A* **13** (1998), 3601–3627, [physics/9802041](#).
- [22] Cariñena J.F., Ramos A., Integrability of the Riccati equation from a group-theoretical viewpoint, *Internat. J. Modern Phys. A* **14** (1999), 1935–1951, [math-ph/9810005](#).
- [23] Cariñena J.F., Rañada M.F., Santander M., Lagrangian formalism for nonlinear second-order Riccati systems: one-dimensional integrability and two-dimensional superintegrability, *J. Math. Phys.* **46** (2005), 062703, 18 pages, [math-ph/0505024](#).
- [24] Conte R., Singularities of differential equations and integrability, in *An Introduction to Methods of Complex Analysis and Geometry for Classical Mechanics and Non-Linear Waves* (Chamonix, 1993), Frontières, Gif, 1994, 49–143.
- [25] Ermakov V.P., Second-order differential equations: conditions of complete integrability, *Appl. Anal. Discrete Math.* **2** (2008), 123–145.
- [26] Flores-Espinoza R., Periodic first integrals for Hamiltonian systems of Lie type, *Int. J. Geom. Methods Mod. Phys.* **8** (2011), 1169–1177, [arXiv:1004.1132](#).
- [27] Flores-Espinoza R., de Lucas J., Vorobiev Y.M., Phase splitting for periodic Lie systems, *J. Phys. A: Math. Theor.* **43** (2010), 205208, 11 pages, [arXiv:0910.2575](#).
- [28] González-López A., Kamran N., Olver P.J., Lie algebras of vector fields in the real plane, *Proc. London Math. Soc. (3)* **64** (1992), 339–368.
- [29] Grabowski J., de Lucas J., Mixed superposition rules and the Riccati hierarchy, *J. Differential Equations* **254** (2013), 179–198, [arXiv:1203.0123](#).
- [30] Grammaticos B., Ramani A., Lafortune S., The Gambier mapping, revisited, *Phys. A* **253** (1998), 260–270, [solv-int/9804011](#).

- [31] Grundland A.M., Levi D., On higher-order Riccati equations as Bäcklund transformations, *J. Phys. A: Math. Gen.* **32** (1999), 3931–3937.
- [32] Guha P., Ghose Choudhury A., Grammaticos B., Dynamical studies of equations from the Gambier family, *SIGMA* **7** (2011), 028, 15 pages, [arXiv:1103.4210](https://arxiv.org/abs/1103.4210).
- [33] Harnad J., Winternitz P., Anderson R.L., Superposition principles for matrix Riccati equations, *J. Math. Phys.* **24** (1983), 1062–1072.
- [34] Ibragimov N.H., Utilization of canonical variables for integration of systems of first-order differential equations, *Arch. ALGA* **6** (2009), 1–18.
- [35] Ince E.L., Ordinary differential equations, Dover Publications, New York, 1944.
- [36] Inselberg A., Superpositions for nonlinear operators. I. Strong superpositions and linearizability, *J. Math. Anal. Appl.* **40** (1972), 494–508.
- [37] Karasu A., Leach P.G.L., Nonlocal symmetries and integrable ordinary differential equations: $\ddot{x} + 3x\dot{x} + x^3 = 0$ and its generalizations, *J. Math. Phys.* **50** (2009), 073509, 17 pages.
- [38] Königsberger L., Über die einer beliebigen Differentialgleichung erster Ordnung angehörigen selbständigen Transcendenten, *Acta Math.* **3** (1883), 1–48.
- [39] Lafortune S., Grammaticos B., Ramani A., Constructing integrable third-order systems: the Gambier approach, *Inverse Problems* **14** (1998), 287–298, [solv-int/9705007](https://arxiv.org/abs/solv-int/9705007).
- [40] Lafortune S., Grammaticos B., Ramani A., Winternitz P., Discrete systems related to some equations of the Painlevé–Gambier classification, *Phys. Lett. A* **270** (2000), 55–61, [nlin.SI/0104019](https://arxiv.org/abs/nlin.SI/0104019).
- [41] Lax P.D., Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.* **21** (1968), 467–490.
- [42] Leach P.G.L., Feix M.R., Bouquet S., Analysis and solution of a nonlinear second-order differential equation through rescaling and through a dynamical point of view, *J. Math. Phys.* **29** (1988), 2563–2569.
- [43] Leach P.G.L., Govinder K.S., On the uniqueness of the Schwarzian and linearisation by nonlocal contact transformation, *J. Math. Anal. Appl.* **235** (1999), 84–107.
- [44] Levi D., Ragnisco O. (Editors), SIDE III – symmetries and integrability of difference equations, *CRM Proceedings & Lecture Notes*, Vol. 25, Amer. Math. Soc., Providence, RI, 2000.
- [45] Lie S., Theorie der Transformationsgruppen I, *Math. Ann.* **16** (1880), 441–528.
- [46] Lie S., Scheffers G., Vorlesungen über kontinuierliche Gruppen mit geometrischen und anderen Anwendungen, Teubner, Leipzig, 1893.
- [47] Olver P.J., Applications of Lie groups to differential equations, *Graduate Texts in Mathematics*, Vol. 107, 2nd ed., Springer-Verlag, New York, 1993.
- [48] Olver P.J., Symmetry and explicit solutions of partial differential equations, *Appl. Numer. Math.* **10** (1992), 307–324.
- [49] Olver P.J., Rosenau P., Group-invariant solutions of differential equations, *SIAM J. Appl. Math.* **47** (1987), 263–278.
- [50] Pickering A., The singular manifold method revisited, *J. Math. Phys.* **37** (1996), 1894–1927.
- [51] Pietrzkowski G., Explicit solutions of the \mathfrak{a}_1 -type Lie–Scheffers system and a general Riccati equation, *J. Dyn. Control Syst.* **18** (2012), 551–571.
- [52] Pinney E., The nonlinear differential equation $y'' + p(x)y + cy^{-3} = 0$, *Proc. Amer. Math. Soc.* **1** (1950), 681.
- [53] Rourke D.E., Augustine M., Exact linearization of the radiation-damped spin system, *Phys. Rev. Lett.* **84** (2000), 1685–1688.
- [54] Shnider S., Winternitz P., Classification of systems of nonlinear ordinary differential equations with superposition principles, *J. Math. Phys.* **25** (1984), 3155–3165.
- [55] Vessiot E., Sur une classe d'équations différentielles, *Ann. Sci. École Norm. Sup. (3)* **10** (1893), 53–64.
- [56] Winternitz P., Lie groups and solutions of nonlinear differential equations, in Nonlinear Phenomena (Oaxtepec, 1982), *Lecture Notes in Phys.*, Vol. 189, Springer, Berlin, 1983, 263–331.