

## Linking first occurrence polynomials over $\mathbb{F}_p$ by Steenrod operations

Pham Anh Minh  
 Grant Walker

**Abstract** This paper provides analogues of the results of [16] for odd primes  $p$ . It is proved that for certain irreducible representations  $L(\lambda)$  of the full matrix semigroup  $M_n(\mathbb{F}_p)$ , the first occurrence of  $L(\lambda)$  as a composition factor in the polynomial algebra  $\mathbf{P} = \mathbb{F}_p[x_1, \dots, x_n]$  is linked by a Steenrod operation to the first occurrence of  $L(\lambda)$  as a submodule in  $\mathbf{P}$ . This operation is given explicitly as the image of an admissible monomial in the Steenrod algebra  $A_p$  under the canonical anti-automorphism  $\tau$ . The first occurrences of both kinds are also linked to higher degree occurrences of  $L(\lambda)$  by elements of the Milnor basis of  $A_p$ .

**AMS Classification** 55S10; 20C20

**Keywords** Steenrod algebra, anti-automorphism,  $p$ -truncated polynomial algebra  $\mathbf{T}$ ,  $\mathbf{T}$ -regular partition/representation

### 1 Introduction

Our aim is to obtain results corresponding to those of [16] for the case where the prime  $p > 2$ . In this we are only partly successful. The main theorem of [16] gives a Steenrod operation which links the first occurrence of each irreducible representation  $L(\lambda)$  of the full matrix semigroup  $M_n(\mathbb{F}_2)$  in the polynomial algebra  $\mathbf{P} = \mathbb{F}_2[x_1, \dots, x_n]$  with the first occurrence of  $L(\lambda)$  as a submodule in  $\mathbf{P}$ . Here  $M_n(\mathbb{F}_2)$  acts on  $\mathbf{P}$  on the right by linear substitutions, which commute with the action of the Steenrod algebra  $A_2$  on  $\mathbf{P}$  on the left. By 'first occurrence' we have in mind the decomposition  $\mathbf{P} = \sum_{d \geq 0} \mathbf{P}^d$ , where  $\mathbf{P}^d$  is the module of homogeneous polynomials of total degree  $d$ , and the known facts that there are minimum degrees  $d_c(\lambda)$  and  $d_s(\lambda)$  in which  $L(\lambda)$  occurs, uniquely in each case, as a composition factor and as a submodule respectively.

For an odd prime  $p$ , we have again the commuting actions of  $M_n = M_n(\mathbb{F}_p)$  on the right of the polynomial algebra  $\mathbf{P} = \mathbb{F}_p[x_1, \dots, x_n]$  and the algebra  $A_p$

of Steenrod  $p$ th powers (no Bocksteins) on the left. We refer to  $A_p$ , somewhat inaccurately, as the Steenrod algebra, and grade it so that  $P^r$  raises degree by  $r(p-1)$ . There are  $p^n$  isomorphism classes of irreducible  $\mathbb{F}_p[M_n]$ -modules  $L(\lambda)$ , indexed by partitions  $\lambda = (\lambda_1; \lambda_2; \dots; \lambda_n)$ , which are column  $p$ -regular, i.e.  $0 \leq \lambda_i - \lambda_{i+1} \leq p-1$  for  $1 \leq i < n$ , where  $\lambda_{n+1} = 0$  [8, 9, 10]. The problem solved in [16] is certainly more difficult in this context. The submodule degree  $d_s(\lambda)$  has recently been determined [12] for every irreducible representation  $L(\lambda)$  of  $M_n$ , but  $d_c(\lambda)$  is not known in general. In particular, the first occurrence problem appears to be difficult even for the 1-dimensional representations  $\det^k$ ,  $1 \leq k \leq p-3$ ,  $p > 3$ , see [2, 3], although it is solved for  $\det^{p-2}$  [1]. (The partition indexing  $\det^k$  is  $(k; \dots; k) = (k^n)$ , i.e.  $k$  repeated  $n$  times.) Further, it is not known in general whether  $\mathbf{P}^{d_c(\lambda)}$  has a unique composition factor isomorphic to  $L(\lambda)$ . Here we identify a class of irreducible representations  $L(\lambda)$  which behave systematically. Since they arise naturally by considering tensor powers of the  $p$ -truncated polynomial algebra  $\mathbf{T} = \mathbf{P} = (x_1^p; \dots; x_n^p)$ , we call them  $\mathbf{T}$ -regular.

Our main result, Theorem 5.7, gives a Steenrod operation  $\sigma(\lambda)$  which links the first occurrence and the first submodule occurrence in  $\mathbf{P}$  of a  $\mathbf{T}$ -regular  $L(\lambda)$ . This determines  $d_c(\lambda)$  in the  $\mathbf{T}$ -regular case. The operation  $\sigma(\lambda)$  is given explicitly as the image of an admissible monomial under the canonical anti-automorphism of  $A_p$ . Calculations for  $n = 3$  suggest that such an operation  $\sigma(\lambda)$  may exist for every irreducible representation  $L(\lambda)$  of  $M_n$ , but we do not pursue this here. Tri [14] has given an ‘algebraic’ alternative to this ‘topological’ method of finding  $d_c(\lambda)$ , using coefficient functions of  $\mathbb{F}_p[M_n]$ -modules.

For  $p = 2$ ,  $\mathbf{T}$  may be identified with the exterior algebra  $(x_1; \dots; x_n)$ , and all the irreducible representations  $L(\lambda)$  of  $M_n$  are  $\mathbf{T}$ -regular. For  $p > 2$ , the only irreducible 1-dimensional  $\mathbf{T}$ -regular representations of  $M_n$  are the ‘trivial’ representation, in which all matrices act as 1, and the  $\det^{p-1}$  representation, in which non-singular matrices act as 1 and singular matrices as 0. The ‘trivial’ representation, for which  $\lambda = (0)$ , occurs in  $\mathbf{P}$  only as  $\mathbf{P}^0$ , the constant polynomials. Our key example is the  $\det^{p-1}$  representation. This occurs first as a composition factor as the top degree  $\mathbf{T}^{n(p-1)}$  of  $\mathbf{T}$ , where it is generated by the monomial  $(x_1 x_2 \dots x_n)^{p-1}$  modulo  $p$ th powers, and first as a submodule in degree  $\rho_n = (p^n - 1)/(p - 1)$ , where it is generated by the Vandermonde determinant

$$w(n) = \begin{vmatrix} x_1 & x_2 & \dots & x_n \\ x_1^p & x_2^p & \dots & x_n^p \\ \vdots & \vdots & \ddots & \vdots \\ x_1^{p^{n-1}} & x_2^{p^{n-1}} & \dots & x_n^{p^{n-1}} \end{vmatrix}$$

**Theorem 1.1** Let  $\theta$  be the canonical anti-isomorphism of  $A_p$ . Then for  $n \geq 1$ ,

$$(P^{p_n-n})(x_1 x_2 \dots x_n)^{p-1} = w(n)^{p-1};$$

where  $p_n = (p^n - 1)/(p - 1)$ .

This result is true for  $p = 2$  if we interpret  $P^r$  as  $Sq^r$  [16]. The operation  $(P^{p_n-n})$  may be replaced by the admissible monomial  $P^{p^{n-1}-1} \dots P^{p^2-1} P^{p-1}$ , which is identical to the Milnor basis element  $P(p-1; \dots; p-1)$  of length  $n-1$  (see Proposition 3.2). In general the operation  $(P^{r_1} P^{r_2} \dots P^{r_m})$  used in Theorem 5.7 can not be replaced by an admissible monomial or a Milnor basis element.

The structure of the paper is as follows. Section 2 contains basic facts about the action of  $(P^r)$  and Milnor basis elements on polynomials. Section 3 contains independent proofs of Theorem 1.1 using invariant theory and by direct computation. In Section 4 we introduce the class of  $\mathbf{T}$ -regular partitions to which our main results apply, and extend Theorem 1.1 to  $\mathbf{T}^d$  for all  $d$ . The main results are stated in Section 5 and proved in Section 6. Section 7 relates these results to the  $\mathbb{F}_p[M_n]$ -module structure of  $\mathbf{P}$ . Section 8 gives Milnor basis elements which link the first occurrence and (in certain cases) the first submodule occurrence of a  $\mathbf{T}$ -regular representation of  $M_n$  with submodules in higher degrees.

The remarks which follow are intended to place our results in topological, combinatorial and algebraic contexts. As for topology, recall (e.g. [17]) that there is an  $A_p$ -module decomposition  $\mathbf{P} = \sum_i \mathbf{P}(i)$ , where the  $i$ -isotypical summand  $\mathbf{P}(i)$  is an indecomposable  $A_p$ -module, and where  $i = \dim L(i)$ , the dimension of  $L(i)$ . Identifying  $\mathbf{P}$  with the cohomology algebra  $H(\mathbb{C}P^1; \mathbb{F}_p)$ , this decomposition can be realized (after localization at  $p$ ) by a homotopy equivalence  $(\mathbb{C}P^1 \times \dots \times \mathbb{C}P^1) \xrightarrow{W} (i)Y$ , which splits the suspension of the product of  $n$  copies of infinite complex projective space  $\mathbb{C}P^1$  as a topological sum of spaces  $Y$  such that  $H(Y; \mathbb{F}_p) = \mathbf{P}(i)$ . The family of  $A_p$ -modules  $\mathbf{P}(i)$  is of major interest in algebraic topology. From this point of view, we determine the connectivity of  $Y$  for  $\mathbf{T}$ -regular (Corollary 5.8) and find a nonzero cohomology operation  $(i)$  on its bottom class (Theorem 5.7).

As for combinatorics and algebra, our aim is to provide information relating the  $A_p$ -module structure of  $\mathbf{P}(i)$  to combinatorial properties of  $i$  and representation theoretic properties of  $L(i)$ . The operation  $(i)$  and its source and target polynomials are combinatorially determined by  $i$ . The target polynomial is

defined by  $w(\lambda) = \prod_{j=1}^n w(\lambda_j)$ , where  $\lambda_j$  is the conjugate of  $\lambda$ , so that  $w(\lambda)$  is a product of determinants corresponding to the columns of the diagram of  $\lambda$ . This polynomial has already appeared in various forms in the literature. In Green's description [8, (5.4d)] of the highest weight vector of the dual Weyl module  $H^0(\lambda)$ ,  $w(\lambda)$  appears as a 'bideterminant' in the coordinate ring of  $M_n(K)$ , where  $K$  is an infinite field of characteristic  $p$ . A proof that  $w(\lambda)$  generates a submodule of  $\mathbf{P}^{ds(\lambda)}$  isomorphic to  $L(\lambda)$  was given in [7, Proposition 1.3], and a proof that this is the first occurrence of  $L(\lambda)$  as a submodule in  $\mathbf{P}$  was given in [12].

We would like to thank the referee of this paper for a very careful reading and for a number of helpful suggestions.

## 2 Preliminary results

In this section we use variants of the Cartan formula  $P^r(fg) = \sum_{r=s+t} P^s f P^t g$  to study the action on polynomials of the elements  $(P^r)$  and Milnor basis elements  $P(R)$  in the Steenrod algebra  $A_p$ . We begin with the standard formula

$$P^i(x^{p^b}) = \begin{cases} x^{p^{b+1}} & \text{if } i = p^b; \\ 0 & \text{otherwise for } i > 0; \end{cases} \tag{1}$$

In particular, we wish to evaluate Steenrod operations on Vandermonde determinants of the form

$$[x_{i_1}^{s_1}; x_{i_2}^{s_2}; \dots; x_{i_n}^{s_n}] = \begin{vmatrix} x_{i_1}^{s_1} & x_{i_2}^{s_1} & \dots & x_{i_n}^{s_1} \\ x_{i_1}^{s_2} & x_{i_2}^{s_2} & \dots & x_{i_n}^{s_2} \\ \vdots & \vdots & \ddots & \vdots \\ x_{i_1}^{s_n} & x_{i_2}^{s_n} & \dots & x_{i_n}^{s_n} \end{vmatrix};$$

where the exponents  $s_1; \dots; s_n$  are powers of  $p$ . As above, we shall abbreviate such determinants by listing their diagonal entries in square brackets: in particular,  $w(n) = [x_1; x_2^p; \dots; x_n^{p^{n-1}}]$ . As in Theorem 1.1, we write  $\rho_n = (p^n - 1)/(p - 1)$ , so that  $\rho_0 = 0$  and  $\rho_n - \rho_j = (p^n - p^j)/(p - 1)$ . The following result is a straightforward calculation using the Cartan formula and (1).

**Lemma 2.1** *If  $r = \rho_n - \rho_j$ ,  $0 \leq j < n$ , then*

$$P^r w(n) = [x_1; x_2^p; \dots; x_j^{p^{j-1}}; x_{j+1}^{p^{j+1}}; \dots; x_n^{p^n}];$$

and  $P^r w(n) = 0$  otherwise. In particular,  $P^r w(n) = 0$  for  $0 < r < \rho^{n-1}$ .  $\square$

To simplify signs, we usually write  $\beta^r$  for  $(-1)^r (P^r)$ . Thus if  $v$  is one of the generators  $x_i$  of  $\mathbf{P}$ , or more generally any linear form  $v = \sum_{i=1}^n a_i x_i$  in  $\mathbf{P}^1$ ,

$$\beta^r v = \begin{cases} v^{p^b} & \text{if } r = p^b, b \geq 0; \\ 0 & \text{otherwise;} \end{cases} \tag{2}$$

Formula (2) follows from (1) by using the identity  $\sum_{i+j=r} (-1)^i P^i P^j = 0$  in  $A_p$  and induction on  $r$ . Using the identity  $\sum_{i+j=r} (-1)^i P^i P^j = 0$  and induction on  $k$ , (2) can be generalized to

$$\beta^r x^{p^k} = \begin{cases} x^{p^b} & \text{if } r = p^b - p^k, b \geq k; \\ 0 & \text{otherwise;} \end{cases} \tag{3}$$

This leads to the following generalization of [16, Lemma 2.2].

**Lemma 2.2**

$$\beta^r [x_1^{p^k}; x_2^{p^{k+1}}; \dots; x_n^{p^{k+n-1}}] = \begin{cases} [x_1^{p^k}; \dots; x_{n-1}^{p^{k+n-2}}; x_n^{p^b}] & \text{if } r = p^b - p_{k+n-1}; \\ 0 & \text{otherwise;} \end{cases}$$

The modifications required to the proof given in [16] are straightforward. □

In evaluating the operations  $\beta^r$ , we shall frequently make use of the Cartan formula expansion for polynomials  $f, g \in \mathbf{P}$ :

$$\beta^r (fg) = \sum_{s+t=r} \beta^s f \beta^t g; \tag{4}$$

which holds because  $\beta$  is a coalgebra homomorphism.

**Lemma 2.3** For all polynomials  $f, g$  in  $\mathbf{P}$  and all  $r \geq 0$ ,

$$\beta^r (f^p g) = \sum_{r=ps+t} (\beta^s f)^p \beta^t g;$$

**Proof** By (4) it suffices to prove the case  $g = 1$ , i.e.

$$\beta^r f^p = \begin{cases} (\beta^s f)^p & \text{if } r = ps; \\ 0 & \text{if } r \text{ is not divisible by } p; \end{cases}$$

In this case, the Cartan formula (4) gives  $\beta^r f^p = \sum_{\substack{r \\ r_i \geq 0}} \beta^{r_1} f \dots \beta^{r_p} f$ , where the sum is over all ordered decompositions  $r = \sum_{i=1}^p r_i, r_i \geq 0$ . Except in the case where  $r_1 = \dots = r_p = s$ , cyclic permutation of  $r_1; \dots; r_p$  gives  $p$  equal terms which cancel in the sum. □

We write  $s(k)$  for the sum of the digits in the base  $p$  expansion of a positive integer  $k$ , i.e. if  $k = \sum_{i=0}^{\infty} a_i p^i$  where  $0 \leq a_i < p$ , then  $s(k) = \sum_{i=0}^{\infty} a_i$ . Thus  $s(k)$  is the minimum number of powers of  $p$  which have sum  $k$ , and  $s(k) \equiv k \pmod{p-1}$ . Formula (2) leads to the following simple sufficient condition for the vanishing of  $\mathfrak{p}^r$  on a homogeneous polynomial of degree  $d$ .

**Lemma 2.4** *If  $(r(\rho - 1) + d) > d$ , then  $\mathfrak{p}^r f = 0$  for all  $f \in \mathbf{P}^d$ .*

**Proof** Since the action of  $\mathfrak{p}^r$  is linear and commutes with specialization of the variables, it is sufficient to prove this when  $f = x_1 x_2 \cdots x_d$ . By (4)  $\mathfrak{p}^r f = \sum \mathfrak{p}^{r_1} x_1 \mathfrak{p}^{r_2} x_2 \cdots \mathfrak{p}^{r_d} x_d$ , where the sum is over all ordered decompositions  $r = r_1 + r_2 + \cdots + r_d$  with  $r_1, r_2, \dots, r_d \geq 0$ . By (2), the only non-zero terms are those in which  $r_i = p k_i$  for some non-negative integers  $k_1, k_2, \dots, k_d$ . But then  $r(\rho - 1) + d = \sum p k_i$ , and the result follows by definition.  $\square$

**Lemma 2.5** *Let  $k \geq 0$  and let  $v = \sum_{i=1}^n a_i x_i$  be a linear form in  $\mathbf{P}^1$ . Then*

$$\mathfrak{p}^{p^k - 1} v^{p-1} = v^{p^k(p-1)};$$

**Proof** There is a unique way to write  $p^k - 1$  as the sum of  $p - 1$  integers of the form  $p_i$  for  $i \geq 0$ , namely  $p^k - 1 = (p - 1)p_k$ . The result now follows from (2) and the Cartan formula (4).  $\square$

**Remark 2.6** The same method can be used to evaluate  $\mathfrak{p}^r v^{p-1}$  for all  $r$ . The result is

$$\mathfrak{p}^r v^{p-1} = \begin{cases} c_r v^{(r+1)(p-1)} & \text{if } ((r + 1)(p - 1)) = p - 1; \\ 0 & \text{otherwise;} \end{cases}$$

where if  $(r + 1)(p - 1) = j_1 p^{a_1} + \cdots + j_s p^{a_s}$ , with  $a_1 > \cdots > a_s \geq 0$  and  $\sum_{i=1}^s j_i = p - 1$ , then  $c_r = (p - 1)! / (j_1! j_2! \cdots j_s!)$ .

The following result, the ‘Cartan formula for Milnor basis elements’ is well-known (cf. [16, Lemma 5.3]).

**Lemma 2.7** *For a Milnor basis element  $P(R) = P(r_1; \dots; r_n)$  and polynomials  $f, g \in \mathbf{P}$ ,*

$$P(R)(fg) = \sum_{R=S+T} P(S)f P(T)g;$$

where the sum is over all sequences  $S = (s_1; \dots; s_n)$  and  $T = (t_1; \dots; t_n)$  of non-negative integers such that  $r_i = s_i + t_i$  for  $1 \leq i \leq n$ .  $\square$

In the same way as for Lemma 2.3, this gives the following result.

**Lemma 2.8** *Let  $P(R) = P(r_1; \dots; r_n)$  be a Milnor basis element and let  $f, g \in \mathbf{P}$  be polynomials. Then*

$$P(R)(f^p g) = \sum_{R=\rho S+T} (P(S)f)^\rho P(T)g; \quad \square$$

Here  $R = \rho S + T$  means that  $r_i = \rho s_i + t_i$  for  $1 \leq i \leq n$ .

### 3 The $\det^{\rho-1}$ representation

In this section we give three proofs of Theorem 1.1. The first uses the results of [12] on submodules, while the second is a variant of this which uses only classical invariant theory. The third proof is computational. The first two proofs use the following preliminary result, which shows that the operation  $\mathfrak{b}^{\rho n-n}$  maps to 0 all monomials of degree  $n(\rho - 1)$  other than the generating monomial  $(x_1 x_2 \dots x_n)^{\rho-1}$  for  $\det^{\rho-1}$ .

**Lemma 3.1** *Let  $f$  be a polynomial in  $\mathbf{P}^{n(\rho-1)}$  which is divisible by  $x^\rho$  for some variable  $x = x_i, 1 \leq i \leq n$ . Then  $\mathfrak{b}^{\rho n-n} f = 0$ .*

**Proof** Let  $f = x^\rho g$ , where  $g \in \mathbf{P}$ . Then by Lemma 2.3

$$\mathfrak{b}^{\rho n-n} f = \sum_{\rho n-n=\rho s+t} (\mathfrak{b}^s x)^\rho \mathfrak{b}^t g; \quad (5)$$

By (2),  $\mathfrak{b}^s x = 0$  if  $s \notin \rho_k$  for some  $k$  with  $0 \leq k \leq n-2$ . Thus it is sufficient to prove that  $\mathfrak{b}^t g = 0$  for  $t = \rho n - n - \rho \rho_k$ , where  $g \in \mathbf{P}^{n(\rho-1)-\rho}$ . By Lemma 2.4, this holds when  $((t+n)(\rho-1) - \rho) > n(\rho-1) - \rho$ . Now  $(t+n)(\rho-1) - \rho = \rho n(\rho-1) - \rho \rho_k(\rho-1) - \rho = \rho^n - \rho^{k+1} - 1$ , hence  $((t+n)(\rho-1) - \rho) = n(\rho-1) - 1 > n(\rho-1) - \rho$  as required. Thus  $\mathfrak{b}^t g = 0$  in all terms of (5) in which  $\mathfrak{b}^s x \neq 0$ , and so  $\mathfrak{b}^{\rho n-n} f = 0$ .  $\square$

**First Proof of Theorem 1.1** We first show that the monomial  $m = (x_1 x_2^\rho x_n^{\rho n-1})^{\rho-1}$  appears in  $\mathfrak{b}^{\rho n-n}(x_1 \dots x_n)^{\rho-1}$  with coefficient 1. In the Cartan formula expansion (4),  $m$  can appear only in the term arising from the decomposition  $\rho n - n = r_1 + r_2 + \dots + r_n$ , where  $r_k = \rho^{k-1} - 1$  for  $1 \leq k \leq n$ . By Lemma 2.5,  $m$  appears in this term with coefficient 1.

By Lemma 3.1,  $\mathfrak{p}^{\rho n - n}$  maps all other monomials in degree  $n(\rho - 1)$  to 0. Hence  $\mathfrak{p}^{\rho n - n}(x_1 \dots x_n)^{\rho - 1}$  generates a 1-dimensional  $\mathbb{F}_\rho[M_n]$ -submodule of  $\mathbf{P}^{\rho^n - 1}$ . Since  $(x_1 \dots x_n)^{\rho - 1}$  generates the 1-dimensional quotient  $\mathbf{T}^{n(\rho - 1)}$  of  $\mathbf{P}^{n(\rho - 1)}$  and since  $\mathbf{T}^{n(\rho - 1)} = \det^{\rho - 1}$ , this submodule of  $\mathbf{P}^{\rho^n - 1}$  is also isomorphic to  $\det^{\rho - 1}$ .

It is known [12] that the first submodule occurrence of  $\det^{\rho - 1}$  for  $M_n$  in  $\mathbf{P}$  is generated by  $w(n)^{\rho - 1}$ , and that this is the unique submodule occurrence of  $\det^{\rho - 1}$  in degree  $\rho^n - 1$ . Since  $m$  is the product of the leading diagonal terms in  $w(n)^{\rho - 1} = [x_1; x_2^{\rho}; \dots; x_n^{\rho^{n-1}}]^{\rho - 1}$ ,  $m$  also has coefficient 1 in  $w(n)^{\rho - 1}$ .  $\square$

**Second Proof of Theorem 1.1** We recall that  $D(n; \rho)$  is the ring of  $GL_n(\mathbb{F}_\rho)$ -invariants in  $\mathbf{P}$ , and that it is a polynomial algebra over  $\mathbb{F}_\rho$  with generators  $Q_{n,i}$  in degree  $\rho^n - \rho^i$  for  $0 \leq i \leq n - 1$ . We may identify  $Q_{n,0}$  with  $w(n)^{\rho - 1}$ . Since  $\mathbf{T}^{n(\rho - 1)}$  is isomorphic to the trivial  $GL_n(\mathbb{F}_\rho)$ -module, it follows as in our first proof that  $\mathfrak{p}^{\rho n - n}(x_1 \dots x_n)^{\rho - 1} \in D(n; \rho)$ .

We shall prove that  $w(n)$  divides  $\mathfrak{p}^{\rho n - n}(x_1 \dots x_n)^{\rho - 1}$ . Recall that  $w(n)$  is the product of linear factors  $c_1 x_1 + \dots + c_n x_n$ , where  $c_1, \dots, c_n \in \mathbb{F}_\rho$ . If we specialize the variables in  $(x_1 \dots x_n)^{\rho - 1}$  by imposing the relation  $c_1 x_1 + \dots + c_n x_n = 0$ , then every monomial in the resulting polynomial is divisible by  $x^{\rho}$  for some variable  $x = x_i$ . By Lemma 3.1, such a monomial is in the kernel of  $\mathfrak{p}^{\rho n - n}$ . Thus  $\mathfrak{p}^{\rho n - n}(x_1 \dots x_n)^{\rho - 1}$  is divisible by  $c_1 x_1 + \dots + c_n x_n$ , and so it is divisible by  $w(n)$ .

Now an element of  $D(n; \rho)$  in degree  $\rho^n - 1$  which is divisible by  $w(n)$  must be a scalar multiple of  $Q_{n,0} = w(n)^{\rho - 1}$ . For if a polynomial in the remaining generators  $Q_{n,1}, \dots, Q_{n,n-1}$  of  $D(n; \rho)$  is divisible by  $w(n)$ , the quotient would be  $SL_n(\mathbb{F}_\rho)$ -invariant, giving a non-trivial polynomial relation between  $Q_{n,1}, \dots, Q_{n,n-1}$  and  $w(n)$ . This contradicts Dickson's theorem that these are algebraically independent generators of the polynomial algebra of  $SL_n(\mathbb{F}_\rho)$ -invariants in  $\mathbf{P}$ .  $\square$

Our third proof of Theorem 1.1 is by direct calculation. We shall evaluate the Milnor basis element  $P(\rho - 1; \dots; \rho - 1)$  of length  $n - 1$  on  $(x_1 \dots x_n)^{\rho - 1}$ . The following result relates the element  $P(\rho - 1; \dots; \rho - 1; b)$  of length  $n$  to admissible monomials and to the anti-automorphism  $\sigma$ . In particular, we show that  $P(\rho - 1; \dots; \rho - 1)$  and  $\mathfrak{p}^{\rho n - n}$  have the same action on  $(x_1 \dots x_n)^{\rho - 1}$ .

**Proposition 3.2** For  $1 \leq b \leq \rho - 1$ ,

$$(i) \quad P(\rho - 1; \dots; \rho - 1; b) = P^{(b+1)\rho^{n-1} - 1} P^{(b+1)\rho - 1} P^b \text{ for } n \geq 1,$$



- (ii)  $\beta^{(b+1)p_n-n}g = P(p-1; \dots; p-1; b)g$  if  $\deg g = n(p-1) + b$  for  $n \geq 1$ ,
- (iii)  $\beta^{(b+1)p_n-n} = \rho^{(b+1)p^{n-1}} \beta^{(b+1)p_{n-1}-n} + P(p-1; \dots; p-1; b)$  for  $n \geq 2$ .

**Proof** Statement (i) is a special case of [4, Theorem 1.1]. For (ii), recall [11] that  $\beta^d$  is the sum of all Milnor basis elements  $P(R)$  in degree  $d(p-1)$ . Here  $R = (r_1; r_2; \dots; r_m)$  is a finite sequence of non-negative integers, and  $P(R)$  has degree  $jRj = \sum (p^j - 1)r_j$  and excess  $e(R) = \sum r_j$ . In particular,  $P^d = P(d)$  is the unique Milnor basis element of maximum excess  $d$  in degree  $d(p-1)$ , but in general there may be more than one element of minimum excess in a given degree.

We will show that  $P(p-1; \dots; p-1; b)$  is the unique element of minimum excess  $e = (n-1)(p-1) + b$  in degree  $d(p-1)$  when  $d = (b+1)p_n - n$ . By [11, Lemma 8] a bijection  $P(r_1; r_2; \dots; r_m) \xrightarrow{P^{t_1} P^{t_2} \dots P^{t_m}}$  between the Milnor basis and the admissible basis of  $A_p$  is defined by  $t_m = r_m$  and  $t_i = r_i + p t_{i+1}$  for  $1 \leq i < m$ . This preserves both the degree and the excess. Thus it is equivalent to prove that  $m = \rho^{(b+1)p^{n-1}-1} \rho^{(b+1)p-1} \rho^b$  is the unique admissible monomial of minimum excess in degree  $d(p-1)$ . Now the excess of an admissible monomial  $P^{t_1} P^{t_2} \dots P^{t_m}$  is  $pt_1 - d(p-1)$  where  $d = \sum_i t_i$ , and so it is minimal when  $t_1$  is minimal. It is easy to verify that  $m$  is the unique admissible monomial in degree  $d(p-1)$  for which  $t_1 = (b+1)p^{n-1} - 1$ , and that this value of  $t_1$  is minimal.

Note that  $p$  divides  $jRj + e(R)$  for all  $R$ . Hence  $\beta^{(b+1)p_n-n} - P(p-1; \dots; p-1; b)$  has excess  $> e + p - 1 = n(p-1) + b$ , and so  $\beta^{(b+1)p_n-n}g = P(p-1; \dots; p-1; b)g$  when  $g$  is a polynomial of degree  $= n(p-1) + b$ .

(iii) Recall Davis's formula [5]

$$\rho^u \beta^v = \sum_{jRj = (p-1)(u+v)} \times \frac{jRj + e(R)}{\rho^u} P(R); \tag{6}$$

which we may apply in the case  $u = (b+1)p^{n-1}$ ,  $v = (b+1)p_{n-1} - n$  to show that  $\rho^u \beta^v$  is the sum of all Milnor basis elements in degree  $d(p-1)$  other than the element  $P(p-1; \dots; p-1; b)$  of minimal excess.

For  $R = (p-1; \dots; p-1; b)$  we have  $jRj + e(R) = (b+1)p^n - p$ , and since  $\rho^u = (b+1)p^n$  the coefficient in (6) is zero. Since  $p$  divides  $jRj + e(R)$  for all  $R$ ,  $jRj + e(R) = (b+1)p^n$  for all other  $R$  with  $jRj = d(p-1)$ . As remarked above, the unique element of maximal excess is  $P^d$  itself, and so for all  $R$  we have  $jRj + e(R) = pd = (b+1)(p + p^2 + \dots + p^n) - pn$ . It is clear from this inequality that the coefficient in (6) is 1 for all  $R \neq (p-1; \dots; p-1; b)$ .  $\square$

**Third Proof of Theorem 1.1** Let  $w_n = p^{p^n - 1} - p^{p^2 - 1} p^{p - 1}$  for  $n \geq 1$ , and  $w_0 = 1$ . We assume that  $w_{n-1}(x_1 \dots x_n)^{p-1} = w(n)^{p-1}$  as induction hypothesis on  $n$ , the case  $n = 1$  being trivial.

The cofactor expansion of  $w(n+1) = [x_1; x_2^p; \dots; x_{n+1}^{p^n}]$  by the top row gives  $w(n+1) = \sum_{i=1}^{n+1} (-1)^i x_i^p \det_i$ , where  $\det_i = [x_1; \dots; x_{i-1}^{p^{i-2}}; x_{i+1}^{p^{i-1}}; \dots; x_{n+1}^{p^{n-1}}]$ . Hence  $w(n+1) (x_1 \dots x_{n+1})^{p-1} = \sum_{i=1}^{n+1} (-1)^i x_i^p \det_i (x_1 \dots x_{i-1} x_{i+1} \dots x_{n+1})^{p-1}$ .

By Proposition 3.2(i),  $\det_i = p \binom{p-1}{i-1} \det_i$  of length  $n$ , and so by Lemma 2.8  $w_n(w(n+1) (x_1 \dots x_{n+1})^{p-1}) = \sum_{i=1}^{n+1} (-1)^i x_i^p \det_i w_n(x_1 \dots x_{i-1} x_{i+1} \dots x_{n+1})^{p-1}$ . Since  $w_n = p^{p^n - 1} w_{n-1}$ ,  $w_n(x_1 \dots x_{i-1} x_{i+1} \dots x_{n+1})^{p-1} = p^{p^n - 1} \det_i^{p-1}$  by the induction hypothesis. Since  $\det_i^{p-1}$  has degree  $p^n - 1$ ,  $p^{p^n - 1} \det_i^{p-1} = \det_i^{p(p-1)}$ . Hence  $w_n(w(n+1) (x_1 \dots x_{n+1})^{p-1}) = \sum_{i=1}^{n+1} (-1)^i x_i^p \det_i^{p^2} = w(n+1)^p$ .

By Lemma 2.1,  $P^r w(n+1) = 0$  for  $0 < r < p^n$ . As  $w_n = p^{p^n - 1} - p^{p^2 - 1} p^{p - 1}$ , iterated application of the Cartan formula gives  $w_n(w(n+1) (x_1 \dots x_{n+1})^{p-1}) = w(n+1) w_n(x_1 \dots x_{n+1})^{p-1}$ . Hence  $w(n+1) w_n(x_1 \dots x_{n+1})^{p-1} = w(n+1)^p$ . Cancelling the factor  $w(n+1)$ , the inductive step is proved.

### 4 T-regular partitions

In this section we define the special class of **T-regular** partitions, and extend Theorem 1.1 to give a Steenrod operation  $\beta^r$  which links the first occurrence and first submodule occurrence of  $\mathbf{T}^d$  for all  $d$ . In fact we prove a more general result which links the first occurrence to a family of higher degree occurrences.

The truncated polynomial module  $\mathbf{T}^d = \mathbf{P}^d / (\mathbf{P}^d \setminus (x_1^p; \dots; x_n^p))$  has a  $\mathbb{F}_p$ -basis represented in  $\mathbf{P}^d$  by the set of all monomials  $x_1^{s_1} x_2^{s_2} \dots x_n^{s_n}$  of total degree  $d = \sum_i s_i$  with  $s_i < p$  for  $1 \leq i \leq n$ . By [2, Theorem 6.1]  $\mathbf{T}^d = L((p-1)^{n-1} b)$ , where  $d = (n-1)(p-1) + b$  and  $1 \leq b \leq p-1$ . We regard the corresponding diagram as a block of  $p-1$  columns, in which the first  $b$  columns have length  $n$  and the remaining  $p-b-1$  columns have length  $n-1$ . Given a partition  $\lambda$ , we can divide its diagram into  $m$  blocks of  $p-1$  columns and compare the blocks with the diagrams corresponding to these. (The  $m$ th block may have  $< p-1$  columns.) For  $1 \leq j \leq m$ , let  $\lambda^{(j)}$  be the partition whose diagram is the  $j$ th block, and let  $\ell_j = \deg \lambda^{(j)}$  be the number of boxes in the  $j$ th block.

**Definition 4.1** A column  $p$ -regular partition  $\lambda$  is **T-regular** if  $L(\lambda^{(j)}) = \mathbf{T}^j$  for all  $j$ . Equivalently, for all  $a \geq 1$ , there is at most one value of  $i$  for which  $(a-1)(p-1) < \ell_i < a(p-1)$ . If  $\lambda$  is **T-regular**, we call  $\lambda$  the **T-conjugate** of  $\dots$ .

In the case  $p = 2$ , all column 2-regular partitions are  $\mathbf{T}$ -regular, and  $\bar{\lambda} = \lambda'$ , the conjugate of  $\lambda$ . If  $\lambda$  is column 2-regular, then the partition  $\lambda' = (\rho - 1)$  obtained by multiplying each part of  $\lambda$  by  $\rho - 1$  is  $\mathbf{T}$ -regular. Since  $\lambda$  is column  $\rho$ -regular,  $\lambda_j - \lambda_{j+1} \leq \rho - 1$  for all  $j$ , and  $m \leq n$ . Thus there is a bijection  $\mathcal{S}$  between the set of  $\mathbf{T}$ -regular partitions  $\lambda = (\lambda_1; \dots; \lambda_n)$  and the set of partitions  $\mu = (\mu_1; \dots; \mu_n)$  which satisfy  $\mu_1 \leq n(\rho - 1)$  and  $\mu_j - \mu_{j+1} \leq \rho - 1$  for  $1 \leq j \leq n - 1$ . In terms of the Mullineux involution  $M$  on the set of all row  $\rho$ -regular partitions,  $\lambda$  and  $\mu$  are related by  $M(\lambda) = \mu'$  [15, Proposition 3.13].

We next extend Theorem 1.1 to give linking formulae for the representations  $\mathbf{T}^d$ . It will be convenient to introduce abbreviated notation for some further Vandermonde determinants. Let  $w(n; a) = [x_1; \dots; x_a^{p^a-1}; x_{a+1}^{p^{a+1}}; \dots; x_n^{p^n}]$  for  $0 \leq a \leq n$ , where the exponent  $p^a$  is omitted. In particular,  $w(n; n) = w(n)$  and  $w(n; 0) = w(n)^p$ .

**Proposition 4.2** For  $n \geq 1$  and  $1 \leq i \leq p - 1$ , let  $i = i_1 + \dots + i_s$  where  $i_1; \dots; i_s > 0$ , and let  $j = i_1 p_{a_1} + \dots + i_s p_{a_s}$ , where  $a_1 > \dots > a_s \geq 0$ . Then

$$p^{p_{n-n-j}} (x_1 x_2 \dots x_{n-1})^{p-1} x_n^{p-i-1} = (-1)^{i(n-1)-j} w(n)^{p-i-1} \prod_{r=1}^s w(n-1; a_r)^{i_r}.$$

Specializing to the case  $s = 1$ ,  $j = i p_{n-1}$  and putting  $b = p - 1 - i$ , we obtain an operation linking the first occurrence and the first submodule occurrence of the representation  $\mathbf{T}^d$ , as follows. Theorem 1.1 can be taken as the case  $b = 0$  or as the case  $b = p - 1$ ; we choose  $b = p - 1$  to fit notation later.

**Corollary 4.3** For  $n \geq 1$  and  $1 \leq b \leq p - 1$ ,

$$p^{(b+1)p_{n-1}-(n-1)} (x_1 x_2 \dots x_{n-1})^{p-1} x_n^b = w(n)^b w(n-1)^{p-b-1}.$$

**Proof of Proposition 4.2** We introduce a parameter into Theorem 1.1, by working in  $\mathbb{F}_p[x_1; \dots; x_{n+1}]$  and writing  $x_{n+1} = t$  in order to distinguish this variable. Since the action of  $A_p$  commutes with the linear substitution which maps  $x_n$  to  $x_n + t$  and fixes  $x_i$  for  $i \neq n$ , we obtain

$$p^{p_{n-n}} (x_1 \dots x_{n-1} (x_n + t))^{p-1} = [x_1; x_2^p; \dots; x_{n-1}^{p^{n-2}}; (x_n + t)^{p^{n-1}}]^{p-1}. \tag{7}$$

Expanding the left hand side of (7) by the binomial theorem, we obtain

$$\sum_{i=0}^{p-1} (-1)^i p^{p_{n-n}} ((x_1 \dots x_{n-1})^{p-1} x_n^{p-1-i} t^i).$$

The right hand side of (7) is

$$[x_1; x_2^{\rho}; \dots; x_{n-1}^{\rho^{n-2}}; x_n^{\rho^{n-1}} + t^{\rho^{n-1}}]^{\rho-1} = \sum_{i=0}^{\rho-1} (-1)^i w(n)^{\rho-1-i} [x_1; x_2^{\rho}; \dots; x_{n-1}^{\rho^{n-2}}; t^{\rho^{n-1}}]^i;$$

since  $w(n) = [x_1; x_2^{\rho}; \dots; x_n^{\rho^{n-1}}]$ . The summands in (7) corresponding to  $i = 0$  give the original result, Theorem 1.1, and so are equal. In fact we can equate the  $i$ th summands for all  $i$ . This happens because  $\rho^r$  raises degree by  $r(\rho - 1)$ , so that the powers  $t^k$  which occur in the  $i$ th summand on the left have  $k \equiv i \pmod{\rho - 1}$ , while if  $t^k$  occurs in the  $i$ th summand on the right, then  $k$  is the sum of  $i$  powers of  $\rho$ , so that again  $k \equiv i \pmod{\rho - 1}$ . Hence for  $1 \leq i \leq \rho - 1$  we have

$$\rho^{\rho n - n} ((x_1 \dots x_{n-1})^{\rho-1} x_n^{\rho-1-i} t^i) = w(n)^{\rho-1-i} [x_1; x_2^{\rho}; \dots; x_{n-1}^{\rho^{n-2}}; t^{\rho^{n-1}}]^i. \tag{8}$$

Since the powers  $t^k$  of  $t$  which can appear here are such that  $k$  is the sum of  $i$  powers of  $\rho$ , we can write  $k = i_1 \rho^{a_1} + \dots + i_s \rho^{a_s}$ , where  $a_1 > \dots > a_s \geq 0$  and  $i_1 + \dots + i_s = i$ . Using the expansion

$$[x_1; x_2^{\rho}; \dots; x_{n-1}^{\rho^{n-2}}; t^{\rho^{n-1}}] = \sum_{a=0}^{\rho-1} (-1)^{n-1-a} w(n-1; a) t^{\rho^a}$$

we can evaluate the coefficient of  $t^k$  on the right hand side of (8) as

$$(-1)^{i(n-1)-j} \frac{i!}{i_1! \dots i_s!} w(n)^{\rho-1-i} w(n-1; a_1)^{i_1} \dots w(n-1; a_s)^{i_s};$$

where we have simplified the sign by noting that  $a_1 i_1 + \dots + a_s i_s \equiv j \pmod{2}$  since  $\rho_{a_s} \equiv a \pmod{2}$ . By the Cartan formula (4), the left hand side of (8) is

$$\sum_{j=0}^{\rho n - n} \rho^{\rho n - n - j} (x_1 \dots x_{n-1})^{\rho-1} x_n^{\rho-1-i} \rho^j t^i$$

Here the term in  $t^k$  arises from  $\rho^j t^i$  where  $k = j(\rho - 1) + i$ , so that  $j = i_1 \rho_{a_1} + \dots + i_s \rho_{a_s}$ , and since this decomposition of  $j$  as a sum of at most  $i$  powers of  $\rho$  is unique, formulas (2) and (4) give  $\rho^j t^i = (i! = i_1! \dots i_s!) t^k$ . Thus equating coefficients of  $t^k$  in (8) gives the result.

### 5 Linking for T-regular representations

In this section we state our main results. We fix an odd prime  $\rho$  and a positive integer  $n$  throughout. As in [16], our results will be statements about polynomials in  $n$  variables when  $\mathcal{X}$  has length  $n$ , i.e.  $\mathcal{X}$  has  $n$  nonzero parts. There

is no loss of generality, since the projection in  $M_n$  which sends  $x_n$  to 0 and  $x_i$  to  $x_i$  for  $i < n$  maps  $L(\lambda)$  to zero if  $\lambda_n > 0$  and on to the corresponding  $\mathbb{F}_p[M_{n-1}]$ -module  $L(\lambda)$  if  $\lambda_n = 0$  (cf. [2, Section 3]). Hence we shall always assume that  $\lambda_n \notin 0$ .

We first establish some notation. Given a  $\mathbf{T}$ -regular partition  $\lambda$  of length  $n$ , we define a polynomial  $v(\lambda)$  whose degree  $d_c(\lambda)$  is given by (9) and which ‘represents’  $L(\lambda)$ , in the sense that the submodule of  $\mathbf{P}^{d_c(\lambda)}$  generated by  $v(\lambda)$  has a quotient module isomorphic to  $L(\lambda)$ . We index the diagram of  $\lambda$  using matrix coordinates  $(i;j)$ , so that  $1 \leq i \leq n$  and  $1 \leq j \leq \lambda_i$ .

**Definition 5.1** *The  $k$ th antidiagonal of the diagram of  $\lambda$  is the set of boxes such that  $j + i(\rho - 1) = k + \rho - 1$ . If the lowest box is in row  $i$  and the highest is in row  $i - s + 1$ , let  $v_k(\lambda) = [x_{i-s+1}; x_{i-s+2}^{\rho}; \dots; x_i^{\rho^{s-1}}]$ , and let  $v(\lambda) = \bigoplus_{k=1}^1 v_k(\lambda)$ .*

Thus an antidiagonal is the set of boxes which lie on a line of slope  $1=(\rho - 1)$  in the diagram, and  $v(\lambda)$  is a product of corresponding Vandermonde determinants. Indenting successive rows by  $\rho - 1$  columns, we obtain a shifted diagram whose columns correspond to these antidiagonals. The  $\mathbf{T}$ -conjugate  $\lambda'$  of  $\lambda$  records the number of antidiagonals  $\lambda'_s$  of length  $\lambda'_s$  for all  $s \geq 1$ .

**Example 5.2** Let  $\rho = 5$ ,  $\lambda = (9;6;3)$ , so that  $\lambda' = (11;6;1)$ . The shifted diagram

$$\text{gives } v(\lambda) = x_1^4 [x_1; x_2^5]^4 [x_1; x_2^5; x_3^{25}] [x_2; x_3^5] x_3.$$

Recall [12] that  $w(\lambda) = \bigoplus_{j=1}^1 w(\lambda_j)$  generates the first occurrence of  $L(\lambda)$  as a submodule in  $\mathbf{P}$ . Thus we can rewrite the linking theorem for  $\mathbf{T}^d$ , Corollary 4.3, as follows.

**Theorem 5.3** *Let  $d = (n - 1)(\rho - 1) + b$ , where  $n \geq 1$  and  $1 \leq b \leq \rho - 1$ , so that  $\mathbf{T}^d = L(\lambda)$  where  $\lambda = ((\rho - 1)^{n-1}b)$ . Then  $\mathbf{p}^r v(\lambda) = w(\lambda)$ , where  $r = (b + 1)\rho_{n-1} - (n - 1)$  and  $\rho_{n-1} = (\rho^{n-1} - 1)/(\rho - 1)$ .  $\square$*

By the *leading monomial* of a polynomial we mean the monomial  $\bigoplus_{i=1}^n x_i^{s_i}$  occurring in it (ignoring the nonzero coefficient) whose exponents are highest in left lexicographic order. The leading monomial  $s(\lambda)$  of  $v(\lambda)$  is obtained by

multiplying the principal antidiagonals in the determinants  $v_k(\lambda)$ ,  $1 \leq k \leq n-1$ . (In Example 5.2,  $s(\lambda) = x_1^{49}x_2^{14}x_3^3$ .) The base  $p$  expansion of every exponent in  $s(\lambda)$  has the form  $s_i = c_k p^k + (p-1)p^{k-1} + \dots + (p-1)p + (p-1)$ , i.e.  $s_i \equiv -1 \pmod{p^k}$ , where  $p^k < s_i < p^{k+1}$ . We adapt the terminology introduced by Singer [13], by calling such a monomial a ‘spike’. In the case  $p = 2$ ,  $s(\lambda) = x_1^{2^{i-1}-1} \dots x_n^{2^n-1}$ . A polynomial which contains such a spike can not be ‘hit’, i.e. it can not be the image of a polynomial of lower degree under a Steenrod operation. This is easily seen by considering the 1-variable case. Hence the polynomial  $v(\lambda)$  is not hit.

**Proposition 5.4** *Let  $\lambda$  be  $\mathbf{T}$ -regular with  $\mathbf{T}$ -conjugate  $\mu$ .*

- (i) *If  $\lambda_i = a_i(p-1) + b_i$ ,  $a_i \in \{0, 1\}$ ,  $b_i \in \{0, \dots, p-1\}$ , then  $s(\lambda) = \prod_{i=1}^n X_i^{(b_i+1)p^{a_i}-1}$ .*
- (ii) *With  $\mu^{(j)}$  as in Definition 4.1,  $s(\lambda) = v(\mu^{(1)}) v(\mu^{(2)})^p \dots v(\mu^{(m)})^{p^{m-1}}$ .*
- (iii) *The coefficient of  $s(\lambda)$  in  $v(\lambda)$  is  $(-1)^{\ell(\lambda)}$ , with  $\ell(\lambda) = \prod_{j=1}^{[m-2]} (-1)^{j-1} 2^j$ .*

**Proof** Formulae (i) and (ii) are easily read off from a tableau obtained by entering  $p^{j-1}$  in each box in the  $j$ th block of  $p-1$  columns of the diagram of  $\lambda$ , and reading this according to rows and to blocks of columns. For (iii), note that the sign of the term arising from the leading antidiagonal in the expansion of an  $s \times s$  determinant is  $+1$  for  $s \equiv 0, 1 \pmod{4}$  and  $-1$  for  $s \equiv 2, 3 \pmod{4}$ , and that the diagram of  $\lambda$  has  $j$  antidiagonals of length  $2^j$ . □

In Theorem 5.5 we establish (i) a ‘level 0 formula’, which gives a sufficient condition for  $\mathfrak{p}^r v(\lambda) = 0$ , and (ii) a ‘level 1 formula’, which gives a sufficient condition for  $\mathfrak{p}^r v(\lambda)$  to be a product related to the decomposition  $\lambda = \mu^{(1)} + \mu^{(-)}$  which splits off the first  $p-1$  columns of the diagram. Thus  $\mu^{(1)} = ((p-1)^{n-1}b)$ , where  $\mu_1 = (n-1)(p-1) + b$  and  $1 \leq b \leq p-1$ , and  $\mu^{(-)}$  is defined by  $\mu_i^- = \mu_i - (p-1)$  if  $\mu_i \geq p-1$ , and  $\mu_i^- = 0$  otherwise. Our main linking result, Theorem 5.7, follows from Theorem 5.5 by induction on  $m$ , the length of  $\lambda$ . The proofs of Theorems 5.5 and 5.7 are deferred to Section 6.

**Theorem 5.5** *Let  $\lambda$  be  $\mathbf{T}$ -regular with  $\mathbf{T}$ -conjugate  $\mu$ , let  $d_c$  be defined by (9) below, and let  $R(r; \lambda) = r(p-1) + d_c(\lambda) - d_c(\mu^{(-)})$ . Recall that  $\ell(k)$  is the sum of the digits in the base  $p$  expansion of  $k$ .*

- (i) *If  $\ell(R(r; \lambda)) > 1$ , then  $\mathfrak{p}^r v(\lambda) = 0$ .*
- (ii) *If  $\ell(R(r; \lambda)) = 1$ , then  $\mathfrak{p}^r v(\lambda) = \mathfrak{p}^{r+d_c(\mu^{(-)})} v(\mu^{(1)}) v(\mu^{(-)})$ .*

**Remark 5.6** Taking  $p = 2$  and  $P^r = Sq^r$ , this reduces to [16, Theorem 2.1], since that theorem can be applied to  $\lambda_{(1)} = (1^n)$  to obtain  $Sq^{r+d_c(\lambda_{(1)})} v(\lambda_{(1)}) = [x_1^{2^{a_1}}; \dots; x_n^{2^{a_n}}]$ , where  $a_1 < \dots < a_n$ . The hypothesis on  $r$  is satisfied since  $r + d_c(\lambda_{(1)}) + n = r + d_c(\lambda_{(1)}) - d_c(\lambda_{(1)}) = 2^{a_1} + \dots + 2^{a_n}$ .

Combining Theorem 5.3 with Theorem 5.5, we obtain our main theorem.

**Theorem 5.7** Let  $\lambda$  be  $\mathbf{T}$ -regular with  $\mathbf{T}$ -conjugate  $\mu$  of length  $m$ . For  $1 \leq k \leq m$ , let  $r_k = (n_k - 1)(p - 1) + b_k$ , where  $n_k \geq 1$  and  $1 \leq b_k \leq p - 1$ . Then

$$\beta^{r_m} \beta^{r_2} \beta^{r_1} v(\lambda) = w(\mu);$$

where  $r_k = (b_k + 1)p_{n_k-1} - (n_k - 1) - \prod_{j=k+1}^m p^{j-k-1}$ .

This theorem determines the first occurrence degree  $d_c(\lambda)$  when  $\lambda$  is  $\mathbf{T}$ -regular.

**Corollary 5.8** Let  $\lambda$  be  $\mathbf{T}$ -regular with  $\mathbf{T}$ -conjugate  $\mu$ . Then the degree in which the irreducible module  $L(\lambda)$  first occurs as a composition factor in the polynomial algebra  $\mathbf{P}$  is given by

$$d_c(\lambda) = \sum_{i=1}^n p^{i-1} \mu_i; \tag{9}$$

and the  $\mathbb{F}_p[M_n]$ -submodule of  $\mathbf{P}^{d_c(\lambda)}$  generated by  $v(\lambda)$  has a quotient module isomorphic to  $L(\lambda)$ .

**Proof** By [7] or [12]  $w(\mu)$  generates a submodule of  $\mathbf{P}^{d_c(\lambda)}$  isomorphic to  $L(\lambda)$ . By Theorem 5.7, there is a Steenrod operation  $\beta = (\beta^r)$  and a polynomial  $v(\lambda) \in \mathbf{P}^d$ , where  $d$  is given by (9), such that  $\beta(v(\lambda)) = w(\mu)$ . Hence the quotient of the submodule generated by  $v(\lambda)$  in  $\mathbf{P}^d$  by the intersection of this submodule with the kernel of  $\beta$  is a composition factor of  $\mathbf{P}^d$  which is isomorphic to  $L(\lambda)$ . Hence the first occurrence degree  $d_c(\lambda) \leq d$ . But  $d_c(\lambda) \geq d$  by [3, Proposition 2.13], and hence  $d_c(\lambda) = d$ .  $\square$

As an example, for  $p = 3$  the partition  $\lambda = (5;3;2)$  is  $\mathbf{T}$ -regular with  $\mathbf{T}$ -conjugate  $\mu = (6;3;1)$ . The module  $L(5;3;2)$  first occurs as a composition factor in degree  $6 + 3 \cdot 3 + 1 \cdot 9 = 24$ , and as a submodule in degree  $5 + 3 \cdot 3 + 2 \cdot 9 = 32$ . The calculations of [1] and [6] for  $n \leq 3$  support the conjecture that the first occurrence degree  $d_c(\lambda)$  is given by the formula above if and only if  $\lambda$  is  $\mathbf{T}$ -regular.

The integers  $r_i$  in Theorem 5.7 can be calculated from a tableau  $\text{Tab}(\lambda)$  obtained by entering integers into the diagram of  $\lambda$  as follows: if a box in row  $i$  is the highest box in its antidiagonal, write  $\rho_{i-1}$  in that box and continue down the antidiagonal, multiplying the number entered at each step by  $\rho$ .

**Lemma 5.9** *The sum of the numbers entered in the  $k$ th block of  $\rho-1$  columns using the above rule is  $r_k$ . The element  $P^{r_1} P^{r_2} \cdots P^{r_m}$  is an admissible monomial in  $A_\rho$ , i.e.  $r_k \leq \rho r_{k+1}$  for  $1 \leq k \leq m-1$ .*

**Example 5.10** For  $\rho = 3$ ,  $\lambda = (6; 5; 4; 3; 2)$ , we obtain  $(r_1; r_2; r_3) = (100; 20; 1)$  using the tableau below.

$$\text{Tab}(\lambda) = \begin{array}{|c|c|c|c|c|c|} \hline 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 & \\ \hline 0 & 0 & 3 & 4 & & \\ \hline 9 & 12 & 13 & & & \\ \hline 39 & 40 & & & & \\ \hline \end{array}$$

Noting that  $\lambda^r = (-1)^r (P^r \lambda)$ , in this case Theorem 5.7 states that in  $\mathbf{P}^{300}$ ,

$$\begin{aligned} (P^{100} P^{20} P^1) x_1^2 [x_1; x_2^3]^2 [x_1; x_2^3; x_3^9]^2 [x_2; x_3^3; x_4^9] [x_3; x_4^3] [x_4; x_5^3] x_5 \\ = -[x_1; x_2^3; x_3^9; x_4^{27}; x_5^{81}]^2 [x_1; x_2^3; x_3^9; x_4^{27}] [x_1; x_2^3; x_3^9] [x_1; x_2^3] x_1 \end{aligned}$$

**Proof of Lemma 5.9** The inequality  $r_k \leq \rho r_{k+1}$  for  $1 \leq k \leq m-1$  is clear from the algorithm, and can also be checked directly from the definition of  $r_k$ . Since  $r_2(\lambda) = r_1(\lambda^-)$ , and so on, we need only check the algorithm for  $r_1$ .

To do this, we introduce a second tableau by entering  $\rho_{i-1}$  in the  $i$ th row of the first block of  $\rho-1$  columns and  $-\rho^{j-2}$  in all the boxes in the  $j$ th block of  $\rho-1$  columns for  $j > 1$ . In Example 5.10 this is as follows.

$$\begin{array}{|c|c|c|c|c|c|} \hline 0 & 0 & -1 & -1 & -3 & -3 \\ \hline 1 & 1 & -1 & -1 & -3 & \\ \hline 4 & 4 & -1 & -1 & & \\ \hline 13 & 13 & -1 & & & \\ \hline 40 & 40 & & & & \\ \hline \end{array}$$

The entries in a antidiagonal running from the  $(i; j)$  box for  $1 \leq j \leq \rho-1$  are then  $\rho_{i-1}; -1; -\rho; \dots; -\rho^{s-2}$ , and their sum  $\rho_{i-1} - \rho_{s-1} = \rho^{s-1} \rho_{i-s}$  is the number entered in this box in  $\text{Tab}(\lambda)$ .

It remains to check that the sum of all the entries in the second tableau is  $r_1 = (b_1 + 1)\rho_{n-1} - (n-1) - d_c(\lambda^-)$ . To see this, note that the entries in  $\lambda^-$  sum to  $-d_c(\lambda^-)$ , while the entries in the last row of  $\lambda^{(1)}$  sum to  $b\rho_{n-1}$  and the entries in the first  $n-1$  rows sum to  $(\rho-1)(\rho_0 + \rho_1 + \dots + \rho_{n-2}) = \rho_{n-1} - (n-1)$ .  $\square$



Since  $w(n)$  is a product of linear factors, so also is  $v(\lambda)$ , and by Theorems 5.3 and 5.5 so also is  $\mathcal{P}^{r_1} v(\lambda)$ . The following calculation shows that  $v(\lambda)$  divides  $\mathcal{P}^{r_1} v(\lambda)$ , and that the quotient can be read off from  $\text{Tab}(\lambda)$  as follows: replace the entry  $\rho_{i-1} - \rho_{s-1}$  in the  $(i; j)$  box,  $1 \leq j \leq \rho - 1$ , by the product of all linear polynomials of the form  $x_i + \sum_{k < i} c_k x_k$ , excluding those where  $c_k = 0$  for  $1 \leq k \leq i - s$ .

**Corollary 5.11** *Let  $\lambda$  be a  $\mathbf{T}$ -regular partition. Let the  $k$ th antidiagonal in the diagram of  $\lambda$  have length  $s_k$  and lowest box in row  $n_k$ . Then*

$$\frac{\mathcal{P}^{r_1} v(\lambda)}{v(\lambda)} = \prod_{k=1}^{\ell(\lambda)} \sum_{\mathbf{c}} (c_1 x_1 + \dots + c_{n_k-1} x_{n_k-1} + x_{n_k});$$

where the inner product is over all vectors  $\mathbf{c} = (c_1; \dots; c_{n_k-1}) \in \mathbb{F}_p^{n_k-1}$  such that  $(c_1; \dots; c_{n_k-s_k}) \neq (0; \dots; 0)$ .

In Theorem 1.1,  $w(\lambda) = ((p - 1)^n)$ ,  $v(\lambda) = (x_1 x_2 \dots x_n)^{p-1}$  and  $\mathcal{P}^{r_1} v(\lambda) = [x_1; x_2^p; \dots; x_n^{p^{n-1}}]^{p-1}$ . Since  $s_k = 1$  for  $1 \leq k \leq n(p - 1)$ , the quotient is the product of all linear polynomials in  $x_1; \dots; x_n$  which are not monomials.

**Proof of Corollary 5.11** The proof is by induction on the number of antidiagonals  $\ell(\lambda)$ . Let  $\mu = \mathcal{P}^{r_1} v(\lambda) = v(\lambda)$ , where  $r_1 = r_1(\lambda)$ . Let  $s$  denote the length of the last antidiagonal in the diagram of  $\lambda$ , and let  $\lambda'$  be the  $\mathbf{T}$ -regular partition obtained by removing this antidiagonal from the diagram of  $\lambda$ . Then by Theorems 5.3 and 5.5,

$$\frac{\mu(\lambda)}{\mu(\lambda')} = \frac{[x_1; x_2^p; \dots; x_n^{p^{n-1}}]}{[x_1; x_2^p; \dots; x_{n-1}^{p^{n-2}}]} \frac{v(\lambda')}{v(\lambda')} \frac{v(\lambda)}{v(\lambda')}.$$

Note that  $\mu(\lambda) = \mu(\lambda')$  when  $s = 1$ . Now  $[x_1; x_2^p; \dots; x_n^{p^{n-1}}] = [x_1; x_2^p; \dots; x_{n-1}^{p^{n-2}}] = \sum_{\mathbf{c}} (c_1 x_1 + \dots + c_{n-1} x_{n-1} + x_n)$ , where the product is taken over all vectors  $\mathbf{c} = (c_1; \dots; c_{n-1}) \in \mathbb{F}_p^{n-1}$ . Also  $v(\lambda) = v(\lambda') = v_{\lambda'}(\lambda) = [x_{n-s+1}; x_{n-s+2}^p; \dots; x_n^{p^{s-1}}]$ . Similarly  $v(\lambda') = v(\lambda'') = [x_{n-s+1}; x_{n-s+2}^p; \dots; x_{n-1}^{p^{s-2}}]$ . The quotient of these determinants is the product of all  $p^{s-1}$  linear polynomials  $c_{n-s+1} x_{n-s+1} + \dots + c_{n-1} x_{n-1} + x_n$ , so  $\mu(\lambda) = \mu(\lambda') = \sum_{\mathbf{c}} (c_1 x_1 + \dots + c_{n-1} x_{n-1} + x_n)$ , where the product is over all  $\mathbf{c} = (c_1; \dots; c_{n-1}) \in \mathbb{F}_p^{n-1}$  with  $c_i \neq 0$  for some  $i$  such that  $1 \leq i \leq n - s$ .

### 6 Proof of the linking theorem

In this section we prove Theorems 5.5 and 5.7. The following lemma will help in checking conditions on the numerical function  $\nu(\cdot)$ .

**Lemma 6.1** (i) Let  $R \geq 1$  have base  $p$  expansion  $R = j_1 p^{a_1} + \dots + j_t p^{a_t}$ , where  $1 \leq j_1, \dots, j_t \leq p-1$ ,  $0 \leq a_1 < \dots < a_t$ , and let  $k \geq 0$ . Then  $(R - p^k) \geq (R) - 1$ , with equality if and only if  $k = a_i$ ,  $1 \leq i \leq t$ .

(ii) With notation as in Theorem 5.5, and with  $\nu$  and  $s$  as in the proof of Corollary 5.11, for  $r \geq 1$  and  $k \geq 0$  we have

$$R(r - p_k + p_{s-1}; \nu) = R(r - p_k + d_c(\nu); \nu) = R(r; \nu) - p^k;$$

**Proof** If  $k \neq a_i$  for  $1 \leq i \leq t$ , then subtraction of  $p^k$  must yield at least one new term  $(p-1)p^a$  in the base  $p$  expansion. This proves (i). For (ii), since  $d_c(\nu) = d_c(\nu) + p d_c(\nu)$  and  $d_c(\nu) = \nu - 1$  we have  $R = R(r; \nu) = (p-1)(r + d_c(\nu)) + \nu - 1$ . Comparing the first occurrence degrees for  $L(\nu)$  and  $L(\nu)$  given by (9),

$$d_c(\nu) = d_c(\nu) + p_s; \quad d_c(\nu) = d_c(\nu) + p_{s-1}; \quad d_c(\nu) = d_c(\nu) + 1; \quad (10)$$

Hence we have  $R(r - p_k + p_{s-1}; \nu) = (p-1)(r - p_k + p_{s-1} + d_c(\nu)) + d_c(\nu) = (p-1)(r - p_k + d_c(\nu)) + d_c(\nu) = R(r - p_k + d_c(\nu); \nu) = R - (p-1)p_k - 1 = R - p^k$ . □

**Proof of Theorem 5.5(i)** We argue by induction on  $\nu - 1$ , the number of antidiagonals of  $\nu$ . With  $\nu$  and  $s$  as above,  $\nu(\nu) = [x_{n-s+1}; x_{n-s+2}^p; \dots; x_n^{p^{s-1}}]$   $\nu(\nu)$ . Using formula (4) and Lemma 2.2, for all  $r \geq 1$  we have

$$p^r \nu(\nu) = \sum_{k=s-1}^{\infty} [x_{n-s+1}; x_{n-s+2}^p; \dots; x_{n-1}^{p^{s-2}}; x_n^{p^k}] p^{r-p_k+p_{s-1}} \nu(\nu); \quad (11)$$

By Lemma 6.1, if  $(R(r; \nu)) > \nu - 1$  then  $(R(r - p_k + p_{s-1}; \nu)) > \nu - 1 - 1$  for all  $k \geq 0$ . Since  $\nu$  has  $\nu - 1$  antidiagonals, the second factor in each term of (11) is zero by the induction hypothesis. Hence  $p^r \nu(\nu) = 0$  if  $(R(r; \nu)) > \nu - 1$ , completing the induction.

**Proof of Theorem 5.5(ii)** As in Lemma 6.1, let  $R = R(r; \nu)$  have base  $p$  expansion  $R = j_1 p^{a_1} + \dots + j_t p^{a_t}$ , let  $(R) = \nu - 1$  and let  $R^0 = R(r - p_k + p_{s-1}; \nu)$ . Then the lemma gives  $(R^0) = \nu - 1 - 1$  if  $k = a_i$ ,  $1 \leq i \leq t$ , and  $(R^0) > \nu - 1 - 1$  otherwise. Hence, applying part (i) of the theorem to (11), we have

$$p^r \nu(\nu) = \sum_{i=1}^t [x_{n-s+1}; x_{n-s+2}^p; \dots; x_{n-1}^{p^{s-2}}; x_n^{p^{a_i}}] p^{r-p_{a_i}+p_{s-1}} \nu(\nu);$$

Since  $(R(r - p_{a_i} + p_{s-1}; \quad)) = \quad - 1 = d_c(\quad)$  by the lemma, and  $p_{s-1} + d_c(\quad) = d_c(\quad)$ , the inductive hypothesis on  $\quad$  gives

$$\mathfrak{p}^{r-p_{a_i}+p_{s-1}}v(\quad) = \mathfrak{p}^{r-p_{a_i}+d_c(\quad)}v(\quad) \quad v(\quad); \quad 1 \leq i \leq t.$$

We can similarly use the lemma to simplify the right hand side of the required identity. Since  $v(\quad) = x_n v(\quad)$ , from (4) and (2) we have

$$\mathfrak{p}^{r+d_c(\quad)}v(\quad) = \sum_{k=0}^{\infty} x_n^{p^k} \mathfrak{p}^{r+d_c(\quad)-p_k}v(\quad):$$

By the lemma,  $R(r + d_c(\quad) - p_k; \quad) = R - p^k$ , so that by (i) we can again reduce to the sum over  $k = a_i, 1 \leq i \leq t$ . As  $v(\quad) = [x_{n-s+1}; x_{n-s+2}^{p^{s-2}}; \dots; x_{n-1}^{p^{s-2}}] v(\quad)$ , it remains after cancelling the factor  $v(\quad)$  and rearranging terms to prove that

$$\sum_{i=1}^t [x_{n-s+1}; x_{n-s+2}^{p^{s-2}}; \dots; x_{n-1}^{p^{s-2}}; x_n^{p^{a_i}}] - [x_{n-s+1}; x_{n-s+2}^{p^{s-2}}; \dots; x_{n-1}^{p^{s-2}}] x_n^{p^{a_i}} f_i = 0;$$

where  $f_i = \mathfrak{p}^{r-p_{a_i}+d_c(\quad)}v(\quad)$ . The expansion of the  $s \times s$  determinant in the  $p^{a_i}$  powers of the variables is

$$\sum_{j=1}^s (-1)^{s-j} [x_{n-s+1}; \dots; x_{n-s+j-1}^{p^{j-2}}; x_{n-s+j+1}^{p^{j-1}}; \dots; x_n^{p^{s-2}}] x_{n-s+j}^{p^{a_i}};$$

Thus the term with  $j = s$  cancels, and interchanging the  $i$  and  $j$  summations, the required formula becomes

$$\sum_{j=1}^{s-1} (-1)^{s-j} [x_{n-s+1}; \dots; x_{n-s+j-1}^{p^{j-2}}; x_{n-s+j+1}^{p^{j-1}}; \dots; x_n^{p^{s-2}}] \sum_{i=1}^t x_{n-s+j}^{p^{a_i}} f_i = 0;$$

Since  $\mathfrak{p}^{r+d_c(\quad)}(x_{n-s+j} v(\quad)) = \sum_{i=1}^t x_{n-s+j}^{p^{a_i}} f_i$  by a similar argument using (4), (1) and Lemma 6.1, it suffices to prove that the monomial  $x_{n-s+j} v(\quad)$  is in the kernel of  $\mathfrak{p}^{r+d_c(\quad)}$  for  $1 \leq j \leq s-1$ . This monomial is divisible by  $x_{n-s+j}^p$ . By permuting the variables, it suffices to consider the case where it is divisible by  $x_1^p$ . Hence the proof of Theorem 5.5 is completed by the following calculation.

**Proposition 6.2** *Let  $R = R(r; \quad)$  and let  $(R) = \quad$ , where  $\quad = (n-1)(p-1) + b$  and  $1 \leq b \leq p-1$ . Then*

$$\mathfrak{p}^{r+d_c(\quad)}(x_1^p(x_2 \dots x_{n-1})^{p-1} x_n^{b-1}) = 0;$$

**Proof** By Lemma 2.3, with  $f = x_1$  and  $g = (x_2 \cdots x_{n-1})^{p-1} x_n^{b-1}$ ,

$$\mathfrak{p}^u(x_1^p | g) = \sum_{u=pv+w} (\mathfrak{p}^v x_1)^p \mathfrak{p}^w(g):$$

Note that  $g = v(\lambda)$  where  $\lambda = ((p-1)^{n-2}(b-1))$ . By (2),  $\mathfrak{p}^v x_1 = 0$  for  $v \notin p_k$ ,  $k \geq 0$ , so we may assume that  $w = u - pv = r + d_c(\lambda) - p \cdot p_k$ . Since  $p \cdot p_k = p_{k+1} - 1$  and  $d_c(\lambda_{(1)}) = p - 1 + d_c(\lambda)$ ,  $R(w; \lambda) = R(r - p_{k+1} + d_c(\lambda); \lambda_{(1)}) = R - p^{k+1}$  by Lemma 6.1(ii). Since  $R(\lambda) = 1$ , Lemma 6.1(i) gives  $(R(w; \lambda))_{(1)} = 1 - 1 > 1 - p$ . Since  $d_c(\lambda) = 1 - p$ ,  $\mathfrak{p}^w g = 0$  by Theorem 5.5(i).  $\square$

**Proof of Theorem 5.7** This follows from Theorem 5.5 by induction on  $m$ . Let  $\lambda_1 = (n-1)(p-1) + b$ ,  $1 \leq b \leq p-1$ . We wish to apply Theorem 5.5 with  $r = r_1$ , so we must check that  $(R(r_1; \lambda_1))_{(1)} = 1$ . For this, note that (9) gives  $d_c(\lambda_1) = \sum_{j=2}^m p^{j-2} j$ , so that  $r_1 + d_c(\lambda_1) = (b+1)p_{n-1} - (n-1)$ . Thus  $R(r_1; \lambda_1) = (p-1)(r_1 + d_c(\lambda_1)) + 1 = (b+1)(p^{n-1} - 1) - (p-1)(n-1) + 1 = bp^{n-1} + (p^{n-1} - 1)$ . Hence  $r_1$  satisfies the hypothesis of Theorem 5.5, so that  $\mathfrak{p}^{r_1} v(\lambda_1) = \mathfrak{p}^{r_1 + d_c(\lambda_1)} v(\lambda_{(1)}) = v(\lambda)$ . By Theorem 5.3,  $\mathfrak{p}^{r_1 + d_c(\lambda_1)} v(\lambda_{(1)}) = w(\lambda_{(1)})$ .

Now  $r_i(\lambda) = r_{i-1}(\lambda)$  for  $2 \leq i \leq m$ , and so the inductive step reduces to showing that

$$\mathfrak{p}^{r_m} \mathfrak{p}^{r_2} w(\lambda_{(1)}) = w(\lambda_{(1)}) \mathfrak{p}^{r_m} \mathfrak{p}^{r_2} v(\lambda): \tag{12}$$

Recall from Lemma 5.9 that  $r_1, \dots, r_m$  is an admissible sequence, i.e.  $r_k \leq p r_{k+1}$  for  $k \geq 1$ . Since  $r_1 \leq (b+1)p_{n-1}$ ,  $r_1 < p^{n-1}$  if  $b < p-1$  and  $r_1 < p^n$  if  $b = p-1$ . Thus we can deduce (12) from Lemma 2.2 and the coproduct formula (4), as follows. We have  $w(\lambda_{(1)}) = w(n)^b w(n-1)^{p-1-b}$ . Now  $\mathfrak{p}^r w(n) = 0$  for  $0 < r < p^{n-1}$  and  $\mathfrak{p}^r w(n-1) = 0$  for  $0 < r < p^{n-2}$ . If there are any factors  $w(n-1)$  in  $w(\lambda_{(1)})$ , then  $r_2 < p^{n-2}$ , and otherwise it suffices to have  $r_2 < p^{n-1}$ .

## 7 First occurrence submodules

For a  $\mathbf{T}$ -regular partition  $\lambda$ , the  $\mathbb{F}_p[M_n]$ -submodule of  $\mathbf{P}^{d_c(\lambda)}$  generated by the first occurrence polynomial  $v(\lambda)$  is a ‘representative polynomial’ for  $L(\lambda)$  in the sense that this module has a quotient isomorphic to  $L(\lambda)$  (see Corollary 5.8). In the case where  $\lambda = (p-1)$  for a column 2-regular partition  $\lambda$ , the leading monomial  $s(\lambda) = x_1^{p-1} \cdots x_n^{p-1}$  has the same property. This is implicit in

the work of Carlisle and Kuhn [2], who identify a subquotient  $\mathbf{T}$  of  $\mathbf{P}^{d_c(\lambda)}$  such that  $\mathbf{T} = \mathbf{T}^1 \oplus \dots \oplus \mathbf{T}^m$ , where  $\lambda^i$  is the  $\mathbf{T}$ -conjugate of  $\lambda$ . Explicitly, if  $v_i \in \mathbf{T}^i$  corresponds to a monomial in  $x_1, \dots, x_n$  with all exponents  $< p$ , then  $v_1 \oplus \dots \oplus v_m \in \mathbf{T}^1 \oplus \dots \oplus \mathbf{T}^m$  corresponds to the equivalence class of  $v_1 \oplus v_2^p \oplus \dots \oplus v_m^{p^{m-1}}$  in the appropriate subquotient of  $\mathbf{P}^{d_c(\lambda)}$ . Proposition 5.4(ii) shows that, taking  $v_j = v(\lambda^{(j)})$ , this monomial is  $s(\lambda)$ . Tri [14] has recently proved that if  $\lambda$  is  $\mathbf{T}$ -regular, then  $L(\lambda)$  is a composition factor in  $\mathbf{T}$ .

We recall from [16, Section 4] the notion of a base  $p$  !-vector.

**Definition 7.1** Given a prime  $p$ , the base  $p$  !-vector  $!_j(s)$  of a sequence of non-negative integers  $s = (s_1; \dots; s_n)$  is defined as follows. Write each  $s_i$  in base  $p$  as  $s_i = \sum_{j=1}^n s_{i,j} p^{j-1}$ , where  $0 \leq s_{i,j} < p$ , and let  $!_j(s) = \sum_{i=1}^n s_{i,j}$ , i.e. add the base  $p$  expansions without ‘carries’. Then  $!(s) = (!_1(s); \dots; !_l(s))$ , with length  $l = \max\{j : !_j(s) > 0\}$  and degree  $d = \sum_{i=1}^n s_i = \sum_{j=1}^l !_j(s) p^{j-1}$ .

Given !-vectors  $\lambda$  and  $\mu$ , we say that  $\lambda$  dominates  $\mu$ , and write  $\lambda \succ \mu$  or  $\mu \prec \lambda$ , if and only if  $\sum_{i=1}^k \lambda_i \geq \sum_{i=1}^k \mu_i$  for all  $k \geq 1$ . By the !-vector of a monomial  $\sum_{i=1}^n x_i^{s_i}$  we mean the !-vector of its sequence of exponents  $s = (s_1; \dots; s_n)$ . The dominance order on !-vectors of the same degree is compatible with left lexicographic order.

**Example 7.2** The lattice of base  $p$  !-vectors of degree  $1 + p + p^2$  is shown below.

$$\begin{array}{c}
 (1 + p + p^2) \\
 \# \\
 (1 + p^2; 1) \\
 \# \\
 \vdots \\
 \# \\
 (1 + p; p) \\
 \cdot \\
 (1; 1 + p) \quad \& \quad (1 + p; 0; 1) \\
 \& \quad \cdot \\
 (1; 1; 1)
 \end{array}$$

**Proposition 7.3** Let  $\lambda$  be a  $\mathbf{T}$ -regular partition. Then the !-vector of the spike monomial  $s(\lambda)$  is the partition  $\mathbf{T}$ -conjugate to  $\lambda$ , and the polynomial  $v(\lambda)$  is the sum of  $(-1)^{l(\lambda)} s(\lambda)$  and monomials  $f$  such that  $!(f) \prec !(s(\lambda))$ .

**Proof** The proof is the same as that given in [16, proposition 4.5], with 2 replaced by  $\rho$  and  $\theta$  replaced by  $\lambda$ . For ( ), see Proposition 5.4(iii).  $\square$

Corollary 5.8 and Proposition 7.3 together provide a ‘topological’ proof that the  $\mathbb{F}_\rho[M_n]$ -submodule of  $\mathbf{P}^{d_c(\lambda)}$  generated by  $s(\lambda)$  has a quotient module isomorphic to  $L(\lambda)$ . The next result provides a further comparison between the spike monomial  $s(\lambda)$  and the polynomial  $v(\lambda)$  in a special case. We conjecture that the corresponding statement holds for all **T**-regular partitions  $\lambda$ .

**Proposition 7.4** *Assume that  $\lambda_i = (\rho - 1) \mu_i$  for  $1 \leq i \leq n$ , where  $\mu = (\mu_1, \dots, \mu_n)$  is a column 2-regular partition. Then the submodule of  $\mathbf{P}^{d_c(\lambda)}$  generated by the polynomial  $v(\lambda)$  is contained in the submodule generated by the spike monomial  $s(\lambda)$ .*

The proof requires a preliminary lemma.

**Lemma 7.5** *If  $f \in \mathbb{F}_\rho[x_2, \dots, x_n]$  and  $1 \leq s \leq n$ , then the  $\mathbb{F}_\rho[M_n]$ -submodule of  $\mathbf{P}$  generated by  $x_1^{\rho^s-1} f$  contains  $[x_1, x_2^\rho, \dots, x_s^{\rho^{s-1}}]^{p-1} f$ .*

**Proof** For each linear form  $v = a_1 x_1 + \dots + a_s x_s$ , where  $a_i \in \mathbb{F}_\rho$  for  $1 \leq i \leq s$ , let  $t_v : \mathbf{P} \rightarrow \mathbf{P}$  be the transvection mapping  $x_1$  to  $v$  and fixing  $x_2, \dots, x_n$ . We claim that the following equation holds in  $\mathbb{F}_\rho[x_1, \dots, x_s]$ .

$$(-1)^s [x_1, x_2^\rho, \dots, x_s^{\rho^{s-1}}]^{p-1} = \sum_v v^{\rho^s-1}; \tag{13}$$

Since  $t_v$  does not change the variables  $x_2, \dots, x_n$  which can occur in  $f$ , it follows from (13) that  $\sum_v t_v$  is an element of the semigroup algebra  $\mathbb{F}_\rho[M_n]$  which maps  $x_1^{\rho^s-1} f$  to  $(-1)^s [x_1, x_2^\rho, \dots, x_s^{\rho^{s-1}}]^{p-1} f$ .

To prove (13), first note that the right hand side is  $GL_s(\mathbb{F}_\rho)$ -invariant. Further, it is mapped to 0 by every singular matrix  $g \in M_s$ , since vectors  $(a_1, \dots, a_s)$  and  $(a_1^g, \dots, a_s^g)$  in  $\mathbb{F}_\rho^s$  in the same coset of the kernel of  $g$  yield terms in (13) with the same image under  $g$ , and  $\rho$  divides the order of this coset. Arguing as in the first or second proof of Theorem 1.1, with  $s$  in place of  $n$ , it follows that (13) holds up to a (possibly zero) scalar.

Finally we verify that the monomial  $m = x_1^{\rho-1} x_2^{\rho(\rho-1)} \dots x_s^{\rho^{s-1}(\rho-1)}$  has coefficient  $(-1)^s$  in the right hand side of (13). For each linear form  $v$ , we have  $v^{\rho^s-1} = v^{\rho^{s-1}(\rho-1)} \dots v^{\rho(\rho-1)} v^{\rho-1}$ , where  $v^{\rho^j(\rho-1)} = (a_1 x_1^{\rho^j} + \dots + a_s x_s^{\rho^j})^{\rho-1}$  for  $0 \leq j \leq s-1$ . The exponent  $\rho-1$  in  $m$  must come from the last factor in this

product, so we must choose the term  $(a_1 x_1)^{\rho-1} = x_1^{\rho-1}$  from the last factor, and  $a_1 \neq 0$ . In the same way, we must choose the term  $(a_2 x_2^\rho)^{\rho-1} = x_2^{\rho(\rho-1)}$  from the last but one factor, and  $a_2 \neq 0$ . Continuing in this way, we see that each of the  $(\rho - 1)^s$  linear forms  $v$  with all coefficients  $a_i \neq 0$  gives a term containing  $m$  (with coefficient 1), while other choices of  $v$  give terms not containing  $m$ . Thus the scalar coefficient in (13) is  $(-1)^s$ .  $\square$

The following example shows how to apply Lemma 7.5 to a partition  $\lambda$  of the form  $(\rho - 1)^s$ , so as to generate  $v(\lambda)$  from  $s(\lambda)$ .

**Example 7.6** Let  $\rho = 3$  and let  $\lambda = (6; 6; 4; 4; 2)$ , so that  $s(\lambda) = x^{26} y^{26} z^8 t^8 u^2$  and  $v(\lambda) = x^2 [x; y^3]^2 [x; y^3; z^9]^2 [y; z^3; t^9]^2 [t; u^3]^2$ .

Begin by permuting the variables, so as to work with the spike  $u^8 t^{26} z^{26} y^8 x^2$ . Apply Lemma 7.5 with  $x_1 = y$  and  $s = 2$  to generate  $[y; x^3]^2 u^8 t^{26} z^{26} x^2$ . Repeat with  $x_1 = z$  and  $s = 3$  to generate  $[z; y^3; x^9]^2 u^8 t^{26} [y; x^3]^2 x^2$ , then with  $x_1 = t$  and  $s = 3$  to generate  $[t; z^3; y^9]^2 u^8 [z; y^3; x^9]^2 [y; x^3]^2 x^2$ , and finally with  $x_1 = u$  and  $s = 2$  to generate  $v(\lambda)$ .

**Proof of Proposition 7.4** We first observe (see [16, Proposition 4.9]) that the (multi)set of lengths of the antidiagonals of the column 2-regular partition  $\lambda$  is equal to the (multi)set of lengths of the rows. Hence the spike monomial  $s(\lambda) = x_n^{\rho^{s_n-1}} x_{n-1}^{\rho^{s_{n-1}-1}} \dots x_1^{\rho^{s_1-1}}$ , where  $s_k$  is the length of the  $k$ th antidiagonal of the diagram of  $\lambda$ , can be obtained from  $s(\lambda)$  by a suitable permutation of the variables. We can now obtain  $v(\lambda)$  from  $s(\lambda)$  by  $n - 1$  successive applications of Lemma 7.5, following the method illustrated by Example 7.6.

## 8 T-regular partitions and the Milnor basis

In this section we link the first occurrence polynomial  $v(\lambda)$  and its leading monomial  $s(\lambda)$  to the polynomial  $\rho(\lambda) = \sum_{j=1}^m w(\lambda_{(j)})^{\rho^j-1}$ , which generates a submodule occurrence of  $L(\lambda)$  in a higher degree. Here, as in Proposition 5.4,  $\lambda_{(j)}$  is the partition given by the  $j$ th block of  $\rho - 1$  columns in the diagram of the **T**-regular partition  $\lambda$ , and  $m$  is the length of  $\lambda$ , the **T**-conjugate of  $\lambda$ . In the case  $\lambda = (\rho - 1)^s$ , we also link the first submodule occurrence polynomial  $w(\lambda)$  to  $\rho(\lambda)$ . The linking is achieved by Milnor basis elements in  $A_\rho$  which are combinatorially related to  $\lambda$ . We also obtain a relation between monomials in **P** and Milnor basis elements in terms of  $!$ -vectors. These results extend some of the results of [16, Section 5].

As in Proposition 5.4, let  $\lambda_i = a_i(p - 1) + b_i$ , where  $a_i \in \{0, 1\}$ ,  $b_i \in \{0, \dots, p - 1\}$ . Following [16], for  $R = ((b_1 + 1)p^{a_1} - 1; \dots; (b_n + 1)p^{a_n} - 1)$  we call the Milnor basis element  $P(R)$  the *Milnor spike* associated to  $\lambda$ . We note that  $!(R) = \lambda$ . A Milnor spike is an admissible monomial [4]. For example, if  $p = 3$  and  $\lambda = (4; 3; 1)$  then the corresponding Milnor spike is  $P(8; 5; 1) = P^{3^2}P^8P^1$ , and for the **T**-conjugate partition  $\mu = (5; 3)$  it is  $P(17; 5) = P^{3^2}P^5$ . In this example,  $\lambda_{(1)} = (3; 2)$  and  $\lambda_{(2)} = (2; 1)$ , so that  $\rho(\lambda) = w(3)w(2) = (w(2)w(1))^3 = [x_1; x_2^3; x_3^9] [x_1; x_2^3]^4 x_1^3$ .

**Theorem 8.1** Let  $\lambda$  be **T**-regular with **T**-conjugate  $\mu$ .

- (i)  $P(R)s(\lambda) = (-1)^{|\lambda|}P(R)v(\lambda) = \rho(\lambda)$ , where  $P(R)$  is the Milnor spike associated to  $(\lambda_2; \dots; \lambda_n)$ .
- (ii) If  $\mu = (p - 1; \dots)$ , where  $\mu$  is column 2-regular,  $P(S)w(\mu) = \rho(\mu)$ , where  $P(S)$  is the Milnor spike associated to  $(\mu_2; \dots; \mu_m)$ .
- (iii) There are formulae corresponding to (i) and (ii) for the Milnor spikes associated to  $\lambda$  and  $\mu$ , with  $\rho(\lambda)$  replaced by  $\rho(\lambda)^p$ .

**Remark 8.2** (iii) follows immediately from (i) and (ii) for degree reasons. The omission of the first terms in  $R$  and  $S$  corresponds to omitting the highest Steenrod power  $P^d$  in the admissible monomial forms of  $P(R)$  and  $P(S)$ . In fact  $d = \deg \rho(\lambda)$ , so that  $P^d \rho(\lambda) = \rho(\lambda)^p$ . In the example  $p = 3$ ,  $\lambda = (4; 3; 1)$  above, (i) states that  $P^8 P^1 (x_1^8 x_2^5 x_3) = -P^8 P^1 (x_1^2 [x_1; x_2^3]^2 [x_2; x_3^3]) = [x_1; x_2^3; x_3^9] [x_1; x_2^3]^4 x_1^3$ . The case  $\mu = (4; 3; 1)$  is excluded from (ii), but in fact  $P^5 w(\mu) = -\rho(\mu)$ . We believe that (ii) holds, up to sign, for all **T**-regular  $\mu$ .

We begin by proving the equivalence of the two statements in (i). For this we use the following generalization of [16, Theorem 5.9(i)]. The proof is based on Lemma 2.8, and follows that given in [16].

**Theorem 8.3** Let  $R = (r_1; \dots; r_t)$  and let  $!(R) = \lambda$ . If the  $!$ -vector  $(x_1^{s_1}; \dots; x_n^{s_n})$  of  $x_1^{s_1}; \dots; x_n^{s_n}$  does not dominate  $\lambda$ , then  $P(R)(x_1^{s_1}; \dots; x_n^{s_n}) = 0$ . □

**Proof of Theorem 8.1(i)** By Proposition 7.3, if the monomial  $f$  occurs in  $v(\lambda)$  and  $f \notin s(\lambda)$ , then  $!(f) < \lambda$ . If  $R = (r_1; \dots; r_n)$  where  $r_i = (b_i + 1)p^{a_i} - 1$ , so that  $P(R)$  is the Milnor spike associated to  $\lambda$ , then, as noted above,  $!(R) = \lambda$ . Hence, by Theorem 8.3,  $P(R)$  takes the same value on  $v(\lambda)$  and on its leading term  $(-1)^{|\lambda|}s(\lambda)$ .

We evaluate  $P(R)s(\lambda)$  by induction on the length  $m$  of  $\lambda$ . The base case  $m = 1$  holds by our previous results, as follows. In this case,  $\lambda = (p - 1; \dots; p - 1; b)$ ,



with  $1 \leq b \leq p - 1$ , and has length  $n$ , while (i) states that  $P(R)s(\sigma) = w(\sigma)$ , where  $R = (p - 1; \dots; p - 1; b)$  has length  $n - 1$ . By Proposition 3.2(ii),  $P(R)g = p^{(b+1)p_{n-1} - (n-1)}g$  when  $\deg g = (n - 1)(p - 1) + b$ , and we may choose  $g = s(\sigma)$ . Hence the result follows from Theorem 5.3.

For the inductive step, we use Proposition 5.4(ii) to write  $s(\sigma) = f^p g$ , where  $g = v(\sigma_{(1)})$  and  $f = s(\sigma)$ . Hence  $P(R)s(\sigma) = (P(S)f)^p P(T)g$  by Lemma 2.8, where the sum is over sequences  $S = (s_2; \dots; s_n)$ ,  $T = (t_2; \dots; t_n)$  such that  $r_i = ps_i + t_i$  for  $2 \leq i \leq n$ . Thus  $t_n = b_1$ ,  $s_n = 0$  and  $t_i \leq p - 1$  for  $2 \leq i \leq n - 1$ . If  $t_i \not\leq p - 1$  for some  $i < n$ , then  $P(T)$  has excess  $\sum_{i=2}^n t_i > \deg v(\sigma_{(1)}) = p - 1$ , so that  $P(T)(v(\sigma_{(1)})) = 0$ . Hence we may assume that  $T = (p - 1; \dots; p - 1; b_1)$ , so that  $s_i = (b_i + 1)p^{a_i - 1} - 1$  for  $2 \leq i \leq n - 1$ . By the argument for the case  $m = 1$ ,  $P(T)(v(\sigma_{(1)})) = w(\sigma_{(1)})$ , and by the induction hypothesis applied to  $\sigma$ ,  $P(S)s(\sigma) = \rho(\sigma)$ . Since  $\rho(\sigma) = w(\sigma_{(1)}) \rho(\sigma)^p$ , the induction is complete.  $\square$

**Proof of Theorem 8.1(ii)** Let  $\sigma = (p - 1)$ , where  $\sigma$  is column 2-regular. Then  $\sigma = (p - 1)$  has length  $m = p - 1$ , and  $\sigma_{(i)} = ((p - 1)^i)$ , so that  $w(\sigma_{(i)}) = w(\sigma_{(i)})^{p-1}$ . Also  $S = (p^{\frac{p}{2}} - 1; \dots; p^{\frac{p}{m}} - 1)$ , so that  $P(S) = P^{t_2} \dots P^{t_m}$ , where  $t_m = p^{\frac{p}{m}} - 1$  and  $t_i = pt_{i+1} + p^{\frac{p}{i}} - 1$  for  $1 \leq i < m$ . We shall argue by induction on  $m$ , the case  $m = 1$ , where  $P(S) = 1$ , being trivial. For  $2 \leq i \leq m$ , let

$$W_i(\sigma) = w(\sigma_{(1)}) w(\sigma_{(i)}) w(\sigma_{(i+1)})^p \dots w(\sigma_{(m)})^{p^{m-i}};$$

so that  $W_1(\sigma) = \rho(\sigma)$  and  $W_m(\sigma) = w(\sigma)$ . We assume as inductive hypothesis on  $j$  that  $P^{t_j} W_j(\sigma) = W_{j-1}(\sigma)$  for  $j > i$ , and prove this for  $j = i$ .

It follows from Lemma 2.1 that  $P^r(w(n)^{p^i}) = 0$  unless  $r = p^i(p_n - p_j)$ , where  $0 \leq j \leq n$ . The largest of these values, equal to the degree of  $w(n)^{p^i}$ , is  $p^i p_n$ . Since  $w(\sigma_{(i)})$  has degree  $p^i - 1$ , it follows by (downward) induction on  $i$  that  $t_i$  is the degree of  $w(\sigma_{(i)}) w(\sigma_{(i+1)})^p \dots w(\sigma_{(m)})^{p^{m-i}}$ . We may express  $t_i$  explicitly as the sum

$$t_i = \sum_{k=i}^m p^{k-i} (p^{\frac{p}{k}} - 1); \tag{14}$$

Hence one term in the expansion of  $P^{t_i}(W_i(\sigma))$  using the Cartan formula is  $W_{i-1}(\sigma)$ . We shall complete the proof by using Lemma 2.1 to show that all other terms in the expansion vanish. Thus we have to consider the possible ways to write  $t_i$  so that

$$(p - 1)t_i = \sum_{v=1}^{p-1} \sum_{k=1}^{p-1} (p^{\frac{p}{k}} - p^{j_{k,v}}) + \sum_{k=i}^m p^{k-i} (p^{\frac{p}{k}} - p^{j_{k,v}}) \tag{15}$$

where  $0 \leq j_{k,v} \leq \binom{0}{k}$  for  $1 \leq k \leq m$ . Equating (14) and (15) and simplifying, we obtain

$$(\rho - 1) \sum_{k=1}^{\infty} \rho^{\binom{0}{k}} + \sum_{k=i}^{\infty} \rho^{k-i} = \sum_{v=1}^{\infty} \sum_{k=1}^{\infty} \rho^{j_{k,v}} + \sum_{k=i}^{\infty} \rho^{k-i} \rho^{j_{k,v}} \quad (16)$$

Since  $\binom{0}{i}$  is column 2-regular,  $\binom{0}{i}$  is strictly decreasing and so  $\binom{0}{i-1} > \binom{0}{i}$ .  $\binom{0}{m} + m - i > m - i$ . Hence the  $m$  powers of  $\rho$  occurring in the left side of (16) are distinct. By uniqueness of base  $\rho$  expansions, there are also  $m$  distinct powers on the right of (16) and these are a permutation of the powers on the left. The argument is now completed as in the case  $\rho = 2$  [16, Section 5].  $\square$

We end with evaluations of certain Milnor basis elements on monomials. While [16, Lemma 5.6] generalizes easily to odd primes, this does not seem to be so useful here as the following (weak) generalization of [16, Proposition 5.8].

**Proposition 8.4** *Let  $R = (r_1; r_2; \dots)$  where  $r_i = \rho - 1$  if  $i = b_1; \dots; b_m$  and  $r_i = 0$  otherwise. Then*

$$P(R)(x_1 \cdots x_n)^{\rho-1} = \begin{cases} [x_1^{\rho^{b_1}}; \dots; x_n^{\rho^{b_n}}]^{\rho-1} & \text{if } m = n; \\ [x_1; x_2^{\rho^{b_1}}; \dots; x_n^{\rho^{b_{n-1}}}]^{\rho-1} & \text{if } m = n - 1; \end{cases}$$

**Proof** This is proved by induction on  $jRj$ . The base of the induction is Theorem 1.1, which is the case  $m = n - 1$ ,  $b_i = i$  for  $1 \leq i \leq n - 1$ . Given a sequence  $R = (r_1; \dots; r_{j-1}; 0; \rho - 1; \rho - 1; \dots; \rho - 1)$ , let  $R^\theta = (r_1; \dots; r_{j-1}; \rho - 1; 0; \rho - 1; \dots; \rho - 1)$ , so that  $jRj - jR^\theta j = (\rho - 1)(\rho^{j+1} - 1) - (\rho - 1)(\rho^j - 1) = (\rho - 1)^2 \rho^j$ . We claim that  $P^{\rho^j(\rho-1)} P(R^\theta)$  and  $P(R)$  have the same value on any polynomial of degree  $n(\rho - 1)$ . To prove this, we use Milnor’s product formula to expand  $P^{\rho^j(\rho-1)} P(R^\theta)$  in the Milnor basis. The Milnor matrix

$$\begin{array}{c|cccccccc} & r_1 & \cdots & r_{j-1} & 0 & 0 & \rho - 1 & \cdots & \rho - 1 \\ \hline 0 & 0 & \cdots & 0 & \rho - 1 & 0 & 0 & \cdots & 0 \end{array}$$

shows that  $P(R)$  occurs with coefficient 1 in the product. Since  $P(R)$  is the unique Milnor basis element of minimal excess  $(n - 1)(\rho - 1)$  in degree  $jRj$ , this proves our claim.

Applying the induction hypothesis to  $P(R^\theta)$ , we have  $P(R)(x_1 \cdots x_n)^{\rho-1} = P^{\rho^j(\rho-1)} [x_1; x_2^{\rho^{b_1}}; \dots; x_j^{\rho^j}; \dots; x_n^{\rho^{b_{n-1}}}]^{\rho-1}$  where  $R$  and  $R^\theta$  differ in the  $i$ th term, i.e.  $b_i = j$  for  $R^\theta$  and  $b_i = j + 1$  for  $R$ . By the Cartan formula, this is  $[x_1; x_2^{\rho^{b_1}}; \dots; x_j^{\rho^{j+1}}; \dots; x_n^{\rho^{b_{n-1}}}]^{\rho-1}$ , and this completes the induction for the case  $m = n - 1$ . The case  $m = n$  is proved similarly.  $\square$

Proposition 8.4 serves as the base of induction for the following generalization of [16, Theorem 5.9(ii)] to odd primes. The proof, by induction on the length of the  $!$ -vector  $\mathbf{s}$ , is essentially the same as in [16].

**Theorem 8.5** *Let  $R_0 = (r_0; r_1; \dots; r_t)$ ,  $R = (r_1; \dots; r_t)$  and  $f = x_1^{s_1} \dots x_n^{s_n}$ , where the base  $p$  expansion of each term  $r_i$  and exponent  $s_j$  contains only the digits 0 and  $p - 1$ . Assume that  $f$  and  $R_0$  have the same  $!$ -vector  $\mathbf{s}$ . Then  $P(R)f = \bigcirc_{k=1}^m \frac{p^{k-1}(\rho-1)}{k}$ , where  $m$  is the length of  $\mathbf{s}$  and  $\Delta_k = [x_{i_1}^{\rho^{j_1}}; \dots; x_{i_j}^{\rho^{j_j}}]$  is the Vandermonde determinant of order  $n = k(\rho - 1)$  defined by the subsequences  $(s_{i_1}; \dots; s_{i_j})$  of  $(s_1; \dots; s_n)$  and  $(r_{j_1}; \dots; r_{j_j})$  of  $R_0$  consisting of the terms whose  $k$ th base  $p$  place is  $p - 1$ .*

**Example 8.6** Using the tables

|       |            |   |         |       |            |   |         |       |                      |  |
|-------|------------|---|---------|-------|------------|---|---------|-------|----------------------|--|
| $r_0$ | $p - 1$    | 0 | $p - 1$ | $x_1$ | $p - 1$    | 0 | $p - 1$ | $r_0$ | $p - 1$              |  |
| $r_1$ | $p - 1$    |   |         | $x_2$ | $p - 1$    |   |         | $r_1$ | $p - 1$ 0 $p - 1$    |  |
| $r_2$ | $p - 1$    |   |         | $x_3$ | $p - 1$    |   |         | $r_2$ | $p - 1$              |  |
| <hr/> |            |   |         | <hr/> |            |   |         | <hr/> |                      |  |
|       | $3(p - 1)$ | 0 | $p - 1$ |       | $3(p - 1)$ | 0 | $p - 1$ |       | $3(p - 1)$ 0 $p - 1$ |  |

we obtain  $P(p - 1; p - 1)x_1^{(\rho^2+1)(\rho-1)}x_2^{\rho-1}x_3^{\rho-1} = [x_1; x_2^{\rho}; x_3^{\rho}]^{\rho-1} x_1^{\rho^2(\rho-1)}$  and  $P((\rho^2 + 1)(\rho - 1); p - 1)x_1^{(\rho^2+1)(\rho-1)}x_2^{\rho-1}x_3^{\rho-1} = [x_1; x_2^{\rho}; x_3^{\rho}]^{\rho-1} (x_1^{\rho})^{\rho^2(\rho-1)}$ .

## References

- [1] D. P. Carlisle, The modular representation theory of  $GL(n; p)$  and applications to topology, Ph.D. thesis, University of Manchester, 1985.
- [2] D. P. Carlisle and N. J. Kuhn, Subalgebras of the Steenrod algebra and the action of matrices on truncated polynomial algebras, J. of Algebra 121 (1989), 370{387.
- [3] D. P. Carlisle and G. Walker, Poincare series for the occurrence of certain modular representations of  $GL(n; p)$  in the symmetric algebra, Proc. Roy. Soc. Edinburgh 113A (1989), 27{41.
- [4] D. P. Carlisle, G. Walker and R. M. W. Wood, The intersection of the admissible basis and the Milnor basis of the Steenrod algebra, J. Pure and Applied Algebra 128 (1998), 1{10.
- [5] D. M. Davis, The antiautomorphism of the Steenrod algebra, Proc. Amer. Math. Soc. 44 (1974), 235{236.
- [6] S. R. Doty and G. Walker, The composition factors of  $F_p[x_1; x_2; x_3]$  as a  $GL(3; p)$ -module, J. Algebra 147 (1992), 411{441.

- [7] S. R. Doty and G. Walker, Truncated symmetric powers and modular representations of  $GL_n$ , *Math. Proc. Camb. Phil. Soc.* 119 (1996), 231{242.
- [8] J. A. Green, Polynomial representations of  $GL_n$ , *Lecture Notes in Mathematics* 830, Springer 1980.
- [9] J. C. Harris and N. J. Kuhn, Stable decomposition of classifying spaces of finite abelian  $p$ -groups, *Math. Proc. Camb. Phil. Soc.* 103 (1988), 427{449.
- [10] G. D. James and A. Kerber, The representation theory of the symmetric group, *Encyclopaedia of Mathematics*, vol. 16, Addison-Wesley (1981).
- [11] J. Milnor, The Steenrod algebra and its dual, *Ann. Math.* 67 (1958), 150{171.
- [12] P. A. Minh and T. T. Tri, The first occurrence for the irreducible modules of general linear groups in the polynomial algebra, *Proc. Amer. Math. Soc.* 128 (2000), 401{405.
- [13] W. Singer, On the action of Steenrod squares on polynomial algebras, *Proc. Amer. Math. Soc.* 111 (1991), 577{583.
- [14] T. T. Tri, On the first occurrence of irreducible representations of semigroup of all matrices as composition factors in the polynomial algebra, *Acta Math. Vietnamica*, to appear.
- [15] G. Walker, Modular Schur functions, *Trans. Amer. Math. Soc.* 346 (1994), 569{604.
- [16] G. Walker and R. M. W. Wood, Linking first occurrence polynomials over  $\mathbb{F}_2$  by Steenrod operations, *J. Algebra* 246 (2001), 739{760.
- [17] R. M. W. Wood, Splitting  $(CP^1 \times \dots \times CP^1)$  and the action of Steenrod squares on the polynomial ring  $\mathbb{F}_2[x_1, \dots, x_n]$ , *Algebraic Topology Barcelona 1986*, *Lecture Notes in Mathematics* 1298, Springer-Verlag (1987), 237{255.

*Department of Mathematics, College of Sciences*  
*University of Hue, Dai hoc Khoa hoc, Hue, Vietnam*  
 and  
*Department of Mathematics, University of Manchester*  
*Oxford Road, Manchester M13 9PL, U.K.*

Email: pami nh@dng.vnn.vn, grant@ma.man.ac.uk

Received: 24 January 2002