


Geometry & Topology
Volume 6 (2002) 355{360
Published: 28 June 2002



Burnside obstructions to the Montesinos{Nakanishi 3{move conjecture

Mieczysław K Dabkowski
Jozef H Przytycki

*Department of Mathematics, The George Washington University
Washington, DC 20052, USA*

Email: mdab@gwu.edu, przytyck@gwu.edu

Abstract

Yasutaka Nakanishi asked in 1981 whether a 3{move is an unknotting operation. In Kirby's problem list, this question is called *The Montesinos{Nakanishi 3{move conjecture*. We define the n th Burnside group of a link and use the 3rd Burnside group to answer Nakanishi's question; ie, we show that some links cannot be reduced to trivial links by 3{moves.

AMS Classification numbers Primary: 57M27

Secondary: 20D99

Keywords: Knot, link, tangle, 3{move, rational move, braid, Fox coloring, Burnside group, Borromean rings, Montesinos{Nakanishi conjecture, branched cover, core group, lower central series, associated graded Lie ring, skein module

Proposed: Robion Kirby
Seconded: Walter Neumann, Vaughan Jones

Received: 5 May 2002
Revised: 19 June 2002

One of the oldest elementary formulated problems in classical Knot Theory is the 3-move conjecture of Nakanishi. A 3-move on a link is a local change that involves replacing parallel lines by 3 half-twists (Figure 1).




Figure 1

Conjecture 1 (Montesinos{Nakanishi, Kirby’s problem list; Problem 1.59(1), [4]) *Any link can be reduced to a trivial link by a sequence of 3-moves.*

The conjecture has been proved to be valid for several classes of links by Chen, Nakanishi, Przytycki and Tsukamoto (eg, closed 4-braids and 4-bridge links).

Nakanishi, in 1994, and Chen, in 1999, have presented examples of links which they were not able to reduce: L_{2BR} , the 2-parallel of the Borromean rings, and $\hat{\Delta}$, the closure of the square of the center of the n th braid group, ie, $\Delta_n = (\sigma_1 \sigma_2 \sigma_3 \dots \sigma_{n-1})^{10}$.

Remark 2 In [6] it was noted that 3-moves preserve the first homology of the double branched cover of a link L with \mathbb{Z}_3 coefficients ($H_1(M_L^{(2)}; \mathbb{Z}_3)$). Suppose that $\hat{\Delta}$ (respectively L_{2BR}) can be reduced by 3-moves to the trivial link T_n . Since $H_1(M_{\hat{\Delta}}^{(2)}; \mathbb{Z}_3) = \mathbb{Z}_3^4$, $H_1(M_{L_{2BR}}^{(2)}; \mathbb{Z}_3) = \mathbb{Z}_3^5$ and $H_1(M_{T_n}^{(2)}; \mathbb{Z}_3) = \mathbb{Z}_3^{n-1}$ where T_n is a trivial link of n components, it follows that $n = 5$ (respectively $n=6$).

We show below that neither $\hat{\Delta}$ nor L_{2BR} can be reduced by 3-moves to trivial links.

The tool we use is a non-abelian version of Fox n -colorings, which we shall call the n th Burnside group of a link, $B_L(n)$.

Definition 3 The n th Burnside group of a link is the quotient of the fundamental group of the double branched cover of S^3 with the link as the branch set divided by all relations of the form $a^n = 1$. Succinctly: $B_L(n) = \pi_1(M_L^{(2)}) / \langle a^n \rangle$.

Proposition 4 $B_L(3)$ is preserved by 3-moves.

Proof In the proof we use the core group interpretation of $\pi_1(M_L^{(2)})$. Let D be a diagram of a link L . We define (after [3, 2]) the associated core group $\pi_D^{(2)}$ of D as follows: generators of $\pi_D^{(2)}$ correspond to arcs of the diagram. Any crossing v_s yields the relation $r_s = y_i y_j^{-1} y_i y_k^{-1}$ where y_i corresponds to the overcrossing and y_j, y_k correspond to the undercrossings at v_s (see Figure 2). In this presentation of $\pi_D^{(2)}$ one relation can be dropped since it is a consequence of others. Wada proved that $\pi_D^{(2)} = \pi_1(M_L^{(2)})$ [10] (see [7] for an elementary proof using only Wirtinger presentation). Furthermore, if we put $y_i = 1$ for any fixed generator, then $\pi_D^{(2)}$ reduces to $\pi_1(M_L^{(2)})$. The last part of our proof is illustrated in Figure 2. \square




Figure 2

Lemma 5 $B^\wedge(3) = \langle x_1, x_2, x_3, x_4 \mid a^3 \rangle$ for any word $a = P_1 P_2 P_3 P_4 g$, where

$$P_i = x_1 x_2^{-1} x_3 x_4^{-1} x_1^{-1} x_2 x_3^{-1} x_4 x_i x_4 x_3^{-1} x_2 x_1^{-1} x_4^{-1} x_3 x_2^{-1} x_1 x_i^{-1}$$

Proof Consider the 5-braid $\sigma = (\sigma_{12} \sigma_{34})^{10}$ (Figure 3). If we label initial arcs of the braid by x_1, x_2, x_3, x_4 and x_5 , and use core relations (progressing from left to right) we obtain labels Q_1, Q_2, Q_3, Q_4 and Q_5 on the final arcs of the braid where

$$Q_i = x_1 x_2^{-1} x_3 x_4^{-1} x_5 x_1^{-1} x_2 x_3^{-1} x_4 x_5^{-1} x_i x_5^{-1} x_4 x_3^{-1} x_2 x_1^{-1} x_5 x_4^{-1} x_3 x_2^{-1} x_1$$

For a group $\pi_\wedge^{(2)}$, of the closed braid \wedge , we have relations $Q_i = x_i$. To obtain $\pi_1(M_\wedge^{(2)})$ we can put $x_5 = 1$, and delete one relation, say $Q_5 x_5^{-1}$. These lead to the presentation of $B^\wedge(3)$ described in the lemma. \square

Theorem 6 *The links \wedge and L_{2BR} are not 3-move reducible to trivial links.*



Figure 3

Proof Let $B(n;3)$ denote the classical free n generator Burnside group of exponent 3. As shown by Burnside [1], $B(n;3)$ is a finite group. Its order, $|B(n;3)|$, is equal to $3^{n + \binom{n}{2} + \binom{n}{3}}$. For a trivial link: $B_{T_k}(3) = B(k-1;3)$. In order to prove that $\hat{\Delta}$ and L_{2BR} are not 3-move reducible to trivial links, it suffices to show that $B_{\hat{\Delta}}(3) \notin B(4;3)$ and $B_{L_{2BR}}(3) \notin B(5;3)$ (see Remark 2). We have demonstrated these to be true both by manual computation, and by using the programs GAP, Magnus and Magma. More details in the case of $\hat{\Delta}$ are provided below.

For the manual calculations, one first observes that for any i , P_i is in the third term of the lower central series of $B(4;3)$. In particular, for $u = x_1 x_2^{-1} x_3 x_4^{-1}$ and $v = x_1^{-1} x_2 x_3^{-1} x_4$, one has $uv \in [B(4;3); B(4;3)]$ and $P_i = [uv; x_i u]$. It is known ([9]), that $B(4;3)$ is of class 3 (the lower central series has 3 terms), and that the third term is isomorphic to Z_3^4 with basis: $e_1 = [[x_2; x_3]; x_4]$, $e_2 = [[x_1; x_3]; x_4]$, $e_3 = [[x_1; x_2]; x_4]$ and $e_4 = [[x_1; x_2]; x_3]$. It now takes an elementary linear algebra calculation (see Lemma 7 below) to show that $P_1; P_2; P_3; P_4$ form another basis of the third term of the lower central series of $B(4;3)$. Thus $|B_{\hat{\Delta}}(3)| = 3^{10}$. \square

Lemma 7 $P_1; P_2; P_3$, and P_4 form a basis of the third term of the lower central series of $B(4;3)$.

Proof In the associated graded Lie ring $L(4;3)$ of $B(4;3)$ ([9]), the third term (denoted L_3) is isomorphic to Z_3^4 with basis $e_1; e_2; e_3; e_4$. In $L(4;3)$, which is a linear space over Z_3 , one uses an additive notation and the bracket in the group becomes a (non-associative) product ([9]). In this notation $e_1 = x_2 x_3 x_4$, $e_2 = x_1 x_3 x_4$, $e_3 = x_1 x_2 x_4$ and $e_4 = x_1 x_2 x_3$. In the calculation expressing P_i in the basis we use the following identities in L_3 ([9]; page 89).

$$xyzt = 0; xyz = yzx = zxy = -xzy = -zyx = -yxz; xyy = 0;$$

Now we have: $P_i = (uv)(x_i u)(uv)^{-1}(x_i u)^{-1} = [(uv)^{-1}; (x_i u)^{-1}] = [uv; x_i u]$ as the last term of the lower central series is in the center of $B(4;3)$. Furthermore, we have $uv = x_1 x_2^{-1} x_3 x_4^{-1} x_1^{-1} x_2 x_3^{-1} x_4 = [x_2^{-1} x_3 x_4^{-1}; x_1^{-1}][x_3 x_4^{-1}; x_2][x_4^{-1}; x_3^{-1}]$.

Writing P_i additively in L_3 one obtains:

$$P_i = ((-x_2 + x_3 - x_4)(-x_1) + (x_3 - x_4)x_2 + x_4x_3)(x_i - x_1 + x_2 - x_3 + x_4):$$

After simplifications one gets:

$$P_1 = -e_1; P_2 = e_1 + e_2; P_3 = e_1 - e_2 - e_3; \text{ and } P_4 = e_1 - e_2 + e_3 + e_4:$$

The matrix expressing P_i 's in terms of e_i 's is the upper triangular matrix with the determinant equal to 1. Therefore the lemma follows. \square

A similar calculation establishes that $jB_{L_2BR}(3)j < jB(5;3)j$. $B(5;3)$ is of class 3 and has 3^{25} elements. Considering L_2BR as a closed 6{braid we note that $B_{L_2BR}(3)$ is obtained from $B(5;3)$ by adding 5 relations $R_1; \dots; R_5$. Relations $fR_i g$ are in the last term of the lower central series of $B(5;3)$ (and of the associated graded algebra $L(5;3)$). Relations form a 4{dimensional subspace in $L_3 = Z_3^{10}$. Thus $jB_{L_2BR}(3)j = 3^{21}$.

For a computer verification showing that $B^\wedge(3) \notin B(4;3)$ consider any presentation of $B(4;3)$ (eg, Magma solution by Mike Newman [5]) and add the relations P_i to obtain a presentation of $B^\wedge(3)$. Using any of the algebra programs mentioned above, one verifies that $jB^\wedge(3)j = 3^{10}$ while $jB(4;3)j = 3^{14}$.

The solution of the Nakanishi{Montesinos 3{move conjecture, presented above, is the first instance of application of Burnside groups of links. It was motivated by the analysis of cubic skein modules of 3{manifolds. The next step is the application of Burnside groups to rational moves on links. This, in turn, should have deep implications to the theory of skein modules [7].

References

- [1] **W Burnside**, *On an Unsettled Question in the Theory of Discontinuous Groups*, Quart. J. Pure Appl. Math. 33 (1902) 230{238
- [2] **R Fenn, C Rourke**, *Racks and links in codimension two*, Journal of Knot Theory and its Ramifications, 1 (1992) 343{406
- [3] **D Joyce**, *A classifying invariant of knots: the knot quandle*, Jour. Pure Appl. Alg. 23 (1982) 37{65
- [4] **R Kirby**, *Problems in low-dimensional topology; Geometric Topology*, from "Proceedings of the Georgia International Topology Conference, 1993", Studies in Advanced Mathematics, Volume 2 part 2 (W Kazez, Editor) AMS/IP (1997) 35{473
- [5] **M Newman**,
<http://magma.maths.usyd.edu.au/magma/Examples/node15.html>

- [6] **JH Przytycki**, t_k moves on links, from "Braids", (JS Birman and A Libgober, Editors) Contemporary Math. Vol. 78 (1988) 615{656
- [7] **JH Przytycki**, Skein modules of 3-manifolds, Bull. Ac. Pol.: Math. 39 (1991) 91{100
- [8] **JH Przytycki**, 3-coloring and other elementary invariants of knots, Banach Center Publications, Vol. 42, "Knot Theory" (1998) 275{295
- [9] **M Vaughan-Lee**, *The restricted Burnside problem*, second edition, London Mathematical Society Monographs no. 8, The Clarendon Press and Oxford University Press, New York (1993)
- [10] **M Wada**, Group invariants of links, Topology, 31 (1992) 399{406