


Geometry & Topology
 Volume 9 (2005) 203–217
 Published: 20 January 2005



A stable classification of Lefschetz fibrations

DENIS AUROUX

*Department of Mathematics, MIT
 Cambridge MA 02139, USA*

Email: auroux@math.mit.edu

Abstract

We study the classification of Lefschetz fibrations up to stabilization by fiber sum operations. We show that for each genus there is a “universal” fibration f_g^0 with the property that, if two Lefschetz fibrations over S^2 have the same Euler–Poincaré characteristic and signature, the same numbers of reducible singular fibers of each type, and admit sections with the same self-intersection, then after repeatedly fiber summing with f_g^0 they become isomorphic. As a consequence, any two compact integral symplectic 4–manifolds with the same values of $(c_1^2, c_2, c_1 \cdot [\omega], [\omega]^2)$ become symplectomorphic after blowups and symplectic sums with f_g^0 .

AMS Classification numbers Primary: 57R17

Secondary: 53D35

Keywords: Symplectic 4–manifolds, Lefschetz fibrations, fiber sums, mapping class group factorizations

Proposed: Tomasz Mrowka
 Seconded: Ronald Fintushel, Ronald Stern

Received: 7 December 2004
 Accepted: 18 January 2005

1 Introduction

Lefschetz fibrations have been the focus of a lot of attention ever since it was shown by Donaldson that, after blow-ups, every compact symplectic 4-manifold admits such structures [2]. We recall the definition:

Definition 1 A Lefschetz fibration on an oriented compact smooth 4-manifold M is a smooth map $f: M \rightarrow S^2$ which is a submersion everywhere except at finitely many non-degenerate critical points p_1, \dots, p_r , near which f identifies in local orientation-preserving complex coordinates with the model map $(z_1, z_2) \mapsto z_1^2 + z_2^2$.

The smooth fibers of f are compact surfaces, and the singular fibers present nodal singularities; each singular fiber is obtained by collapsing a simple closed loop (the *vanishing cycle*) in the smooth fiber. The monodromy of the fibration around a singular fiber is given by a positive Dehn twist along the vanishing cycle.

Denoting by $q_1, \dots, q_r \in S^2$ the images of the critical points (which we will always assume to be distinct), and choosing a reference point $q_* \in S^2 \setminus \text{crit}(f)$, we can characterize the fibration f by its *monodromy homomorphism*

$$\psi: \pi_1(S^2 \setminus \{q_1, \dots, q_r, q_*\}) \rightarrow \text{Map}_g,$$

where $\text{Map}_g = \pi_0 \text{Diff}^+(\Sigma_g)$ is the mapping class group of a genus g surface. It is a classical result (cf. [4]) that the monodromy morphism ψ is uniquely determined up to conjugation by an element of Map_g and the action of a braid on $\pi_1(S^2 \setminus \{q_i\})$ by “Hurwitz moves” (see Section 2); moreover, if the fiber genus is at least 2 then the monodromy determines the isomorphism class of the Lefschetz fibration f .

The classification of Lefschetz fibrations is a difficult problem (essentially as difficult as the classification of symplectic 4-manifolds), and is only understood in genus 1 and 2 (with some assumptions on the nature of singular fibers in the latter case). It is a classical result of Moishezon and Livne [7] that genus 1 Lefschetz fibrations are all holomorphic, and are classified by the number of irreducible singular fibers (which is a multiple of 12) and the number of reducible singular fibers. More recently, Siebert and Tian [9] have obtained a classification result for genus 2 Lefschetz fibrations without reducible singular fibers and with “transitive monodromy” (a technical assumption which we will not discuss here). Namely, these fibrations are all holomorphic, and are classified by their number of vanishing cycles, which is always a multiple of 10. In fact, all such

fibrations can be obtained as fiber sums of two standard holomorphic fibrations f_0 and f_1 with respectively 20 and 30 singular fibers.

In higher genus, or even in genus 2 if one allows reducible singular fibers, the classification appears to be much more complicated. However, we can attempt to determine a minimal set of moves (ie, surgery operations) which can be used to relate to each other any two Lefschetz fibrations with the same genus. In this context, we consider stabilization by *fiber sums* with certain standard fibrations. (The fiber sum of two Lefschetz fibrations is obtained by deleting a neighborhood of a smooth fiber in each of them, and gluing the resulting open manifolds along their boundaries in a fiber-preserving manner). It was shown in [1] that, given two genus 2 symplectic Lefschetz fibrations f, f' with the same numbers of singular fibers of each type (irreducible, reducible with genus 1 components, reducible with components of genus 0 and 2), for all large n the fiber sums $f \# n f_0$ and $f' \# n f_0$ are isomorphic. More generally, as a corollary of a recent result of Kharlamov and Kulikov about braid monodromy factorizations [5], a similar result holds for all Lefschetz fibrations with monodromy contained in the hyperelliptic mapping class group.

Our goal is to obtain a similar stabilization result in the general case (without assumptions on the fiber genus or on the monodromy). In this context we must consider pairs of Lefschetz fibrations f, f' with the same fiber genus and the same numbers of singular fibers of each type (irreducible, or reducible of *type* $(h, g - h)$, ie, with components of genera h and $g - h$, for each $0 \leq h \leq \frac{g}{2}$), but we must also place two additional restrictions (which automatically hold when $g \leq 2$ or in the hyperelliptic case). Namely, we must assume that the intersection forms on the total spaces M and M' have the same signature, and we must assume that the fibrations f and f' admit distinguished sections s, s' which represent classes in $H_2(M, \mathbb{Z})$ (resp. $H_2(M', \mathbb{Z})$) with the same self-intersection number $-k$.

Then, we claim that, after repeatedly fiber summing f and f' with a certain “universal” Lefschetz fibration f_g^0 , constructed in Section 3, we eventually obtain isomorphic Lefschetz fibrations:

Theorem 2 *For every g there exists a genus g Lefschetz fibration f_g^0 with the following property. Let $f: M \rightarrow S^2$ and $f': M' \rightarrow S^2$ be two genus g Lefschetz fibrations, each equipped with a distinguished section. Assume that:*

- (i) *the total spaces M and M' have the same Euler characteristic and signature;*
- (ii) *the distinguished sections of f and f' have the same self-intersection;*

(iii) f and f' have the same numbers of reducible fibers of each type.

Then, for all large enough values of n , the fiber sums $f \#_n f_g^0$ and $f' \#_n f_g^0$ are isomorphic.

A brief remark is in order about assumptions (i) and (ii) in this statement. First, since the Euler characteristic is given by the formula $\chi = 4 - 4g + r$, where g is the fiber genus and r is the total number of singular fibers, the first part of (i) is equivalent to the requirement that f and f' have the same numbers of singular fibers. Moreover, in the hyperelliptic case the assumption on signature can be eliminated, because the signature is given by Endo's formula [3], which involves only the number of singular fibers of each type; however, in general the signature depends on the actual vanishing cycles. It is also worth mentioning that, in general, it is not known whether every Lefschetz fibration admits a section (although there are no known examples without a section). However, all Lefschetz fibrations obtained by blowing up the base points of a pencil (and in particular all those which arise from Donaldson's construction) admit sections of square -1 .

The cases $g = 0$ and $g = 1$ of Theorem 2 are trivial (in that case no stabilization is needed), and the case $g = 2$ is proved in [1] (taking f_2^0 to be the holomorphic genus 2 fibration with 20 singular fibers and total space a rational surface). Thus we will only consider the case $g \geq 3$ in the proof.

As a corollary of Theorem 2 and of Donaldson's result, we have the following statement for *integral* compact symplectic 4-manifolds (ie, such that $[\omega] \in H^2(X, \mathbb{R})$ is the image of an integer cohomology class):

Corollary 3 *Let X, X' be two integral compact symplectic 4-manifolds with the same $(c_1^2, c_2, c_1 \cdot [\omega], [\omega]^2)$. Then X and X' become symplectomorphic after sufficiently many blowups and symplectic sums with the total space X_g^0 of the fibration f_g^0 (for a suitable genus g).*

This corollary follows from Theorem 2 by considering pencils of the same (large) degree d on X and X' , and blowing up the $d^2[\omega]^2$ base points. The resulting Lefschetz fibrations have the same fiber genus (by the assumptions on $c_1 \cdot [\omega]$ and $[\omega]^2$), admit sections of square -1 , and, if d is large enough, can be assumed to contain only irreducible fibers.

The proof of Theorem 2 actually gives a complete classification of Lefschetz fibrations up to fiber sum stabilization. For example, considering only Lefschetz fibrations with irreducible fibers, we have:

Theorem 4 For every $g \geq 3$ there exist Lefschetz fibrations $f_g^A, f_g^B, f_g^C, f_g^D$ with the following property: if f is a genus g Lefschetz fibration without reducible singular fibers, and if f admits a section, then there exist integers $a, b, c, d \in \mathbb{Z}$ such that for all large enough values of n the fiber sums $f \# n f_g^0$ and $(n + a) f_g^A \# (n + b) f_g^B \# (n + c) f_g^C \# (n + d) f_g^D$ are isomorphic.

The Lefschetz fibrations $f_g^A, f_g^B, f_g^C, f_g^D$ are constructed in Section 3 (and f_g^0 is in fact nothing but their fiber sum).

The rest of this paper is organized as follows: in Section 2 we review the description of Lefschetz fibrations by mapping class group factorizations; in Section 3 we introduce the concept of universal positive factorization and construct the Lefschetz fibrations f_g^0 ; and in Sections 4–5 we prove Theorem 2.

This work was partially supported by NSF grant DMS-0244844.

2 Mapping class group factorizations

The monodromy of a Lefschetz fibration can be encoded in a *mapping class group factorization* by choosing an ordered system of generating loops $\gamma_1, \dots, \gamma_r$ for $\pi_1(S^2 \setminus \{q_1, \dots, q_r\})$, such that each loop γ_i encircles only one of the points q_i and $\prod \gamma_i$ is homotopically trivial. The monodromy of the fibration along each of the loops γ_i is a Dehn twist τ_i ; we can then describe the fibration in terms of the relation $\tau_1 \dots \tau_r = 1$ in Map_g . The choice of the loops γ_i (and therefore of the twists τ_i) is of course not unique, but any two choices differ by a sequence of *Hurwitz moves* exchanging consecutive factors: $\tau_i \cdot \tau_{i+1} \rightarrow (\tau_{i+1})_{\tau_i^{-1}} \cdot \tau_i$ or $\tau_i \cdot \tau_{i+1} \rightarrow \tau_{i+1} \cdot (\tau_i)_{\tau_{i+1}}$, where we use the notation $(\tau)_\phi = \phi^{-1} \tau \phi$, ie, if τ is a Dehn twist along a loop δ then $(\tau)_\phi$ is the Dehn twist along the loop $\phi(\delta)$.

Definition 5 A factorization $F = \tau_1 \cdot \dots \cdot \tau_r$ in Map_g is an ordered tuple of positive Dehn twists. We say that two factorizations are *Hurwitz equivalent* ($F \sim F'$) if they can be obtained from each other by a sequence of Hurwitz moves.

A Lefschetz fibration is thus characterized by a factorization of the identity element in Map_g , uniquely determined up to Hurwitz equivalence and simultaneous conjugation of all factors by a same element of Map_g , ie, up to the equivalence relation generated by the moves

$$\tau_1 \cdot \dots \cdot \tau_i \cdot \tau_{i+1} \cdot \dots \cdot \tau_r \longleftarrow \tau_1 \cdot \dots \cdot \tau_{i+1} \cdot (\tau_i)_{\tau_{i+1}} \cdot \dots \cdot \tau_r \quad \forall 1 \leq i < r,$$

$$\tau_1 \cdots \tau_r \longleftrightarrow (\tau_1)_\phi \cdots (\tau_r)_\phi \quad \forall \phi \in \text{Map}_g.$$

We will actually be considering Lefschetz fibrations equipped with a distinguished section. The section determines a marked point in each fiber, and so we can lift the monodromy to a *relative* mapping class group. In fact, even though the normal bundle to the section s is not trivial (it has degree $-k$ for some $k \geq 1$), we can restrict ourselves to the preimage of a large disc Δ containing all the critical values of f , and fix a trivialization of the normal bundle to s over Δ . Deleting a small tubular neighborhood of the section s , we can now view the monodromy of f as a homomorphism

$$\psi: \pi_1(\Delta \setminus \{q_1, \dots, q_r\}) \rightarrow \text{Map}_{g,1},$$

where $\text{Map}_{g,1}$ is the mapping class group of a genus g surface with one boundary component. The product of the Dehn twists $\tau_i = \psi(\gamma_i)$ is not the identity, but the central element $T_\delta^k \in \text{Map}_{g,1}$, where T_δ is the *boundary twist*, ie, the Dehn twist along a loop parallel to the boundary.

With this understood, a Lefschetz fibration with a distinguished section of square $-k$ is described by a factorization of T_δ^k as a product of positive Dehn twists in $\text{Map}_{g,1}$, up to Hurwitz equivalence and global conjugation.

A word about notations: while we use the multiplicative notation for factorizations, and sometimes write $\tau_1 \cdots \tau_r = T_\delta^k$ to express the fact that $\tau_1 \cdots \tau_r$ is a factorization of T_δ^k , it is important not to confuse a factorization (a tuple of Dehn twists) with the product of its factors (an element in $\text{Map}_{g,1}$). We will also use multiplicative notation for the concatenation of factorizations ($F \cdot F'$ is the factorization consisting of the factors in F , followed by those in F' , and $(F)^n$ is the concatenation of n copies of F), and we will denote by $(F)_\phi$ the factorization obtained by conjugating each factor of F by the element $\phi \in \text{Map}_{g,1}$.

To finish this section, we establish the following properties of Hurwitz equivalence for factorizations of central elements:

Lemma 6 *Let T be a central element in a group G . Then:*

- (a) *if $F' \cdot F''$ is a factorization of T , then $F'' \cdot F'$ is also a factorization of T , and $F' \cdot F'' \sim F'' \cdot F'$;*
- (b) *if F is a factorization of T whose factors generate G , then $F \sim (F)_\phi$ for all $\phi \in G$;*
- (c) *if F is a factorization of T , and F' is any factorization, then $F' \cdot F \sim F \cdot F'$.*

Proof (see also Lemma 6 in [1]).

(a) To prove that any cyclic permutation of the factors amounts to a Hurwitz equivalence, it suffices to prove that if $\tau \in G$ and $\tau \cdot F''$ is a factorization of T then $\tau \cdot F'' \sim F'' \cdot \tau$. Denote by ϕ the product of the factors in F'' : using Hurwitz moves to move all the factors in F'' to the left of τ , we have $\tau \cdot F'' \sim F'' \cdot (\tau)_\phi$. The result then follows from the observation that $\phi = \tau^{-1}T$ commutes with τ .

(b) Let τ be any of the factors in F : then by (a) we can perform a cyclic permutation of the factors and obtain a factorization F' such that $F \sim F' \cdot \tau$. Moving τ to the left of F' , we have $F' \cdot \tau \sim \tau \cdot (F')_\tau = (\tau \cdot F')_\tau$. Applying (a) again we have $(\tau \cdot F')_\tau \sim (F)_\tau$. So, for any factor τ of F , we have $F \sim (F)_\tau$, and similarly $F \sim (F)_{\tau^{-1}}$. The result then follows from the assumption that the factors of F generate G , by expressing ϕ in terms of the factors.

(c) Simply move all the factors of F to the left of the factors in F' , to obtain $F' \cdot F \sim F \cdot (F')_T = F \cdot F'$ (since T is central). □

3 Universal positive factorizations

Let us first recall a presentation of $\text{Map}_{g,1}$ due to Matsumoto [6], which is a reformulation of Wajnryb’s classical presentation [10] in a form that is more convenient for our purposes (see Theorem 1.3 and Remark 1.1 of [6]):

Theorem 7 (Matsumoto) *For $g \geq 2$, the mapping class group $\text{Map}_{g,1}$ is generated by the Dehn twists a_0, \dots, a_{2g} along the loops c_0, \dots, c_{2g} represented in Figure 1, with the relations:*

- (i) $a_i a_j = a_j a_i$ if $c_i \cap c_j = \emptyset$, and $a_i a_j a_i = a_j a_i a_j$ if $c_i \cap c_j \neq \emptyset$;
- (ii) $(a_0 a_2 a_3 a_4)^{10} = (a_0 a_1 a_2 a_3 a_4)^6$;
- (iii) for $g \geq 3$: $(a_0 a_1 a_2 a_3 a_4 a_5 a_6)^9 = (a_0 a_2 a_3 a_4 a_5 a_6)^{12}$.




Figure 1: The Dehn–Lickorish–Humphries generators of $\text{Map}_{g,1}$

The relations (i) are the *braid relations*, and realize $\text{Map}_{g,1}$ as a quotient of an Artin group, while (ii) is a reformulation of the *chain relation*, and (iii) is a reformulation of the *lantern relation* (see [6]).

The subgroup of $\text{Map}_{g,1}$ generated by a_1, \dots, a_{2g} is the *hyperelliptic* subgroup, and is closely related to the braid group B_{2g+1} (realizing the genus g surface as a double cover of the disc branched in $2g+1$ points, the Dehn twists a_1, \dots, a_{2g} are the lifts of the standard generators of B_{2g+1}).

Lemma 8 *For every integer $1 < n < 2g$, let*

$$R_n = (a_{n+1} \cdot a_{n+2} \cdot \dots \cdot a_{2g})^{2g-n+1} \cdot \prod_{i=n}^1 (a_i \cdot a_{i+1} \cdot \dots \cdot a_{i+2g-n}) \cdot \prod_{i=2g-n+1}^1 (a_i \cdot a_{i+1} \cdot \dots \cdot a_{i+n-1}).$$

Then $(a_1 \cdot \dots \cdot a_{n-1})^{2n} \cdot (R_n)^2$ is a factorization of T_δ in $\text{Map}_{g,1}$.

Proof We work in the braid group B_{2g+1} with generators x_1, \dots, x_{2g} , and consider the expression obtained from R_n after replacing each a_i by x_i . Then it is easy to see that $a' = (x_1 \dots x_{n-1})^n$ is the full twist rotating the n leftmost strands by 2π , while $a'' = (x_{n+1} \dots x_{2g})^{2g-n+1}$ is the full twist rotating the $2g+1-n$ rightmost strands by 2π . Moreover, $b' = \prod_{i=n}^1 (x_i \dots x_{i+2g-n})$ is the braid which exchanges the n leftmost strands with the $2g-n+1$ rightmost strands in the counterclockwise direction, while $b'' = \prod_{i=2g-n+1}^1 (x_i \dots x_{i+n-1})$ does the same for the $2g-n+1$ leftmost strands and the n rightmost strands. Hence the product $b'b''$ corresponds to a full rotation of the n leftmost strands around the $2g-n+1$ rightmost strands, and $a'a''b'b''$ is the full twist $\Delta^2 = (x_1 \dots x_{2g})^{2g+1}$. Since Δ^2 is a central element in B_{2g+1} , we also have $a''b'b''a' = \Delta^2$.

We now lift things to the double cover; since Δ^2 lifts to the hyperelliptic element H (rotating the surface about its central axis by π), we deduce from the above calculation that $(a_1 \cdot \dots \cdot a_{n-1})^n \cdot R_n$ and $R_n \cdot (a_1 \cdot \dots \cdot a_{n-1})^n$ are factorizations of H , and hence that $(a_1 \cdot \dots \cdot a_{n-1})^n \cdot (R_n)^2 \cdot (a_1 \cdot \dots \cdot a_{n-1})^n$ is a factorization of $H^2 = T_\delta$. Since T_δ is central in $\text{Map}_{g,1}$, the result follows by Lemma 6(a). \square

It is in fact not hard to check explicitly that the factorization considered in Lemma 8 is Hurwitz equivalent to the standard hyperelliptic factorization $(a_1 \cdot \dots \cdot a_{2g})^{4g+2}$.

From now on we assume that $g \geq 3$. By Theorem 1.4 of [6], $(a_0 a_2 a_3 a_4)^{10} = (a_0 a_1 a_2 a_3 a_4)^6 = (a_1 a_2 a_3 a_4)^{10}$ and $(a_0 a_1 a_2 a_3 a_4 a_5 a_6)^9 = (a_0 a_2 a_3 a_4 a_5 a_6)^{12} =$

$(a_1 a_2 a_3 a_4 a_5 a_6)^{14}$. Hence, we can define new factorizations of T_δ by substitution into the factorization of Lemma 8:

Definition 9 Let $\mathcal{A} = (a_0 \cdot a_2 \cdot a_3 \cdot a_4)^{10} \cdot (R_5)^2$, $\mathcal{B} = (a_0 \cdot a_1 \cdot a_2 \cdot a_3 \cdot a_4)^6 \cdot (R_5)^2$, $\mathcal{C} = (a_0 \cdot a_1 \cdot a_2 \cdot a_3 \cdot a_4 \cdot a_5 \cdot a_6)^9 \cdot (R_7)^2$, $\mathcal{D} = (a_0 \cdot a_2 \cdot a_3 \cdot a_4 \cdot a_5 \cdot a_6)^{12} \cdot (R_7)^2$ (where for $g = 3$ we take R_7 to be the empty factorization), and $\mathcal{F}_0 = \mathcal{A} \cdot \mathcal{B} \cdot \mathcal{C} \cdot \mathcal{D}$.

$\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ are factorizations of the central element T_δ in which every factor is one of the $(a_i)_{0 \leq i \leq 2g}$, and every generator appears at least once (except possibly for \mathcal{D} , which does not involve a_1 when $g = 3$).

We also define $f_g^0, f_g^A, f_g^B, f_g^C, f_g^D$ to be the Lefschetz fibrations with monodromy factorizations $\mathcal{F}_0, \mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ respectively (so $f_g^A, f_g^B, f_g^C, f_g^D$ are irreducible and admit sections of square -1 , while f_g^0 is their fiber sum and admits a section of square -4). Let us mention that, as a consequence of Lemma 6(b), when performing a fiber sum with f_g^0 the choice of the identification diffeomorphism between fibers is irrelevant, and all possible ways in which the fiber sum can be carried out are equivalent.

The factorizations $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$ form a “universal” set of positive factorizations, in the sense that their factors are exactly the generators of $\text{Map}_{g,1}$ (out of sequence, and with some repetitions), and every relation in the presentation of Theorem 7 can be interpreted either as a Hurwitz equivalence or as a substitution replacing one of these factorizations by another one of them. We will see below that these properties are the key ingredients for the proof of Theorem 2; since many other groups related to braid groups or mapping class groups can be presented in a similar manner, the methods used here may also be relevant to the study of factorizations in these groups.

4 Stable equivalence of factorizations

In this section, we prove the following result, which implies Theorem 4:

Theorem 10 *Let F, F' be two factorizations of the same element of $\text{Map}_{g,1}$ as a product of positive Dehn twists along non-separating curves. Then there exist integers a, b, c, d, k, l such that $F \cdot (\mathcal{A})^a \cdot (\mathcal{B})^b \cdot (\mathcal{C})^c \cdot (\mathcal{D})^d \sim F' \cdot (\mathcal{A})^{a+l} \cdot (\mathcal{B})^{b-l} \cdot (\mathcal{C})^{c+k} \cdot (\mathcal{D})^{d-k}$.*

In order to prove this result, we consider factorizations where the factors are either positive Dehn twists or their inverses, and the equivalence relation \equiv generated by the following moves:

- Hurwitz moves involving only positive Dehn twists;
- creation or cancellation of pairs of inverse factors: $a_i \cdot a_i^{-1} \equiv a_i^{-1} \cdot a_i \equiv \emptyset$;
- defining relations of the mapping class group: $a_i \cdot a_j \equiv a_j \cdot a_i$ if $c_i \cap c_j = \emptyset$, $a_i \cdot a_j \cdot a_i \equiv a_j \cdot a_i \cdot a_j$ if $c_i \cap c_j \neq \emptyset$, $(a_0 \cdot a_2 \cdot a_3 \cdot a_4)^{10} \equiv (a_0 \cdot a_1 \cdot a_2 \cdot a_3 \cdot a_4)^6$, and $(a_0 \cdot a_1 \cdot a_2 \cdot a_3 \cdot a_4 \cdot a_5 \cdot a_6)^9 \equiv (a_0 \cdot a_2 \cdot a_3 \cdot a_4 \cdot a_5 \cdot a_6)^{12}$.

Lemma 11 *If the factors of F are Dehn twists along non-separating curves, then there exists a factorization \bar{F} in which every factor is of the form $a_i^{\pm 1}$ for some $0 \leq i \leq 2g$, and such that $F \equiv \bar{F}$.*

Proof We use pair creations and Hurwitz moves to replace every factor in F by a factorization involving only the $a_i^{\pm 1}$. Let τ be a factor in F . Since τ is a Dehn twist along a non-separating curve, there exist $g_0, \dots, g_k \in \{a_0^{\pm 1}, \dots, a_{2g}^{\pm 1}\}$ such that $\tau = (\prod_1^k g_j)^{-1} g_0 (\prod_1^k g_j)$. We proceed by induction on k . If $k = 0$ then τ is already one of the generators. Otherwise, if g_k is one of the generators, say a_i , then we can write $\tau = (a_i^{-1} \tilde{\tau} a_i) \equiv a_i^{-1} \cdot a_i \cdot (a_i^{-1} \tilde{\tau} a_i) \equiv a_i^{-1} \cdot \tilde{\tau} \cdot a_i$ (using a pair creation and a Hurwitz move). Similarly, if $g_k = a_i^{-1}$, then we can write $\tau = (a_i \tilde{\tau} a_i^{-1}) \equiv (a_i \tilde{\tau} a_i^{-1}) \cdot a_i \cdot a_i^{-1} \equiv a_i \cdot \tilde{\tau} \cdot a_i^{-1}$. Since $\tilde{\tau}$ is conjugated to one of the generators by a word of length $k - 1$, this completes the proof. \square

Lemma 12 *Under the assumptions of Theorem 10, $F \equiv F'$.*

Proof We first use Lemma 11 to replace F and F' by equivalent factorizations \bar{F} and \bar{F}' whose factors are all of the form $a_i^{\pm 1}$. Next, recall that if a group G admits a presentation with generators $\{a_i, i \in I\}$ and relations $\{r_j, j \in J\}$, then it is generated as a monoid by the elements $\{a_i, a_i^{-1}, i \in I\}$, and a presentation of G as a monoid is given by the set of relations $R' = \{r_j, j \in J\} \cup \{a_i a_i^{-1} = 1, a_i^{-1} a_i = 1, i \in I\}$. Hence, if \bar{F} and \bar{F}' are factorizations of the same element, then we can rewrite one into the other by successively applying the rewriting rules given by the set of relations R' . However, in the case of the mapping class group, each rewriting is one of the moves that generate the equivalence relation \equiv (either one of the defining relations of $\text{Map}_{g,1}$, or the creation or cancellation of a pair of inverses). \square

Denote by \equiv^+ the equivalence relation generated by Hurwitz moves and by the defining relations, ie, without allowing creations of pairs of inverse factors. Then we have:

Lemma 13 *Under the assumptions of Theorem 10, there exists an integer n such that $F \cdot (\mathcal{A})^n \equiv^+ F' \cdot (\mathcal{A})^n$.*

Proof By Lemma 12, $F \equiv F'$, so we can transform F into F' by a sequence of Hurwitz moves, pair creations/cancellations, and defining relations. Call $F = F_0, F_1, \dots, F_m = F'$ the successive factorizations appearing in this sequence of moves; let n_j be the number of factors of the form a_i^{-1} appearing in the factorization F_j , and let $n = \sup\{n_0, \dots, n_m\}$.

Recall that the factors of \mathcal{A} generate $\text{Map}_{g,1}$; therefore, by Lemma 6(a), for every $i \in \{0, \dots, 2g\}$ there exists a factorization \mathcal{A}_i whose factors are elements of $\{a_0, \dots, a_{2g}\}$, and such that $\mathcal{A} \sim a_i \cdot \mathcal{A}_i \sim \mathcal{A}_i \cdot a_i$. (For example \mathcal{A}_i can be obtained by cyclically permuting the factors of \mathcal{A} and deleting an occurrence of a_i). Let F_j^+ be the factorization obtained from F_j by replacing each factor of the form a_i^{-1} by the factorization \mathcal{A}_i . Then we claim that, for all $0 \leq j < m$, $F_j^+ \cdot (\mathcal{A})^{n-n_j} \equiv^+ F_{j+1}^+ \cdot (\mathcal{A})^{n-n_{j+1}}$.

Indeed, if F_{j+1} is obtained from F_j by a Hurwitz move or by applying a defining relation, then the negative factors are not involved and the claim is obvious. If F_{j+1} is obtained from F_j by deleting a pair of mutually inverse factors $a_i \cdot a_i^{-1}$, F_{j+1}^+ is obtained from F_j^+ by deleting an occurrence of the subword $a_i \cdot \mathcal{A}_i$. Hence, we can write $F_j^+ = F_j' \cdot a_i \cdot \mathcal{A}_i \cdot F_j''$ and $F_{j+1}^+ = F_j' \cdot F_j''$ for some F_j', F_j'' , and the claim follows from the sequence of Hurwitz moves

$$F_j' \cdot a_i \cdot \mathcal{A}_i \cdot F_j'' \cdot (\mathcal{A})^{n-n_j} \sim F_j' \cdot \mathcal{A} \cdot F_j'' \cdot (\mathcal{A})^{n-n_j} \sim F_j' \cdot F_j'' \cdot (\mathcal{A})^{n-n_{j+1}},$$

where in the last step we have used Lemma 6(c). The argument is the same for creations of pairs of inverses. The proof is then completed by observing that $F_0^+ = F$ and $F_m^+ = F'$, since F and F' contain no negative factors. \square

We can now proceed with the proof of Theorem 10. By Lemma 13, there exists n such that $F \cdot (\mathcal{A})^n \equiv^+ F' \cdot (\mathcal{A})^n$, so we can transform $F \cdot (\mathcal{A})^n$ into $F' \cdot (\mathcal{A})^n$ by a sequence of Hurwitz moves and applications of the defining relations. Let $F_0 = F \cdot (\mathcal{A})^n, F_1, \dots, F_m = F' \cdot (\mathcal{A})^n$ be the successive factorizations appearing in this sequence of moves. If F_{j+1} is obtained from F_j by a Hurwitz move, or by applying one of the braid relations, then we have $F_{j+1} \sim F_j$. For example, a braid relation of the form $a_i \cdot a_j \cdot a_i \equiv a_j \cdot a_i \cdot a_j$ can be viewed as a succession of two Hurwitz moves $a_i \cdot a_j \cdot a_i \sim a_j \cdot (a_i)_{a_j} \cdot a_i \sim a_j \cdot a_i \cdot (a_i)_{a_j a_i}$, where $(a_i)_{a_j a_i} = (a_j a_i)^{-1} a_i a_j a_i = a_j$.

On the other hand, if F_{j+1} is obtained from F_j by applying the relation (ii) from Theorem 7, then we can write $F_j = F_j' \cdot (a_0 \cdot a_2 \cdot a_3 \cdot a_4)^{10} \cdot F_j''$ for some

F'_j, F''_j , and $F_{j+1} = F'_j \cdot (a_0 \cdot a_1 \cdot a_2 \cdot a_3 \cdot a_4)^6 \cdot F''_j$. It is then easy to check, using Lemma 6 (a) and (c), that $F_j \cdot \mathcal{B} \sim F_{j+1} \cdot \mathcal{A}$; and vice-versa if we apply relation (ii) backwards. Similarly, if F_{j+1} is obtained from F_j by applying relation (iii), then $F_j \cdot \mathcal{D} \sim F_{j+1} \cdot \mathcal{C}$, and vice-versa if we apply relation (iii) backwards.

Hence, if we concatenate each F_j with suitable numbers of copies of \mathcal{A} , \mathcal{B} , \mathcal{C} and \mathcal{D} (depending on j), then we can realize each step as a Hurwitz equivalence. Since we always trade a copy of \mathcal{A} for a copy of \mathcal{B} , and a copy of \mathcal{C} for a copy of \mathcal{D} , Theorem 10 follows.

We can now prove Theorem 4:

Proof of Theorem 4 Let F be a factorization in $\text{Map}_{g,1}$ associated to the Lefschetz fibration f : then the product of the factors in F is equal to T_δ^m , for some integer $m \geq 1$ (such that the chosen section of f has self-intersection $-m$). The result then follows by applying Theorem 10 to F and $F' = (\mathcal{A})^m$. \square

5 Proof of Theorem 2

Let F and F' be factorizations in $\text{Map}_{g,1}$ describing the monodromies of the Lefschetz fibrations f and f' . Assumption (ii) on the self-intersection numbers of the distinguished sections implies that the products of the factors in F and F' are equal to each other, and are of the form T_δ^m for some $m \geq 1$. We first deal with the reducible singular fibers, using the following lemma:

Lemma 14 *If τ, τ' are Dehn twists along separating curves of the same type, then there exists an integer n and a factorization F'' involving only Dehn twists along non-separating curves, such that $\tau \cdot (\mathcal{A})^n \sim \tau' \cdot F''$.*

Proof τ, τ' are conjugated to each other in $\text{Map}_{g,1}$, so there exist $g_1, \dots, g_k \in \{a_0^{\pm 1}, \dots, a_{2g}^{\pm 1}\}$ such that $\tau' = (\prod_1^k g_j)^{-1} \tau (\prod_1^k g_j)$. It is enough to consider the case $k = 1$ (iterating k times in the general case). If $\tau' = a_i^{-1} \tau a_i$ then, with the same notations as in the proof of Lemma 13, we have $\tau \cdot \mathcal{A} \sim \tau \cdot a_i \cdot \mathcal{A}_i \sim a_i \cdot \tau' \cdot \mathcal{A}_i \sim \tau' \cdot (a_i)_{\tau'} \cdot \mathcal{A}_i$, and the result follows by setting $F'' = (a_i)_{\tau'} \cdot \mathcal{A}_i$. Similarly, if $\tau' = a_i \tau a_i^{-1}$ then $\tau \cdot \mathcal{A} \sim \mathcal{A} \cdot \tau \sim \mathcal{A}_i \cdot a_i \cdot \tau \sim \mathcal{A}_i \cdot \tau' \cdot a_i \sim \tau' \cdot (\mathcal{A}_i)_{\tau'} \cdot a_i$. \square

The manner in which we use this lemma is the following: let s be the number of reducible singular fibers of f and f' . Without loss of generality, we can assume that the homologically trivial vanishing cycles correspond to the first

s factors of F and F' , and that they are ordered according to types (this can always be ensured by performing Hurwitz moves). Call these factors τ_1, \dots, τ_s for F , and τ'_1, \dots, τ'_s for F' . Then assumption (iii) on the numbers of reducible singular fibers implies that τ_j and τ'_j are conjugated for each $1 \leq j \leq s$. Hence, applying Lemma 14 to each pair (τ_j, τ'_j) , and adding sufficiently many copies of \mathcal{A} to F (using Lemma 6(c) to move them to the beginning of the factorization), we can replace each τ_j by τ'_j , at the expense of generating extra Dehn twists along nonseparating curves. After suitable Hurwitz moves, we conclude that there exists an integer N and factorizations \tilde{F}, \tilde{F}' involving only Dehn twists along non-separating curves, such that $F \cdot (\mathcal{A})^N \sim \tau'_1 \cdot \dots \cdot \tau'_s \cdot \tilde{F}$ and $F' \cdot (\mathcal{A})^N \sim \tau'_1 \cdot \dots \cdot \tau'_s \cdot \tilde{F}'$.

Since \tilde{F} and \tilde{F}' are factorizations of the same element $(\tau'_1 \dots \tau'_s)^{-1} T_\delta^{m+N}$, we can apply Theorem 10 to them. It follows that there exist integers a, b, c, d, k, l such that $F \cdot (\mathcal{A})^{N+a} \cdot (\mathcal{B})^b \cdot (\mathcal{C})^c \cdot (\mathcal{D})^d \sim F' \cdot (\mathcal{A})^{N+a+l} \cdot (\mathcal{B})^{b-l} \cdot (\mathcal{C})^{c+k} \cdot (\mathcal{D})^{d-k}$. This implies that the fiber sums $\hat{f} = f \# (N+a)f_g^A \# bf_g^B \# cf_g^C \# df_g^D$ and $\hat{f}' = f' \# (N+a+l)f_g^A \# (b-l)f_g^B \# (c+k)f_g^C \# (d-k)f_g^D$ are isomorphic. Performing additional fiber sums if necessary, we can assume that $N+a = b = c = d$. Then, in order to complete the proof of Theorem 2, it is sufficient to prove that $k = l = 0$. For this purpose we use the following lemmas to compare the Euler–Poincaré characteristics and signatures of the total spaces \hat{M} and \hat{M}' of \hat{f} and \hat{f}' :

Lemma 15 $\chi(\hat{M}') - \chi(\hat{M}) = \chi(M') - \chi(M) + 10l - 9k$.

Proof Recall that the Euler characteristic of a genus g Lefschetz fibration over S^2 with r singular fibers is equal to $4 - 4g + r$. Hence, we just have to compare the numbers of singular fibers of \hat{f} and \hat{f}' . Since f_g^A has 10 more singular fibers than f_g^B , and f_g^C has 9 fewer singular fibers than f_g^D , the result follows. □

Lemma 16 $\sigma(\hat{M}') - \sigma(\hat{M}) = \sigma(M') - \sigma(M) - 6l + 5k$.

Proof By Novikov additivity, it is sufficient to show that the signatures of the total spaces M_A, M_B, M_C, M_D of $f_g^A, f_g^B, f_g^C, f_g^D$ satisfy the relations $\sigma(M_A) = \sigma(M_B) - 6$ and $\sigma(M_C) = \sigma(M_D) + 5$.

These signatures can be computed explicitly via an algorithm due to Ozbagci [8]. Since Ozbagci’s formula is a sum of individual contributions which each depend only on one of the factors and on the *product* of all the preceding factors, it

is sufficient to carry out the algorithm for the portions of \mathcal{A} and \mathcal{B} (resp. \mathcal{C} and \mathcal{D}) which differ from each other; the contributions from the common part $(R_5)^2$ (resp. $(R_7)^2$) will be the same in both cases.

In fact, after a closer look at the signature formula it is easy to convince oneself that $\sigma(M_A) - \sigma(M_B)$ and $\sigma(M_C) - \sigma(M_D)$ do not depend on g , and can be computed for a fixed low value of g (e.g., $g = 3$).

Then, rather than Ozbagci's somewhat complicated formula, one can use the following simple recipe to determine the signature – the underlying principle being that, given a Lefschetz fibration $f: M \rightarrow S^2$ admitting a section, the complement to the fiber and section classes in $H_2(M, \mathbb{Z})$ is generated by certain linear combinations of the Lefschetz thimbles of f .

Given the set of vanishing cycles $(\delta_1, \dots, \delta_r)$ (ie, loops in the fiber Σ_g such that each monodromy factor τ_i is the Dehn twist along δ_i), form the $r \times r$ matrix Q whose entries are given by

$$q_{ij} = \begin{cases} 0 & \text{if } i > j, \\ -1 & \text{if } i = j, \\ \delta_i \cdot \delta_j & \text{if } i < j, \end{cases}$$

where $\delta_i \cdot \delta_j$ is the intersection number in $H_1(\Sigma_g, \mathbb{Z})$. In a suitable sense, Q is the matrix of the intersection pairing on the space of formal linear combinations of Lefschetz thimbles, and its antisymmetrization $A = Q - Q^t$ describes the intersection pairing between vanishing cycles inside Σ_g .

Viewing Q and A as bilinear forms, the kernel of A is the space of all combinations of Lefschetz thimbles which have homologically trivial boundary in $H_1(\Sigma_g, \mathbb{Z})$, and can hence be completed to 2-cycles inside M . The restriction $Q' = Q|_{\text{Ker } A}$ is now a (degenerate) symmetric bilinear form, of rank $b_2(M) - 2$; and Q' has the same signature as the intersection form on $H_2(M, \mathbb{Z})$, ie $\sigma(Q') = \sigma(M)$.

Applying this formula, we easily check that for $g = 3$, $\sigma(M_A) = -48$, $\sigma(M_B) = -42$, $\sigma(M_C) = -35$, and $\sigma(M_D) = -40$. \square

The proof of Theorem 2 can now be completed by observing that, since $\chi(M') = \chi(M)$ and $\sigma(M') = \sigma(M)$ by assumption (i), and since \hat{M} and \hat{M}' are diffeomorphic by construction, we must have $10l = 9k$ and $6l = 5k$, which implies that $k = l = 0$.

References

- [1] **D Auroux**, *Fiber sums of genus 2 Lefschetz fibrations*, Turkish J. Math. 27 (2003) 1–10, [arXiv:math.GT/0204285](#) MathReview
- [2] **S K Donaldson**, *Lefschetz pencils on symplectic manifolds*, J. Differential Geom. 53 (1999) 205–236 MathReview
- [3] **H Endo**, *Meyer’s signature cocycle and hyperelliptic fibrations*, Math. Ann. 316 (2000) 237–257 MathReview
- [4] **A Kas**, *On the handlebody decomposition associated to a Lefschetz fibration*, Pacific J. Math. 89 (1980) 89–104 MathReview
- [5] **V Kharlamov**, **V Kulikov**, *On braid monodromy factorizations*, Izv. Ross. Akad. Nauk Ser. Mat. 67 (2003) 79–118; translation: Izvestia Math. 67 (2003) 499–534, [arXiv:math.AG/0302113](#) MathReview
- [6] **M Matsumoto**, *A presentation of mapping class groups in terms of Artin groups and geometric monodromy of singularities*, Math. Ann. 316 (2000) 401–418 MathReview
- [7] **B Moishezon**, *Complex surfaces and connected sums of complex projective planes*, Lecture Notes in Math. 603, Springer–Verlag (1977) MathReview
- [8] **B Ozbagci**, *Signatures of Lefschetz fibrations*, Pacific J. Math. 202 (2002) 99–118 MathReview
- [9] **B Siebert**, **G Tian**, *On the holomorphicity of genus two Lefschetz fibrations*, Ann. Math. to appear, [arXiv:math.SG/0305343](#)
- [10] **B Wajnryb**, *A simple presentation for the mapping class group of an orientable surface*, Israel J. Math. 45 (1983) 157–174; errata: **J S Birman**, **B Wajnryb**, Israel J. Math. 88 (1994) 425–427 MathReview