

Solutions to sheet 6


Solution to exercise 1:

- (a) Let M be the Möbius strip obtained by a suitable identification of two opposite sides of the unit square $[0, 1]^2$. We can identify the boundary ∂M with S^1 . Consider the map $\varphi_0: S^1 \times [0, \frac{1}{2}] \rightarrow M$ which sends (p, t) to the point which is obtained by starting from the boundary point $p \in S^1 = \partial M$ and going distance t orthogonally into the band. This map is continuous and surjective. It is not injective, since $\varphi_0(p, \frac{1}{2}) = \varphi_0(-p, \frac{1}{2})$ for all p . This means however, that the map φ_0 descends to the quotient

$$C := (S^1 \times [0, \frac{1}{2}]) / ((p, \frac{1}{2}) \sim (-p, \frac{1}{2}))$$

which is a crosscap. So we have a map $\varphi: C \rightarrow M$. This map is continuous and bijective, hence a homeomorphism (since C is compact and M is Hausdorff).

- (b) The following square yields the Klein bottle when the indicated glueings are performed. By cutting and identifying the two horizontal lines as indicated, the two (!) depicted regions are then Möbius strips with common boundary.



Solution to exercise 2:

- (a) Let X be a sphere with a twisted handle attached and let Y be a sphere with two crosscaps attached. We show that there is a homeomorphism $\varphi: X \rightarrow Y$. For an open disk $D \subset X$ we then have an induced homeomorphism $X \setminus D \rightarrow Y \setminus \varphi(D)$, where $X \setminus D$ and $Y \setminus \varphi(D)$ are exactly the spaces given in the problem. (We are using that a sphere minus a disk is another disk, and that the fact that it is irrelevant where in a surface one removes an open disk.) The space X is nothing but a Klein bottle, since a twisted handle is a Klein bottle minus a disk. To obtain the space Y we first remove two disks, which yields a cylinder. Glueing a crosscap on each of the boundary components of the cylinder is (by problem 1) the same as glueing two Möbius strips on the cylinder. This in turn is the same as glueing two Möbius strips along their boundary, which (again by problem 1) yields a Klein bottle. Hence X and Y are both Klein bottles and therefore there is a homeomorphism $\varphi: X \rightarrow Y$.

- (b) A similar discussion can be made.

Solution to exercise 3:

- (a) Take an annulus surrounding each disc, and define a map which is the identity on the rest of the disc, but compresses the annulus radially by a factor of 2 towards the outside. This map shrinks the ‘disc with holes’ into itself. Simultaneously, chop the cylinder into three segments, and map the outer ones to the ‘missing’ parts of the annuli (the inner halves) and stretch the middle segment to cover the whole cylinder now.
- (b) In this case X is a disc with a twisted handle attached.

Solution to exercise 4:

(This is Lemma 5.4.6. in the Knotes of J. Roberts) The map is defined by re-identifying the edges in F' which we just ‘unidentified’. It is a quotient map and therefore continuous. One can check that restricted to the boundary of F' , the map p is a 2 : 1 covering map onto C . But there are only two double covers of the circle, the connected one and the disconnected one.

Solution to exercise 5:

- (a) This follows from the statement in the previous problem.
- (b) This follows from the fact that a one-sided curve has a neighborhood which is homeomorphic to a Möbius strip, together with the observation that a Möbius strip does not disconnect when cut a long its middle circle.

Solution to exercise 6:

- (a) We need to make the problem more precise: We need to assume that X is a combinatorial surface which we can write as a combinatorial union of sub-surfaces $X = A \cup B$. To compute $\chi(X) = \chi(A \cup B)$ we note that

$$\begin{aligned} & \#\{\text{vertices in } X\} \\ &= \#\{\text{vertices in } A\} + \#\{\text{vertices in } B\} - \#\{\text{vertices in } A \cap B\} \end{aligned}$$

and the same holds for the respective numbers of edges and triangles, hence

$$\chi(X) = \chi(A) + \chi(B) - \chi(A \cap B).$$


- (b) A combinatorial circle C is nothing but a polygonal loop. If C has n vertices then C has n edges, hence $\chi(C) = n - n = 0$.

Solution to exercise 7:

- (a) From problem 4 we know that the boundary of F' is the boundary of F plus one or two combinatorial circles. By the previous problem these circles do not contribute to the Euler characteristic.
- (b) The Euler characteristic of a combinatorial disc is equal to 1. There are 1 or 2 new boundary components, if we cap them off with discs the Euler characteristic increases by 1 or 2 respectively, which follows from Problem 6a.

Solution to exercise 8:

- (a) Start with an arrangement X of d disjoint discs. We have $\chi(X) = d$. Now let X' be the space obtained by adding a combinatorial band (a rectangle subdivided diagonally into two triangles) as in this picture:




We have 2 new triangles and 3 new edges. Hence we have $\chi(X') = \chi(X) - 1$. So if Y is obtained by adding b bands to X then $\chi(Y) = d - b$.

- (b) This surface S is non-orientable, it has one boundary component and Euler characteristic -1 , since a cylinder has $\chi = 0$ and adding a band decreased χ by 1 as we have seen above. The classification of surfaces tells us that $-1 = \chi(S) = 2 - g(S) - 1$, which implies that the genus $g(S)$ is equal to 2. Hence S is a Klein bottle with a disc removed.

Solution to exercise 9:

A triangulation of the sphere can be obtained, for example, by radially projecting a tetrahedron T whose vertices lie on the sphere. The fact that any two triangulations yield the same Euler characteristic follows from these two facts:

- (i) The Euler characteristic of a triangulation does not change when we refine the triangulation:
- (ii) Any two triangulations (of the sphere) have a common refinement, e.g.



Solution to exercise 10:


This follows from previous problems: Start with $F_1 \sqcup F_2$ and remove a disc in both surfaces to obtain $F'_1 \sqcup F'_2$. Removing a disc decreases χ by 1, so we have $\chi(F'_1 \sqcup F'_2) = \chi(F_1 \sqcup F_2) - 2 = \chi(F_1) + \chi(F_2) - 2$. Now glueing F'_1 with F'_2 along the new boundary components doesn't change χ by Problem 7a, hence $\chi(F_1 \# F_2) = \chi(F_1) + \chi(F_2) - 2$. The Euler characteristic and the genus of an *orientable* surface F are linked via $\chi(F) = 2 - 2g(F)$. By what we just proved we have

$$\begin{aligned} 2 - 2g(F_1 \# F_2) &= \chi(F_1 \# F_2) \\ &= \chi(F_1) + \chi(F_2) - 2 \\ &= 2 - 2g(F_1) + 2 - 2g(F_2) - 2 \\ &= 2 - 2(g(F_1) + g(F_2)) \end{aligned}$$


and hence, $g(F_1 \# F_2) = g(F_1) + g(F_2)$.

Solution to exercise 11:

- (a) This glueing yields a cone, which, as a topological surface, is a disc.
- (b) The simplest model is a bigon:




- (c) In the following we take two copies of the above planar model for Σ_1 . A disc can be removed in the surface by adding an additional edge to the polygon. I.e. we start with




and

Then we glue the two new edges, which means that we glue the surfaces

along the boundaries of the removed discs. We get:




This can also be seen as this octagon:



(d) Convince yourself by cutting it out and glueing the corresponding edges.

(e) The model looks as follows:



Solution to exercise 12:

- (a) Proceed as in the previous exercise sheets to deform the boundary curve into a diagram of the trefoil knot.
- (b) This is because an untwisted band and a band twisted by 360 degrees are homeomorphic via a homeomorphism that is the identity on the vertical boundary segments. In other words, the twisting of the bands is a feature of the given embedding, not of the surface itself.

- (c) An ambient isotopy between the two surfaces (i.e. between the one with twisted bands and the one with untwisted bands) induces an isotopy of the boundaries. This is impossible, since the surface with untwisted bands has the unknot as its boundary, while the initial surface has a trefoil boundary.

Solution to exercise 13:

As in problem 12b the untwisted band in the left surface is homeomorphic, via a map preserving the boundary, to the knotted and twisted ‘pretzel’ band in the right surface. A homeomorphism from the left to the right surface will be this map on the untwisted band and the identity everywhere else.

Solution to exercise 14:

These are two diagrams of the trefoil knot. The left one is easily transformed into the right one by taking the left-most string to the right. We have seen in an earlier sheet that the right one is the trefoil knot. We make a chess-board colouring and leave the outside white. Then the left surface is non-orientable, the right surface is orientable.

Solution to exercise 15:

The surface obtained by the chessboard coloring can be seen as a surface made from disks and bands. This allows us to compute the Euler characteristic (using problem 8a). Together with the orientability we can then identify the surfaces. The surface S_1 on the left has 2 disks and 5 bands, i.e. $\chi(S_1) = 2 - 5 = -3$. It is orientable and it has 5 boundary components. We have

$$-3 = \chi(S_1) = 2 - 2g(S_1) - 5$$

and hence, S_1 is the orientable surface of genus 0 with 5 boundary components, i.e. a sphere with 5 disks removed.

The surface S_2 on the right has 3 disks and 8 bands, i.e. $\chi(S_2) = 3 - 8 = -5$. It is non-orientable and it has 3 boundary components. We have

$$-5 = \chi(S_2) = 2 - g(S_2) - 3$$

and hence, S_2 is the non-orientable surface of genus 4 with 3 boundary components.

Solution to exercise 16:

This surface S is made from 4 disks and 7 bands, i.e. $\chi(S) = 4 - 7 = -3$. It is orientable and it has 3 boundary components. We have

$$-3 = \chi(S) = 2 - 2g(S) - 3$$

and hence, S is the orientable surface of genus 1 with 3 boundary components, i.e. a torus with 3 disks removed.