

Graph Theory

Instructor: Benny Sudakov

Assignment 5

To be completed by April 7

Unless noted otherwise, all graphs considered are simple. The solution of every problem should be no longer than one page.

Problem 1: Let G be a connected graph on more than 2 vertices such that every edge is contained in some perfect matching of G . Show that G is 2-edge-connected.

Problem 2:

(a) Let G be a graph on $2n$ vertices that has exactly one perfect matching. Show that G has at most n^2 edges.

[Hint: .segde gnihtam owt neewteb gnissorc segde owt tsom ta sah G taht evresbO]

(b) Construct such a G containing exactly n^2 edges.

Problem 3: Let A be a finite set with subsets A_1, \dots, A_n , and let d_1, \dots, d_n be positive integers. Show that there are disjoint subsets $D_k \subseteq A_k$ with $|D_k| = d_k$ for all $k \in [n]$ if and only if

$$|\cup_{i \in I} A_i| \geq \sum_{i \in I} d_i$$

for all $I \subseteq [n]$.

Problem 4: Suppose M is a matching in a bipartite graph $G = (A \cup B, E)$. We say that a path $P = a_1 b_1 \cdots a_k b_k$ is an *augmenting path* in G if $b_i a_{i+1} \in M$ for all $i \in [k-1]$ and a_1 and b_k are not covered by M . The name comes from the fact that the size of M can be increased by flipping the edges along P (in other words, taking the symmetric difference of M and P): by deleting the edges $b_i a_{i+1}$ from M and adding the edges $a_i b_i$ instead.

(a) Prove Hall's theorem by showing that if Hall's condition is satisfied and M does not cover A , then there is an augmenting path in G .

- (b) Show that if M is not a maximum matching (i.e. there is a larger matching in G) then the graph contains an augmenting path. Is this true for non-bipartite graphs as well?

Problem 5: Show that for $k \geq 1$, every k -regular $(k - 1)$ -edge-connected graph on an even number of vertices contains a perfect matching.

[Hint: .ytirap no desab sesac owt otni kaerB]